




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : C07D 211/52, 295/12, 211/70, 401/04, 401/06, 401/12, 223/08, 409/06, 211/58, A61K 31/445		A1	(11) International Publication Number: <b>WO 97/24325</b>  (43) International Publication Date: 10 July 1997 (10.07.97)
(21) International Application Number: PCT/JP96/03820  (22) International Filing Date: 26 December 1996 (26.12.96)  (30) Priority Data: 7/343905 28 December 1995 (28.12.95) JP 8/187375 17 July 1996 (17.07.96) JP  (71) Applicant (for all designated States except US): TAKEDA CHEMICAL INDUSTRIES, LTD. [JP/JP]; 1-1, Doshomachi 4-chome, Chuo-ku, Osaka-shi, Osaka 541 (JP).  (72) Inventors; and (75) Inventors/Applicants (for US only): KATO, Kaneyoshi [JP/JP]; 2-40, Maruyamadai 2-chome, Kawanishi-shi, Hyogo 666-01 (JP). YAMAMOTO, Mitsuo [JP/JP]; 5-20, Shikanjima 1-chome, Konohana-ku, Osaka-shi, Osaka 554 (JP). HONDA, Susumu [JP/JP]; 6-22, Izumicho, Nishinomiya-shi, Hyogo 662 (JP). FUJISAWA, Tomoyuki [JP/JP]; 18-D76-207, Tsukumodai 5-chome, Suita-shi, Osaka 565 (JP).		(74) Agents: ASAHINA, Tadao et al.; Osaka Plant of Takeda Chemical Industries, Ltd., 17-85, Jusohonmachi 2-chome, Yodogawa-ku, Osaka-shi, Osaka 532 (JP).  (81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, IL, IS, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  Published With international search report.	


(54) Title: DIPHENYLMETHANE DERIVATIVES AS MIP-1 $\alpha$ /RANTES RECEPTOR ANTAGONISTS

## (57) Abstract

An MIP-1 $\alpha$ /RANTES-receptor antagonist which comprises the compound of formula (I), wherein Ar<sup>1</sup> and Ar<sup>2</sup> independently represent an optionally substituted aromatic group; Q<sup>1</sup> and Q<sup>2</sup> independently represent an optionally substituted divalent C<sub>1-6</sub> aliphatic hydrocarbon group which may have either oxygen or sulfur within the carbon chain; R<sup>1</sup> represents hydrogen atom, an optionally substituted lower alkyl group or an optionally substituted lower alkyl-carbonyl group; R<sup>2</sup> represents an optionally substituted hydrocarbon group or an optionally substituted acyl group, or R<sup>1</sup> and R<sup>2</sup>, taken together with the adjacent nitrogen atom, form an optionally substituted nitrogen containing heterocyclic group; and a group of formula (a) represents an optionally substituted nitrogen-containing mono or fused heterocyclic group, or a salt thereof.



[ I ]



(a)

BEST AVAILABLE COPY

**BEST AVAILABLE COPY**

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

DESCRIPTION

## DIPHENYLMETHANE DERIVATIVES AS MIP-1ALPHA/RANTES RECEPTOR ANTAGONISTS

TECHNICAL FIELD

5           This invention relates to a compound which has a  
MIP-1 $\alpha$ /RANTES receptor antagonism and is useful for  
preventing or treating allergic diseases (e.g.  
bronchial asthma, atopic dermatitis, etc.),  
inflammatory diseases (e.g. arteriosclerosis,  
10       rheumatoid arthritis, etc.) and multiple sclerosis.

BACKGROUND ART

Chemokines are a group of cytokines regulating  
chemotaxis of leukocytes and it has recently been  
becoming clear that chemokines and other cytokines have  
15       relevance to the progression and exacerbation of  
conditions of diseases in the acute and chronic periods  
of inflammatories.

It is known that, among chemokines, RANTES  
(regulated on activation, normal T expressed and  
20       secreted) and MIP-1 $\alpha$  (macrophage inflammatory  
protein-1 $\alpha$ ) belong to CC chemokines and act on  
lymphocytes, monocytes, eosinophils and basophils to  
enhance migration and further show a direct leucocyte  
activation, e.g. degranulation, secretion of various  
25       inflammatory mediator, etc. (Clinical Immunotherapy  
Vol. 4, pages 1-8, 1995).

Particularly, an increase in amount of gene  
expression of RANTES is observed in synovia of  
rheumatism patients (Clinical & Experimental  
30       Immunology, Vol. 101, page 398, 1995; and Lancet, Vol.  
343, page 547, 1994) or focus of arteriosclerosis,  
which suggests that they are concerned with the  
diseases. It has also been reported that in  
administration of a MIP-1 $\alpha$  antibody to mice delays  
35       crisis of arthritis and ameliorates the symptoms (The  
Journal of American Society for Clinical Investigation,

Vol. 95, page 2868, 1995). However, antibodies are macromolecules and have a problem about oral absorption and stability.

5 MIP-1 $\alpha$ /RANTES receptor described in this specification means a mutual receptor among chemokines, for example, MIP-1 $\alpha$ , RANTES or MCP-3 (monocyte chemoattractant protein-3), etc., which is called CCR1 (Nature Medicine, page 1174, 1996).

10 According to the above background, it has been desired to develop a novel drug as a CCR1 receptor antagonist/agonist. Although a peptide antagonist for a RANTES receptor is known (Journal of Biological Chemistry, 27, 18, page 12521-10527(1996)), it has a problem about oral absorption and stability.

15 It has been becoming apparent that eosinophils and basophils are concerned in recruitment, progression and exacerbation of various allergic diseases and inflammatory diseases due to aggregation to the inflammatory site and activation. Therefore, It is  
20 considered that immunopathy diseases (e.g. bronchial asthma, atopic dermatitis, arteriosclerosis, articular rheumatism, etc.) may be prevented or treated by inhibiting the action of the above chemokines (Clinical Immunotherapy Vol. 4, pages 1-8, 1995). However, such  
25 antagonists have never been reported so far.


On the other hand, a lot of diphenylmethane derivatives have hitherto been synthesized (Journal of Medicinal Chemistry, Vol. 34, page 12, 1991; Arch. int. Pharmacodyn., Vol. 107, page 194, 1956; Japanese Patent  
30 Kokai (Laid-Open) No. 123164/1987). For example, loperamide is commercially available as antidiarrheic. It is also known that loperamide has a calmodulin antagonism but it is not known that it inhibits migration of cells induced by the chemokines. It is  
35 not known that haloperidol having a 4-hydroxypiperidyl group used as an antipsychotic agent has the action.



It has been desired to develop a MIP-1 $\alpha$ /RANTES receptor antagonist and a novel drug inhibiting diseases caused by RANTES or MIP-1 $\alpha$ .

# DISCLOSURE OF INVENTION

5 The inventors of this invention have intensively studied. As a result, it has been found that a compound of the formula:




wherein Ar<sup>1</sup> and Ar<sup>2</sup> independently represent an optionally substituted aromatic group;

15 Q<sup>1</sup> and Q<sup>2</sup> independently represent an optionally substituted divalent C<sub>1-6</sub> aliphatic hydrocarbon group which may have oxygen or sulfur within the carbon chain;


R<sup>1</sup> is a hydrogen atom, an optionally substituted lower alkyl group or an optionally substituted lower alkyl-carbonyl group;

R<sup>2</sup> is an optionally substituted hydrocarbon group or an acyl group, or R<sup>1</sup> and R<sup>2</sup>, taken together with the adjacent nitrogen atom, may form an optionally substituted nitrogen-containing heterocyclic ring; and

25 a group of the formula:



30 is an optionally substituted monocyclic or fused nitrogen-containing heterocyclic group, or a salt thereof has an excellent MIP-1 $\alpha$ /RANTES receptor antagonism, unexpectedly, on the basis of a specific chemical structure of the formula:



5           This invention has been accomplished on the basis of the above discovery.

          This invention is, therefore, directed to:

- (1) A MIP-1 $\alpha$ /RANTES receptor antagonist comprising a compound [I] or a salt thereof,  
10       (2) A composition as described in the above item (1), wherein

          Ar<sup>1</sup> and Ar<sup>2</sup> independently represent (A) a monocyclic or fused polycyclic aromatic hydrocarbon group having 6 to 14 carbon atoms, or (B) a 5- to 11-  
15       membered monocyclic or fused heteroaromatic group having at least one of 1 or 2 kinds of hetero atoms selected from nitrogen, sulfur and oxygen in addition to carbon atoms, said heterocyclic group being optionally fused with the monocyclic or fused  
20       polycyclic aromatic hydrocarbon group having 6 to 14 carbon atoms, each of which may have a substituent selected from the group consisting of  
(I) a halogen atom,  
(II) a C<sub>1-3</sub> alkylendioxy group,  
25       (III) a nitro group,  
(IV) a cyano group,  
(V) a C<sub>1-6</sub> alkyl group optionally having 1 to 3 halogen atoms,  
(VI) a C<sub>2-6</sub> alkenyl group optionally having 1 to 3  
30       halogen atoms,  
(VII) a C<sub>2-6</sub> alkynyl group optionally having 1 to 3 halogen atoms,  
(VIII) a C<sub>3-6</sub> cycloalkyl group,  
(IX) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3  
35       halogen atoms,

- (X) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms,  
(XI) a hydroxyl group,  
(XII) an amino group,  
5 (XIII) a mono-C<sub>1-6</sub> alkylamino group,  
(XIV) a di-C<sub>1-6</sub> alkylamino group,  
(XV) a 5- to 7-membered cyclic amino group,  
(XVI) an acylamino group which is shown by the formula:  
10 (i) -NHCOOR<sup>3</sup>, (ii) -NHCONHR<sup>3</sup>, (iii) -NHCOR<sup>3</sup> or (iv) -NHSO<sub>2</sub>R<sup>3</sup> wherein R<sup>3</sup> is (1) a C<sub>1-6</sub> alkyl group, (2) a C<sub>2-6</sub> alkenyl group, (3) a C<sub>2-6</sub> alkynyl group, (4) a C<sub>3-6</sub> cycloalkyl group which may be fused with a benzene ring optionally having 1 to 3 C<sub>1-6</sub> alkoxy groups, (5) a C<sub>6-10</sub> aryl group or (6) a C<sub>7-16</sub> aralkyl group, each of a group  
15 shown by above items (1) to (6) optionally having 1 to 5 substituents selected from the group consisting of (a) a halogen atom, (b) a C<sub>1-3</sub> alkylenedioxy group, (c) a nitro group, (d) a cyano group, (e) a C<sub>1-6</sub> alkyl group optionally having 1 to 3 halogen atoms, (f) a C<sub>3-6</sub> cycloalkyl group, (g) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (h) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (i) a hydroxyl group, (j) an amino group, (k) a mono-C<sub>1-6</sub> alkylamino group,  
20 group, (l) a di-C<sub>1-6</sub> alkylamino group, (m) a C<sub>1-6</sub> alkyl-carbonyl group, (n) a carboxyl group, (o) a C<sub>1-6</sub> alkoxy-carbonyl group, (p) a carbamoyl group, (q) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (r) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (s) a C<sub>6-10</sub> aryl-carbamoyl group, (t) a sulfo group,  
30 group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, (w) a C<sub>6-10</sub> aryloxy group and (x) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being  
35 optionally fused with a benzene ring,

- (XVII) a C<sub>1-6</sub> alkyl-carbonyl group,  
(XVIII) a carboxyl group,  
(XIX) a C<sub>1-6</sub> alkoxy-carbonyl group,  
(XX) a carbamoyl group,  
5 (XXI) a mono-C<sub>1-6</sub> alkyl-carbamoyl group,  
(XXII) a di-C<sub>1-6</sub> alkyl-carbamoyl group,  
(XXIII) a C<sub>6-10</sub> aryl-carbamoyl group,  
(XXIV) a sulfo group,  
(XXV) a C<sub>1-6</sub> alkylsulfonyl group,  
10 (XXVI) a C<sub>6-10</sub> aryl group, and  
(XXVII) a C<sub>6-10</sub> aryloxy group;  
Q<sup>1</sup> and Q<sup>2</sup> independently represent  
(I) a C<sub>1-6</sub> alkylene group,  
(II) a C<sub>2-6</sub> alkenylene group, or  
15 (III) a C<sub>2-6</sub> alkynylene group, each of a group shown by  
the above items (I) to (III) may have oxygen or  
optionally oxydized sulfur within the carbon chain;  
R<sup>1</sup> is  
(I) a hydrogen atom,  
20 (II) a C<sub>1-6</sub> alkyl group which may have 1 to 5  
substituents selected from the group consisting of (a)  
a halogen atom, (b) a C<sub>1-3</sub> alkylendioxy group, (c) a  
nitro group, (d) a cyano group, (e) a C<sub>1-6</sub> alkyl group  
optionally having 1 to 3 halogen atoms, (f) a C<sub>3-6</sub>  
25 cycloalkyl group, (g) a C<sub>1-6</sub> alkoxy group optionally  
having 1 to 3 halogen atoms, (h) a C<sub>1-6</sub> alkylthio group  
optionally having 1 to 3 halogen atoms, (i) a hydroxyl  
group, (j) an amino group, (k) a mono-C<sub>1-6</sub> alkylamino  
group, (l) a di-C<sub>1-6</sub> alkylamino group, (m) a C<sub>1-6</sub> alkyl-  
30 carbonyl group, (n) a carboxyl group, (o) a C<sub>1-6</sub> alkoxy-  
carbonyl group, (p) a carbamoyl group, (q) a mono-C<sub>1-6</sub>  
alkyl-carbamoyl group, (r) a di-C<sub>1-6</sub> alkyl-carbamoyl  
group, (s) a C<sub>6-10</sub> aryl-carbamoyl group, (t) a sulfo  
group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl

group, (w) a C<sub>6-10</sub> aryloxy group and (x) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, or  
(III) a C<sub>1-6</sub> alkyl-carbonyl group which may have 1 to 5 substituents selected from (a) a halogen atom, (b) a C<sub>1-3</sub> alkylendioxy group, (c) a nitro group, (d) a cyano group, (e) a C<sub>1-6</sub> alkyl group optionally having 1 to 3 halogen atoms, (f) a C<sub>3-6</sub> cycloalkyl group, (g) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (h) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (i) a hydroxyl group, (j) an amino group, (k) a mono-C<sub>1-6</sub> alkylamino group, (l) a di-C<sub>1-6</sub> alkylamino group, (m) a C<sub>1-6</sub> alkyl-carbonyl group, (n) a carboxyl group, (o) a C<sub>1-6</sub> alkoxy-carbonyl group, (p) a carbamoyl group, (q) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (r) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (s) a C<sub>6-10</sub> aryl-carbamoyl group, (t) a sulfo group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, (w) a C<sub>6-10</sub> aryloxy group and (x) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally or fused with a benzene ring;

R<sup>2</sup> is

- (I) a C<sub>1-6</sub> alkyl group,
  - (II) a C<sub>2-6</sub> alkenyl group,
  - (III) a C<sub>2-6</sub> alkynyl group,
  - (IV) a C<sub>3-6</sub> cycloalkyl group which may be fused with a benzene ring optionally having 1 to 3 C<sub>1-6</sub> alkoxy groups,
  - (V) a C<sub>6-10</sub> aryl group,
  - (VI) a C<sub>7-16</sub> aralkyl group,
- each of a group shown by above the items (1) to (6)

optionally having 1 to 5 substituents selected from the group consisting of (a) a halogen atom, (b) a C<sub>1-3</sub> alkylenedioxy group, (c) a nitro group, (d) a cyano group, (e) a C<sub>1-6</sub> alkyl group optionally having 1 to 3 halogen atoms, (f) a C<sub>3-6</sub> cycloalkyl group, (g) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (h) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (i) a hydroxyl group, (j) an amino group, (k) a mono-C<sub>1-6</sub> alkylamino group, (l) a di-C<sub>1-6</sub> alkylamino group, (m) a C<sub>1-6</sub> alkyl-carbonyl group, (n) a carboxyl group, (o) a C<sub>1-6</sub> alkoxy-carbonyl group, (p) a carbamoyl group, (q) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (r) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (s) a C<sub>6-10</sub> aryl-carbamoyl group, (t) a sulfo group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, (w) a C<sub>6-10</sub> aryloxy group and (x) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, or

(VII) an acyl group which is shown by the formula:  
-(C=O)-R<sup>4</sup>, -SO<sub>2</sub>-R<sup>4</sup>, -(C=O)NR<sup>5</sup>R<sup>4</sup>, -(C=O)O-R<sup>4</sup>, -(C=S)O-R<sup>4</sup>,  
or -(C=S)NR<sup>5</sup>R<sup>4</sup>, wherein R<sup>4</sup> is

(i) a hydrogen atom,  
(ii) a C<sub>1-6</sub> alkyl group,  
(iii) a C<sub>2-6</sub> alkenyl group,  
(iv) a C<sub>2-6</sub> alkynyl group,  
(v) a C<sub>3-6</sub> cycloalkyl group which may be fused with a benzene ring optionally having 1 to 3 C<sub>1-6</sub> alkoxy groups,  
(vi) a C<sub>6-10</sub> aryl group,  
(vii) a C<sub>7-16</sub> aralkyl group,  
(viii) a 5- to 11-membered heterocyclic group having at least one hetero atom selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said

- heterocyclic group being optionally fused with a benzene ring,
- (ix) a C<sub>1-6</sub> alkyl-carbonyl group,
- (x) a carboxyl group,
- 5 (xi) a C<sub>1-6</sub> alkoxy-carbonyl group,
- (xii) a mono-C<sub>1-6</sub> alkyl-carbamoyl group,
- (xiii) a di-C<sub>1-6</sub> alkyl-carbamoyl group,
- (xiv) a 5- to 7-membered cyclic amino group, or
- (xv) a C<sub>6-10</sub> aryloxy group,
- 10 each of a group shown by the above items (ii) to (xv) optionally having 1 to 5 substituents selected from the group consisting of (a) a halogen atom, (b) a C<sub>1-3</sub> alkylenedioxy group, (c) a nitro group, (d) a cyano group, (e) a C<sub>1-6</sub> alkyl group optionally substituted
- 15 with (e-1) a halogen atom, (e-2) a C<sub>1-3</sub> alkylenedioxy group, (e-3) a nitro group, (e-4) a cyano group, (e-5) a C<sub>3-6</sub> cycloalkyl group, (e-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (e-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms,
- 20 (e-8) a hydroxyl group, (e-9) an amino group, (e-10) a mono-C<sub>1-6</sub> alkylamino group, (e-11) a di-C<sub>1-6</sub> alkylamino group, (e-12) a C<sub>1-6</sub> alkyl-carbonyl group, (e-13) a carboxyl group, (e-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (e-15) a carbamoyl group, (e-16) a mono-C<sub>1-6</sub> alkyl-
- 25 carbamoyl group, (e-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (e-18) a C<sub>6-10</sub> aryl-carbamoyl group, (e-19) a sulfo group, (e-20) a C<sub>1-6</sub> alkylsulfonyl group, (e-21) a C<sub>6-10</sub> aryl group, (e-22) a C<sub>6-10</sub> aryloxy group or (e-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero
- 30 atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (f) a C<sub>3-6</sub> cycloalkyl group, (g) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (h) a C<sub>1-6</sub> alkylthio group

optionally having 1 to 3 halogen atoms, (i) a C<sub>7-16</sub> aralkyl group, (j) a hydroxyl group, (k) an amino group which may be substituted with a C<sub>1-6</sub> alkyl carbonyl group, (l) a mono-C<sub>1-6</sub> alkylamino group, (m) a di-C<sub>1-6</sub> alkylamino group, (n) a C<sub>1-6</sub> alkyl-carbonyl group whose alkyl portion may be substituted with (n-1) a halogen atom, (n-2) a C<sub>1-3</sub> alkylenedioxy group, (n-3) a nitro group, (n-4) a cyano group, (n-5) a C<sub>3-6</sub> cycloalkyl group, (n-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (n-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (n-8) a hydroxyl group, (n-9) an amino group, (n-10) a mono-C<sub>1-6</sub> alkylamino group, (n-11) a di-C<sub>1-6</sub> alkylamino group, (n-12) a C<sub>1-6</sub> alkyl-carbonyl group, (n-13) a carboxyl group, (n-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (n-15) a carbamoyl group, (n-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (n-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (n-18) a C<sub>6-10</sub> aryl-carbamoyl group, (n-19) a sulfo group, (n-20) a C<sub>1-6</sub> alkylsulfonyl group, (n-21) a C<sub>6-10</sub> aryl group, (n-22) a C<sub>6-10</sub> aryloxy group or (n-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (o) a carboxyl group, (p) a C<sub>1-6</sub> alkoxy-carbonyl group, (q) a formyl group which may be substituted with 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (r) a carbamoyl group, (s) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (s-1) a halogen atom, (s-2) a C<sub>1-3</sub> alkylenedioxy group, (s-3) a nitro group, (s-4) a cyano group, (s-5) a C<sub>3-6</sub> cycloalkyl group, (s-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (s-7) a



C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (s-8) a hydroxyl group, (s-9) an amino group, (s-10) a mono-C<sub>1-6</sub> alkylamino group, (s-11) a di-C<sub>1-6</sub> alkylamino group, (s-12) a C<sub>1-6</sub> alkyl-carbonyl group, (s-13) a carboxyl group, (s-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (s-15) a carbamoyl group, (s-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (s-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (s-18) a C<sub>6-10</sub> aryl-carbamoyl group, (s-19) a sulfo group, (s-20) a C<sub>1-6</sub> alkylsulfonyl group, (s-21) a C<sub>6-10</sub> aryl group, (s-22) a C<sub>6-10</sub> aryloxy group or (s-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (t) a di-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (t-1) a halogen atom, (t-2) a C<sub>1-3</sub> alkylenedioxy group, (t-3) a nitro group, (t-4) a cyano group, (t-5) a C<sub>3-6</sub> cycloalkyl group, (t-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (t-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (t-8) a hydroxyl group, (t-9) an amino group, (t-10) a mono-C<sub>1-6</sub> alkylamino group, (t-11) a di-C<sub>1-6</sub> alkylamino group, (t-12) a C<sub>1-6</sub> alkyl-carbonyl group, (t-13) a carboxyl group, (t-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (t-15) a carbamoyl group, (t-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (t-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (t-18) a C<sub>6-10</sub> aryl-carbamoyl group, (t-19) a sulfo group, (t-20) a C<sub>1-6</sub> alkylsulfonyl group, (t-21) a C<sub>6-10</sub> aryl group, (t-22) a C<sub>6-10</sub> aryloxy group or (t-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (u) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (v)

an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group, (w) a sulfo group which may be substituted with an amino group, (x) a C<sub>1-6</sub> alkylsulfonyl group, (y) a C<sub>6-10</sub> aryl group, (z) a C<sub>6-10</sub> aryloxy group, (aa) a C<sub>2-6</sub> alkenylamino group, (bb) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (cc) a 5- to 7-membered cyclic amino group which may have an oxo group or which may be substituted with a hydroxyl group, (dd) a C<sub>1-6</sub> alkoxy-carbamoyl group, (ee) a carbamoyloxy group, (ff) a sulfamoyl group, (gg) a mono-C<sub>1-6</sub> alkyl-sulfamoyl group, and (hh) a di-C<sub>1-6</sub> alkyl-sulfamoyl group;

15 R<sup>5</sup> is

(I) a hydrogen atom or

(II) a C<sub>1-6</sub> alkyl group;

or R<sup>1</sup> and R<sup>2</sup>, taken together with the adjacent nitrogen atom, form a 4- to 8-membered heterocyclic group

20 optionally having at least one nitrogen and 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, which may have 1 to 5 substituents selected from the group


25 consisting of (a) a halogen atom, (b) a C<sub>1-3</sub> alkylenedioxy group, (c) a nitro group, (d) a cyano group, (e) a C<sub>1-6</sub> alkyl group optionally having 1 to 3 halogen atoms, (f) a C<sub>3-6</sub> cycloalkyl group, (g) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms,

30 (h) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (i) a hydroxyl group, (j) an amino group, (k) a mono-C<sub>1-6</sub> alkylamino group, (l) a di-C<sub>1-6</sub> alkylamino group, (m) a C<sub>1-6</sub> alkyl-carbonyl group, (n) a carboxyl group, (o) a C<sub>1-6</sub> alkoxy-carbonyl group, (p) a

35 carbamoyl group, (q) a mono-C<sub>1-6</sub> alkyl-carbamoyl group,

(r) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (s) a C<sub>6-10</sub> aryl-carbamoyl group, (t) a sulfo group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, and (w) a C<sub>6-10</sub> aryloxy group;

5 a group of the formula:



is (1) a 4- to 9-membered monocyclic ring or (2) 6- to  
 10 14-membered bicyclic ring, each of which may have 1 or  
 2 unsaturated bonds and optionally having 1 or 2  
 substituents selected from the group consisting of  
 (i) a C<sub>1-6</sub> alkyl group,  
 (ii) a C<sub>1-6</sub> alkoxy group,  
 15 (iii) a C<sub>1-6</sub> alkylthio group, each of a group shown by  
 the above items (i) to (iii) may have 1 to 5  
 substituents selected from (a) a halogen atom, (b) a  
 C<sub>1-3</sub> alkylenedioxy group, (c) a nitro group, (d) a cyano  
 group, (e) a C<sub>1-6</sub> alkyl group optionally having 1 to 3  
 20 halogen atoms, (f) a C<sub>3-6</sub> cycloalkyl group, (g) a C<sub>1-6</sub>  
 alkoxy group optionally having 1 to 3 halogen atoms,  
 (h) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3  
 halogen atoms, (i) a hydroxyl group, (j) an amino  
 group, (k) a mono-C<sub>1-6</sub> alkylamino group, (l) a di-C<sub>1-6</sub>  
 25 alkylamino group, (m) a C<sub>1-6</sub> alkyl-carbonyl group, (n) a  
 carboxyl group, (o) a C<sub>1-6</sub> alkyl-carbamoyl group, (p) a  
 carbamoyl group, (q) a mono-C<sub>1-6</sub> alkyl-carbamoyl group,  
 (r) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (s) a C<sub>6-10</sub> aryl-  
 carbamoyl group, (t) a sulfo group, (u) a C<sub>1-6</sub>  
 30 alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, (w) a C<sub>6-10</sub>  
 aryloxy group and (x) a 5- to 7-membered heterocyclic  
 group having 1 to 3 hetero atoms selected from  
 nitrogen, oxygen and sulfur in addition to carbon  
 atoms, said heterocyclic group being optionally fused  
 35 with a benzene ring,

- (iv) a hydroxyl group,
- (v) an amino group,
- (vi) a mono-C<sub>1-6</sub> alkylamino group,
- (vii) a di-C<sub>1-6</sub> alkylamino group,
- 5 (viii) a C<sub>1-6</sub> alkyl-carbonyl group,
- (ix) a carboxyl group,
- (x) a C<sub>1-6</sub> alkoxy-carbonyl group,
- (xi) a carbamoyl group,
- (xii) a mono-C<sub>1-6</sub> alkyl-carbamoyl group,
- 10 (xiii) a di-C<sub>1-6</sub> alkyl-carbamoyl group,
- (xiv) a C<sub>6-10</sub> aryl-carbamoyl group,
- (xv) a sulfo group,
- (xvi) a C<sub>1-6</sub> alkylsulfonyl group,
- (xv) a C<sub>6-10</sub> aryl group, and
- 15 (xvi) a C<sub>6-10</sub> aryloxy group,
- (3) A composition as described in the above item (1) wherein R<sup>1</sup> is a hydrogen atom or a C<sub>1-6</sub> alkyl group,
- (4) A composition as described in the above item (1) wherein R<sup>1</sup> is a hydrogen atom or methyl,
- 20 (5) A composition as described in the above item (1) wherein R<sup>1</sup> is a hydrogen atom,
- (6) A composition as described in the above item (1) wherein R<sup>2</sup> is an acyl group,
- (7) A composition as described in the above item (6)
- 25 wherein the acyl group is of the formula -(C=O)-R<sup>4</sup>, -SO<sub>2</sub>-R<sup>4</sup>, -SO-R<sup>4</sup>, -(C=O)NR<sup>5</sup>R<sup>4</sup>, -(C=O)O-R<sup>4</sup>, -(C=S)O-R<sup>4</sup>, or -(C=S)NR<sup>5</sup>-R<sup>4</sup>,  
wherein R<sup>4</sup> is a hydrogen atom, an optionally substituted hydrocarbon group, an optionally substituted heterocyclic group, an optionally substituted lower alkyl-carbonyl group, a carboxyl group, an optionally substituted lower alkoxy-carbonyl group, an optionally substituted mono-lower alkylaminocarbonyl group, an optionally substituted di-
- 35 lower alkylaminocarbonyl group, an optionally


substituted 5- or 7-membered cyclic amino group or an optionally substituted aryloxy group; and  $R^5$  is a hydrogen atom or a lower alkyl group,

(8) A composition as described in the above item (6),  
 5 wherein the acyl group is of the formula  $-(C=O)-R^4$  or  $-(C=O)NHR^4$ , wherein  $R^4$  is a hydrogen atom, an optionally substituted hydrocarbon group, an optionally substituted heterocyclic group, an optionally substituted lower alkyl-carbonyl group, a carboxyl  
 10 group, an optionally substituted lower alkoxy-carbonyl group, an optionally substituted mono-lower alkylaminocarbonyl group, an optionally substituted di-lower alkylaminocarbonyl group, an optionally substituted  
 15 substituted 5- or 7-membered cyclic amino group or an optionally substituted aryloxy group; and  $R^5$  is a hydrogen atom or a lower alkyl group,

(9) A composition as described in the above item (8), wherein  $R^4$  is a group of the formula:

(A)


20



or

(B)

25



wherein  $R^6$  and  $R^7$  independently represent (a) a hydrogen atom, (b) a  $C_{1-6}$  alkyl group optionally substituted with


30 (b-1) a halogen atom, (b-2) a  $C_{1-3}$  alkylenedioxy group, (b-3) a nitro group, (b-4) a cyano group, (b-5) a  $C_{3-6}$  cycloalkyl group, (b-6) a  $C_{1-6}$  alkoxy group optionally having 1 to 3 halogen atoms, (b-7) a  $C_{1-6}$  alkylthio group optionally having 1 to 3 halogen atoms, (b-8) a  
 35 hydroxyl group, (b-9) an amino group, (b-10) a mono- $C_{1-6}$

alkylamino group, (b-11) a di-C<sub>1-6</sub> alkylamino group, (b-12) a C<sub>1-6</sub> alkyl-carbonyl group, (b-13) a carboxyl group, (b-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (b-15) a carbamoyl group, (b-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (b-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (b-18) a C<sub>6-10</sub> aryl-carbamoyl group, (b-19) a sulfo group, (b-20) a C<sub>1-6</sub> alkylsulfonyl group, (b-21) a C<sub>6-10</sub> aryl group, (b-22) a C<sub>6-10</sub> aryloxy group or (b-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (c) a C<sub>3-6</sub> cycloalkyl group, (d) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (e) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (f) a C<sub>7-16</sub> aralkyl group, (g) a hydroxyl group, (h) an amino group, (i) a mono-C<sub>1-6</sub> alkylamino group, (j) a di-C<sub>1-6</sub> alkylamino group, (k) a C<sub>1-6</sub> alkyl-carbonyl group whose alkyl portion may be substituted with (k-1) a halogen atom, (k-2) a C<sub>1-3</sub> alkylenedioxy group, (k-3) a nitro group, (k-4) a cyano group, (k-5) a C<sub>3-6</sub> cycloalkyl group, (k-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (k-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (k-8) a hydroxyl group, (k-9) an amino group, (k-10) a mono-C<sub>1-6</sub> alkylamino group, (k-11) a di-C<sub>1-6</sub> alkylamino group, (k-12) a C<sub>1-6</sub> alkyl-carbonyl group, (k-13) a carboxyl group, (k-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (k-15) a carbamoyl group, (k-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (k-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (k-18) a C<sub>6-10</sub> aryl-carbamoyl group, (k-19) a sulfo group, (k-20) a C<sub>1-6</sub> alkylsulfonyl group, or (k-21) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said

heterocyclic group being optionally fused with a benzene ring, (l) a carboxyl group, (m) a C<sub>1-6</sub> alkoxy-carbonyl group, (n) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (o) a carbamoyl group, (p) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (p-1) a halogen atom, (p-2) a C<sub>1-3</sub> alkylenedioxy group, (p-3) a nitro group, (p-4) a cyano group, (p-5) a C<sub>3-6</sub> cycloalkyl group, (p-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (p-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (p-8) a hydroxyl group, (p-9) an amino group, (p-10) a mono-C<sub>1-6</sub> alkylamino group, (p-11) a di-C<sub>1-6</sub> alkylamino group, (p-12) a C<sub>1-6</sub> alkyl-carbonyl group, (p-13) a carboxyl group, (p-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (p-15) a carbamoyl group, (p-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (p-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (p-18) a C<sub>6-10</sub> aryl-carbamoyl group, (p-19) a sulfo group, (p-20) a C<sub>1-6</sub> alkylsulfonyl group, (p-21) a C<sub>6-10</sub> aryl group, (p-22) a C<sub>6-10</sub> aryloxy group or (p-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (q) a di-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (q-1) a halogen atom, (q-2) a C<sub>1-3</sub> alkylenedioxy group, (q-3) a nitro group, (q-4) a cyano group, (q-5) a C<sub>3-6</sub> cycloalkyl group, (q-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (q-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (q-8) a hydroxyl group, (q-9) an amino group, (q-10) a mono-C<sub>1-6</sub> alkylamino group, (q-11) a di-C<sub>1-6</sub>

alkylamino group, (q-12) a C<sub>1-6</sub> alkyl-carbonyl group,  
 (q-13) a carboxyl group, (q-14) a C<sub>1-6</sub> alkoxy-carbonyl  
 group, (q-15) a carbamoyl group, (q-16) a mono-C<sub>1-6</sub>  
 alkyl-carbamoyl group, (q-17) a di-C<sub>1-6</sub> alkyl-carbamoyl  
 5 group, (q-18) a C<sub>6-10</sub> aryl-carbamoyl group, (q-19) a  
 sulfo group, (q-20) a C<sub>1-6</sub> alkylsulfonyl group, (q-21) a  
 C<sub>6-10</sub> aryl group, (q-22) a C<sub>6-10</sub> aryloxy group or (q-23) a  
 5- to 7-membered heterocyclic group having 1 to 3  
 hetero atoms selected from nitrogen, oxygen and sulfur  
 10 in addition to carbon atoms, said heterocyclic group  
 being optionally fused with a benzene ring, (r) an  
 optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (s)  
 an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group, (t)  
 a sulfo group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub>  
 15 aryl group, (w) a C<sub>6-10</sub> aryloxy group, (x) a C<sub>2-6</sub>  
 alkenylamino group or (y) a 5- to 7-membered  
 heterocyclic group having 1 to 3 hetero atoms selected  
 from nitrogen, oxygen and sulfur in addition to carbon  
 atoms, said heterocyclic group being optionally fused  
 20 with a benzene ring,  
 (10) A composition as described in the above item (8),  
 wherein R<sup>4</sup> is a group of the formula:

(A)



(B)



30

wherein R<sup>6</sup> and R<sup>7</sup> independently represent (a) a  
 hydrogen atom, (b) a C<sub>1-6</sub> alkyl group optionally  
 substituted with

(b-1) a hydroxyl group, (b-2) a di-C<sub>1-6</sub> alkylamino  
 35 group, (b-3) a C<sub>1-6</sub> alkoxy-carbonyl group, or (b-4) a 5-



- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (c) a C<sub>7-16</sub>
- 5 aralkyl group, (d) a C<sub>1-6</sub> alkyl-carbonyl group whose alkyl portion may be substituted with (d-1) a halogen atom, (d-2) a mono-C<sub>1-6</sub> alkylamino group, (d-3) a C<sub>1-6</sub> alkoxy-carbonyl group, or (d-4) a 5- to 7-membered
- 10 heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (e) a C<sub>1-6</sub> alkoxy-carbonyl group, (f) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero
- 15 atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (g) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (g-1) a halogen atom, or (g-2) a C<sub>1-6</sub>
- 20 alkyl-carbonyl group, (h) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (i) an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group, or (j) a C<sub>6-10</sub> aryloxy group, (11) A composition as described in the above item (1) wherein Q<sup>1</sup> and Q<sup>2</sup> are independently a C<sub>1-6</sub> alkylene group
- 25 which may have an oxo group, (12) A composition as described in the above item (1) wherein Q<sup>1</sup> is a C<sub>1-4</sub> alkylene group and Q<sup>2</sup> is a methylene group, (13) A composition as described in the above item (1)
- 30 wherein the ring of the formula:



35 is a 4- to 9-membered monocyclic ring or 6- to 14-

membered bicyclic ring, which may have 1 or 2 unsaturated bonds and may have 1 or 2 substituents in any position other than N and Z,


(14) A composition as described in the above item (1)

5 wherein the ring of the formula:




is

10



15




(15) A composition as described in the above item (1) wherein the ring of the formula:

20



is

25




(16) A composition as described in the above item (1) wherein the ring of the formula:

30



is



5 (17) A composition as described in the above item (13)  
wherein Z is

- (A) an optionally substituted 1, 2-phenylene,
- (B) a group of the formula:




wherein  $\text{Ar}^3$  is an optionally substituted aromatic group, and n is an integer of 0 to 3,

- (C) a group of the formula:




wherein  $\text{Ar}^3$  and n have the same meanings as defined above; and Y is (i) a hydrogen atom, (ii) an  
20 optionally halogenated lower alkyl group, (iii) an optionally halogenated lower alkoxy group, (iv) an optionally halogenated lower alkylthio group, (v) a hydroxyl group, (vi) a cyano group, (vii) an alkyl-carbonyl group, (viii) a lower alkyl-carbonyloxy group,  
25 (ix) a formylamino group, (x) an amino group, (xi) a mono-lower alkylamino group, (xii) a di-lower alkylamino group, (xiii) a carboxyl group, (xiv) a lower alkoxy-carbonyl group or (xv) a lower alkyl-carbonylamino group, or

- 30 (D) a group of the formula:



wherein  $\text{Ar}^3$  and n have the same meanings as  
35 defined above, or

- (E) a group of the formula:




wherein  $\text{Ar}^3$  and  $n$  have the same meanings as  
 5 defined above,

(18) A composition as described in the above item (1)  
 wherein the ring of the formula:



is pyrrolidine, piperidine, piperazine, azepine or  
 azocine, each of which may be fused with a benzene ring  
 and may have a substituent,

15 (19) A composition as described in the above item (13)  
 wherein  $Z$  is a group of the formula:



20

wherein  $\text{Ar}^3$  is an optionally substituted aromatic  
 group,  $n$  is an integer of 0 to 3, and  $Y$  is a hydrogen  
 atom or a hydroxyl group,

(20) A composition as described in the above item (19)  
 25 wherein  $\text{Ar}^3$  is a  $\text{C}_{6-14}$  aryl group or a 5- to 7-membered  
 heterocyclic group having 1 to 3 hetero atoms of 1 or 2  
 kinds selected from nitrogen, oxygen and sulfur in  
 addition to a carbon atom, each of which may have 1 to  
 3 substituents selected from a halogen atom, an  
 30 optionally halogenated  $\text{C}_{1-6}$  alkyl group, and an  
 optionally halogenated  $\text{C}_{1-6}$  alkoxy group,

(21) A composition as described in the above item (19)  
 wherein  $\text{Ar}^3$  is a phenyl group optionally substituted  
 with a halogen atom,

35 (22) A composition as described in the above item (19)  
 wherein  $n$  is 0,


(23) A composition as described in the above item (19) wherein Y is a hydroxyl group,

(24) A composition as described in the above item (1) wherein Ar<sup>1</sup> and Ar<sup>2</sup> independently represent a C<sub>6-14</sub> aryl group or a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms of 1 or 2 kinds selected from nitrogen, oxygen and sulfur in addition to a carbon atom, each of which may have 1 to 3 substituents selected from a halogen atom, an optionally halogenated C<sub>1-6</sub> alkyl group, and an optionally halogenated C<sub>1-6</sub> alkoxy group,


(25) A composition as described in the above item (1) wherein Ar<sup>1</sup> and Ar<sup>2</sup> independently represent phenyl, 4-chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl,

(26) A composition as described in the above item (1), wherein Ar<sup>1</sup> and Ar<sup>2</sup> independently represent phenyl, 4-chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl;


Q<sup>1</sup> is a C<sub>1-4</sub> alkylene group; Q<sup>2</sup> is a methylene group; the group of the formula:



is




wherein Z is a group of the formula:



wherein Ar<sup>3</sup> is a phenyl group optionally substituted with a halogen atom, n is an integer of 0 to 3, and Y is a hydrogen atom or a hydroxyl group;  
R<sup>1</sup> is a hydrogen atom or methyl;


$R^2$  is (I) an  $C_{1-6}$  alkyl group which may be substituted with a  $C_{1-6}$  alkoxy-carbonyl group, a carboxyl group, a  $C_{1-6}$  alkyl-carbonyl group or a formyl group or (II) an acyl group represented by the formula:



wherein  $R^4$  is

- (i) a hydrogen atom,
- (ii) a  $C_{1-6}$  alkyl group which may have 1 to 5 substituents selected from (a) a hydroxyl group, (b) an amino group which may be substituted with a  $C_{1-6}$  alkyl-carbonyl group, (c) a mono- $C_{1-6}$  alkylamino group, (d) a di- $C_{1-6}$  alkylamino group, (e) a carboxyl group, (f) a  $C_{1-6}$  alkoxy-carbonyl group, (g) a mono- $C_{1-6}$  alkyl-carbamoyl group, (h) a sulfo group which may be substituted with amino group, (i) a 5- to 7-membered cyclic amino group which may have an oxo group or which may be substituted with a hydroxyl group, (j) a  $C_{1-6}$  alkoxy-carbamoyl group, and (k) a carbamoyloxy group,
- (iii) a  $C_{2-6}$  alkenyl group,
- (iv) a  $C_{6-10}$  aryl group,
- (v) a 5- to 11-membered heterocyclic group having at least one hetero atom selected from nitrogen, oxygen and sulfur in addition to a carbon atom, said heterocyclic group being optionally fused with a benzene ring,
- (vi) a  $C_{1-6}$  alkyl group which may be substituted with a  $C_{1-6}$  alkyl-carbonyl group,
- (vii) a carboxyl group which may be substituted with a  $C_{1-6}$  alkyl group,
- (viii) a 5- to 7-membered cyclic amino group which may be substituted with
  - (a) a  $C_{1-6}$  alkyl group optionally substituted with (a-1) a hydroxyl group, (a-2) a di- $C_{1-6}$  alkylamino group, (a-3) a  $C_{1-6}$  alkoxy-carbonyl group or (a-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms

- selected from nitrogen, oxygen and sulfur in addition to carbon or fused with benzene ring,
- (b) a C<sub>7-16</sub> aralkyl group, (c) a C<sub>1-6</sub> alkyl-carbonyl group whose alkyl portion may be substituted with (c-1)
- 5 a halogen atom, (c-2) a mono-C<sub>1-6</sub> alkylamino group, (c-3) a C<sub>1-6</sub> alkoxy-carbonyl group or (c-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to a carbon atom, said heterocyclic group being
- 10 optionally fused with a benzene ring,
- (d) a C<sub>1-6</sub> alkoxy-carbonyl group, (e) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to a
- 15 carbon atoms, said heterocyclic group being optionally fused with a benzene ring,
- (f) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with a halogen atom or a C<sub>1-6</sub> alkyl-carbonyl group, (g) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (h) an optionally halogenated
- 20 C<sub>6-10</sub> aryl carbonyl group or (i) a C<sub>1-6</sub> alkoxy-carbamoyl group, or
- (ix) a C<sub>6-10</sub> aryloxy group; and
- R<sup>5</sup> is a hydrogen atom or a C<sub>1-6</sub> alkyl group,
- 25 (27) A compound of the formula:



30

wherein Ar<sup>1</sup>, Ar<sup>2</sup> and Ar<sup>3</sup> independently represent an optionally substituted aromatic group;

35 Q<sup>1</sup> and Q<sup>2</sup> independently represent a divalent C<sub>1-6</sub> aliphatic hydrocarbon group, which may have oxygen or

sulfur within the carbon chain; and

$R^2$  is an optionally substituted hydrocarbon group or an acyl group or a salt thereof (except N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino-2,2-diphenylpentyl]-1-methanesulfonamide hydrochloride, N-[5-[4-chlorophenyl)-4-hydroxypiperidino-2,2-diphenylpentyl]-1-(p-toluene)sulfonamide hydrochloride and N-[5-(4-(4-chlorophenyl)-4-hydroxypiperidino-2,2-diphenylpentyl)-1-(2-thiophene)sulfonamide hydrochloride),

(28) The compound as described in the above item (27)

wherein  $R^2$  is a group of the formula  $-(C=O)-R^4$ ,  $-(C=O)NR^5R^4$ ,  $-(C=O)O-R^4$ ,  $-(C=S)O-R^4$  or  $-(C=S)NR^5R^4$  wherein  $R^4$  is a hydrogen atom, an optionally

substituted hydrocarbon group, an optionally substituted heterocyclic group, an optionally substituted lower alkyl-carbonyl group, a carboxyl group, an optionally substituted lower alkoxycarbonyl group, an optionally substituted mono-lower alkylaminocarbonyl group, an optionally substituted di-lower alkylaminocarbonyl group or an optionally substituted 5- or 7-membered cyclic amino group; and  $R^1$  is a hydrogen atom or a lower alkyl group,

(29) A compound as described in the above item (27), wherein  $R^2$  is the formula  $-(C=O)-R^4$  or  $-(C=O)NH-R^4$ ,


wherein  $R^4$  is a hydrogen atom, an optionally substituted hydrocarbon group, an optionally substituted heterocyclic group, an optionally substituted lower alkyl-carbonyl group, a carboxyl group, an optionally substituted lower alkoxycarbonyl group, an optionally substituted mono-lower alkylaminocarbonyl group, an optionally substituted di-lower alkylaminocarbonyl group or an optionally substituted 5- or 7-membered cyclic amino group,

(30) A compound as described in the above item (28),

wherein  $R^4$  is of the formula:




(A)



or

5

(B)



wherein R<sup>6</sup> and R<sup>7</sup> independently represent (a) a  
10 hydrogen atom, (b) a C<sub>1-6</sub> alkyl group optionally  
substituted with  
(b-1) a halogen atom, (b-2) a C<sub>1-3</sub> alkylenedioxy group,  
(b-3) a nitro group, (b-4) a cyano group, (b-5) a C<sub>3-6</sub>  
cycloalkyl group, (b-6) a C<sub>1-6</sub> alkoxy group optionally  
15 having 1 to 3 halogen atoms, (b-7) a C<sub>1-6</sub> alkylthio  
group optionally having 1 to 3 halogen atoms, (b-8) a  
hydroxyl group, (b-9) an amino group, (b-10) a mono-C<sub>1-6</sub>  
alkylamino group, (b-11) a di-C<sub>1-6</sub> alkylamino group, (b-  
12) a C<sub>1-6</sub> alkyl-carbonyl group, (b-13) a carboxyl  
20 group, (b-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (b-15) a  
carbamoyl group, (b-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl  
group, (b-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (b-18) a  
C<sub>6-10</sub> aryl-carbamoyl group, (b-19) a sulfo group, (b-20)  
a C<sub>1-6</sub> alkylsulfonyl group, (b-21) a C<sub>6-10</sub> aryl group,  
25 (b-22) a C<sub>6-10</sub> aryloxy group or (b-23) a 5- to 7-  
membered heterocyclic group having 1 to 3 hetero atoms  
selected from nitrogen, oxygen and sulfur in addition  
to carbon atoms, said heterocyclic group being  
optionally fused with a benzene ring, (c) a C<sub>3-6</sub>  
30 cycloalkyl group, (d) a C<sub>1-6</sub> alkoxy group optionally  
having 1 to 3 halogen atoms, (e) a C<sub>1-6</sub> alkylthio group  
optionally having 1 to 3 halogen atoms, (f) a C<sub>7-16</sub>  
aralkyl group, (g) a hydroxyl group, (h) an amino  
group, (i) a mono-C<sub>1-6</sub> alkylamino group, (j) a di-C<sub>1-6</sub>  
35 alkylamino group, (k) a C<sub>1-6</sub> alkyl-carbonyl group whose

alkyl portion may be substituted with (k-1) a halogen atom, (k-2) a C<sub>1-3</sub> alkylenedioxy group, (k-3) a nitro group, (k-4) a cyano group, (k-5) a C<sub>3-6</sub> cycloalkyl group, (k-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (k-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (k-8) a hydroxyl group, (k-9) an amino group, (k-10) a mono-C<sub>1-6</sub> alkylamino group, (k-11) a di-C<sub>1-6</sub> alkylamino group, (k-12) a C<sub>1-6</sub> alkyl-carbonyl group, (k-13) a carboxyl group, (k-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (k-15) a carbamoyl group, (k-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (k-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (k-18) a C<sub>6-10</sub> aryl-carbamoyl group, (k-19) a sulfo group, (k-20) a C<sub>1-6</sub> alkylsulfonyl group, or (k-21) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (l) a carboxyl group, (m) a C<sub>1-6</sub> alkoxy-carbonyl group, (n) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (o) a carbamoyl group, (p) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (p-1) a halogen atom, (p-2) a C<sub>1-3</sub> alkylenedioxy group, (p-3) a nitro group, (p-4) a cyano group, (p-5) a C<sub>3-6</sub> cycloalkyl group, (p-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (p-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (p-8) a hydroxyl group, (p-9) an amino group, (p-10) a mono-C<sub>1-6</sub> alkylamino group, (p-11) a di-C<sub>1-6</sub> alkylamino group, (p-12) a C<sub>1-6</sub> alkyl-carbonyl group, (p-13) a carboxyl group, (p-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (p-15) a carbamoyl group, (p-16) a mono-C<sub>1-6</sub>

alkyl-carbamoyl group, (p-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (p-18) a C<sub>6-10</sub> aryl-carbamoyl group, (p-19) a sulfo group, (p-20) a C<sub>1-6</sub> alkylsulfonyl group, (p-21) a C<sub>6-10</sub> aryl group, (p-22) a C<sub>6-10</sub> aryloxy group or (p-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (q) a di-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (q-1) a halogen atom, (q-2) a C<sub>1-3</sub> alkylenedioxy group, (q-3) a nitro group, (q-4) a cyano group, (q-5) a C<sub>3-6</sub> cycloalkyl group, (q-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (q-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (q-8) a hydroxyl group, (q-9) an amino group, (q-10) a mono-C<sub>1-6</sub> alkylamino group, (q-11) a di-C<sub>1-6</sub> alkylamino group, (q-12) a C<sub>1-6</sub> alkyl-carbonyl group, (q-13) a carboxyl group, (q-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (q-15) a carbamoyl group, (q-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (q-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (q-18) a C<sub>6-10</sub> aryl-carbamoyl group, (q-19) a sulfo group, (q-20) a C<sub>1-6</sub> alkylsulfonyl group, (q-21) a C<sub>6-10</sub> aryl group, (q-22) a C<sub>6-10</sub> aryloxy group or (q-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (r) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (s) an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group, (t) a sulfo group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, (w) a C<sub>6-10</sub> aryloxy group, (x) a C<sub>2-6</sub> alkenylamino group or (y) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused

with a benzene ring,

(31) A compound as described in the above item (27) wherein  $Q^1$  and  $Q^2$  are independently a  $C_{1-6}$  alkylene group which may have an oxo group,

5 (32) A compound as described in the above item (27) wherein  $Q^1$  is a  $C_{1-4}$  alkylene group and  $Q^2$  is a methylene group,


(33) A compound as described in the above item (27) wherein  $Ar^3$  is a phenyl group optionally substituted  
10 with a halogen atom,

(34) A compound as described in the above item (27) wherein  $Ar^1$  and  $Ar^2$  independently represent a  $C_{6-14}$  aryl group or a 5- to 7-membered heterocyclic groups having  
15 1 to 3 hetero atoms of 1 or 2 kinds selected from nitrogen, oxygen and sulfur in addition to a carbon atom, each of which may have 1 to 3 substituents selected from a halogen atom, an optionally halogenated  $C_{1-6}$  alkyl group, and an optionally halogenated  $C_{1-6}$  alkoxy group,

20 (35) A compound as described in the above item (27) wherein  $Ar^1$  and  $Ar^2$  independently represent phenyl, 4-chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl,

(36) A compound as described in the above item (27),  
25 wherein  $Ar^1$  and  $Ar^2$  independently represent phenyl, 4-chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl;

$Q^1$  is a  $C_{1-4}$  alkylene group;  $Q^2$  is a methylene group;  
 $R^2$  is (I) a  $C_{1-6}$  alkyl group which may be substituted  
30 with a  $C_{1-6}$  alkoxy-carbonyl group, a carboxyl group, a  $C_{1-6}$  alkyl-carbonyl group or a formyl group or (II) an acyl group represented by the formula:



wherein  $R^4$  is

35 (i) a hydrogen atom,

- (ii) a C<sub>1-6</sub> alkyl group which may have 1 to 5 substituents selected from (a) a hydroxyl group, (b) an amino group which may be substituted with a C<sub>1-6</sub> alkyl-carbonyl group, (c) a mono-C<sub>1-6</sub> alkylamino group, (d) a 5 di-C<sub>1-6</sub> alkylamino group, (e) a carboxyl group, (f) a C<sub>1-6</sub> alkoxy-carbonyl group, (g) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (h) a sulfo group which may be substituted with amino group (i) a 5- to 7-membered cyclic amino group which may have an oxo group or which may be substituted 10 with a hydroxyl group, (j) a C<sub>1-6</sub> alkoxy-carbamoyl group, and (k) a carbamoyloxy group.
- (iii) a C<sub>2-6</sub> alkenyl group,
- (iv) a C<sub>6-10</sub> aryl group,
- (v) a 5- to 11-membered heterocyclic group having at 15 least one hetero atom selected from nitrogen, oxygen and sulfur in addition to a carbon atom, said heterocyclic group being optionally fused with a benzene ring,
- (vi) a C<sub>1-6</sub> alkyl group which may be substituted with a 20 C<sub>1-6</sub> alkyl-carbonyl group,
- (vii) a carboxyl group which may be substituted with a C<sub>1-6</sub> alkyl group,
- (viii) a 5- to 7-membered cyclic amino group which may be substituted with
- 25 (a) a C<sub>1-6</sub> alkyl group optionally substituted with (a-1) a hydroxyl group, (a-2) a di-C<sub>1-6</sub> alkylamino group, (a-3) a C<sub>1-6</sub> alkoxy-carbonyl group or (a-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition 30 to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,
- (b) a C<sub>7-16</sub> aralkyl group, (c) a C<sub>1-6</sub> alkyl-carbonyl group whose alkyl portion may be substituted with (c-1) a halogen atom, (c-2) a mono-C<sub>1-6</sub> alkylamino group, (c-3) a C<sub>1-6</sub> alkoxy-carbonyl group or (c-4) a 5- to 7-
- 35


membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,

- 5 (d) a C<sub>1-6</sub> alkoxy-carbonyl group, (e) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused  
10 with a benzene ring,  
(f) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with a halogen atom or a C<sub>1-6</sub> alkyl-carbonyl group, (g) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (h) an optionally halogenated  
15 C<sub>6-10</sub> aryl carbonyl group or (i) a C<sub>1-6</sub> alkoxy-carbamoyl group, or  
(ix) a C<sub>6-10</sub> aryloxy group;


R<sup>5</sup> is a hydrogen atom or a C<sub>1-6</sub> alkyl group; and

Ar<sup>3</sup> is a phenyl group optionally substituted with a  
20 halogen atom,

(37) A process for producing a compound of the formula:




wherein R<sup>2</sup> is an acyl group, and the other symbols have the same meanings as described in the above item  
30 (27) or a salt thereof, which comprises subjecting a compound of the formula:



(IX')


wherein the all symbols have the same meanings as described in the above item (27) or a salt thereof to the acylation reaction,

(38) A process for producing a compound of the formula:



(II')

wherein  $\text{R}^4$  represents a hydrogen atom, an optionally substituted hydrocarbon group, an optionally substituted heterocyclic group, an optionally substituted lower alkyl-carbonyl group, a carboxyl group, an optionally substituted lower alkoxy-carbonyl group, an optionally substituted mono-lower alkylaminocarbonyl group, an optionally substituted di-lower alkylaminocarbonyl group or an optionally substituted 5- or 6-membered cyclic amino group; and  $\text{R}^5$  is a hydrogen atom or a lower alkyl group, and the other symbols have the same meanings as defined in Claim 27 or a salt thereof, which comprises reacting a compound of the formula:



(X')

wherein Ph is a phenyl group, and the other symbols have the same meanings as defined above or a salt thereof with a compound of the formula:



[XI]

wherein R<sup>4</sup> and R<sup>5</sup> have the same meanings as defined above or a salt thereof,

(39) A composition as described in the above item (1) which is a prophylactic or therapeutic agent for inflammatory diseases,

(40) A composition as described in the above item (1) which is a prophylactic or therapeutic agent for allergic diseases,


(41) A composition as described in the above item (1) which is a prophylactic or therapeutic agent for arteriosclerosis, bronchial asthma, atopy, multiple sclerosis or rheumatoid arthritis,

(42) A pharmaceutical composition comprising the compound as described in the above item (27),

(43) A MIP-1 $\alpha$ /RANTES receptor antagonist comprising the compound as described in the above item (27),

(44) A method of treating or preventing inflammatory diseases or allergic diseases which comprises administering to a mammal in need an effective amount of a compound of the formula:





5


wherein  $\text{Ar}^1$  and  $\text{Ar}^2$  independently represent an optionally substituted aromatic group;

$\text{Q}^1$  and  $\text{Q}^2$  independently represent an optionally substituted divalent  $\text{C}_{1-6}$  aliphatic hydrocarbon group which may have oxygen or sulfur within the carbon chain;

$\text{R}^1$  is a hydrogen atom, an optionally substituted lower alkyl group or an optionally substituted lower alkyl-carbonyl group;

$\text{R}^2$  is an optionally substituted hydrocarbon group or an acyl group, or  $\text{R}^1$  and  $\text{R}^2$ , taken together with the adjacent nitrogen atom, may form an optionally substituted nitrogen-containing heterocyclic ring; and a group of the formula:

20




is an optionally substituted monocyclic or fused nitrogen-containing heterocyclic ring, or a salt thereof,

25

(45) Use of a compound of the formula:

30



wherein  $\text{Ar}^1$  and  $\text{Ar}^2$  independently represent an optionally substituted aromatic group;

$\text{Q}^1$  and  $\text{Q}^2$  independently represent an optionally substituted divalent  $\text{C}_{1-6}$  aliphatic hydrocarbon group


35

which may have oxygen or sulfur within the carbon chain;

$R^1$  is a hydrogen atom, an optionally substituted lower alkyl group or an optionally substituted lower alkyl-carbonyl group;

$R^2$  is an optionally substituted hydrocarbon group or an acyl group, or  $R^1$  and  $R^2$ , taken together with the adjacent nitrogen atom, form an optionally substituted nitrogen-containing heterocyclic ring; and

a group of the formula:



is an optionally substituted monocyclic or fused nitrogen-containing heterocyclic ring or a salt thereof, for the manufacture of a medicament for treating or preventing inflammatory diseases or allergic diseases, and

(46) A compound as described in the above item (27) which is Examples

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(piperidin-4-yl)urea,

Ethyl 4-[4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino]

piperidino-4-oxobutyrates,

N-Ethyl-4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino-1-piperidinecarboxamide,

N-Ethoxycarbonylmethyl-4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino-1-piperidinecarboxamide,

Ethyl 3-[4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino]piperidino-3-oxopropionate,

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(1-ethylpiperidin-4-yl)urea,

1-[(Piperidin-4-yl)carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane,  
1-[[N-Ethylcarbamoyl)piperidin-4-yl]carboamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane,  
5 e,  
1-[[N-(Ethoxycarbonylacetyl)piperidin-4-yl]carboamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane,  
1-[[N-(3-Methoxycarbonylpropionyl)piperidin-4-yl]carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane or a salt thereof.

#### Detailed description

The aromatic group of the "optionally substituted aromatic group" for Ar<sup>1</sup>, Ar<sup>2</sup> and Ar<sup>3</sup> includes, for  
15 example, "aromatic hydrocarbon groups" and "heteroaromatic groups" and these groups may have any number (preferably 1 to 5, more preferably 1 to 3, further more preferably 1 or 2) of substituents in any substitutable position.

20 The "aromatic hydrocarbon group" mentioned above includes, for example, monocyclic or fused polycyclic aromatic hydrocarbon groups having 6 to 14 carbon atoms. Specific examples thereof include C<sub>6-14</sub> aryl groups such as phenyl, 1-naphthyl, 2-naphthyl, indenyl, anthryl, etc. Among them, phenyl, 1-naphthyl and  
25 2-naphthyl are preferred, and phenyl is particularly preferred.

The "heteroaromatic group" mentioned above includes, for example, 5- to 11-membered monocyclic or  
30 fused heteroaromatic groups having at least one (e.g. 1 to 4, preferably 1 to 3, more preferably 1 or 2) of 1 or 2 kinds of hetero atoms selected from a nitrogen atom, a sulfur atom and an oxygen atom in addition to a carbon atom. Specific examples thereof include  
35 aromatic heterocyclic group such as thiophene, benzo[b]thiophene, benzo[b]furan, benzimidazole,

benzoxazole, benzothiazole, benzisothiazole,  
naphtho[2,3-b]thiophene, thianthrene, furan,  
isoindolizine, xanthrene, phenoxathiin, pyrrole,  
imidazole, pyrazole, pyridine, pyrazine, pyrimidine,  
5 pyridazine, indol, isoindol, 1H-indazole, purine,  
4H-quinolizine, isoquinoline, quinoline, phtharazine,  
naphthyridine, quinoxaline, cinnoline, carbazole,  
 $\beta$ -carboline, phenanthridine, acridine, phenazine,  
isothiazole, phenothiazine, isoxazole, furazane,  
10 phenoxazine, isochroman, etc., or a monovalent group  
obtained by eliminating any hydrogen from a ring formed  
by condensing these rings (preferably monocyclic  
heterocycle mentioned above) with one or a plurality  
(preferably 1 or 2, more preferably 1) of aromatic  
15 rings (e.g. aromatic hydrocarbon group, preferably  
benzene ring, etc.). The preferred "aromatic  
heterocyclic group" include, for example, 2-pyridyl,  
3-pyridyl, 4-pyridyl, 2-quinolyl, 3-quinolyl,  
4-quinolyl, 5-quinolyl, 8-quinolyl, 1-isoquinolyl,  
20 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl, 1-indolyl,  
2-indolyl, 3-indolyl, 2-benzothiazolyl,  
2-benzo[b]thienyl, benzo[b]furanyl, 2-thienyl,  
3-thienyl, etc. The more preferred one include, for  
example, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl,  
25 4-pyridyl, 2-quinolyl, 1-isoquinolyl, 1-indolyl,  
2-indolyl, 2-benzothiazolyl, etc. Among them,  
2-pyridyl is commonly used.

The substituent that may be present on the  
"optionally substituted aromatic ring in any position"  
30 for Ar<sup>1</sup>, Ar<sup>2</sup> and Ar<sup>3</sup> includes, for example, a halogen  
atom (e.g. fluorine, chlorine, bromine, iodine, etc.),  
a lower alkylenedioxy group (e.g. C<sub>1-3</sub> alkylenedioxy  
such as methylenedioxy, ethylenedioxy, etc.), a nitro  
group, a cyano group, an optionally halogenated lower  
35 alkyl group, an optionally halogenated lower alkenyl  
group, an optionally halogenated lower alkynyl group, a

lower cycloalkyl group (e.g. C<sub>3-6</sub> cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc.), an optionally halogenated lower alkoxy group, an optionally halogenated lower alkylthio group, a  
5 hydroxyl group, an amino group, a mono-lower alkylamino group (e.g. mono-C<sub>1-6</sub> alkylamino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), a di-lower alkylamino group (e.g. di-C<sub>1-6</sub>  
10 alkylamino such as dimethylamino, diethylamino, dipropylamino, dibutylamino, etc.), a 5- to 7-membered cyclic amino group (e.g. morpholino, piperazin-1-yl, piperidino, pyrrolidin-1-yl, etc.), an acylamino group, a lower alkyl-carbonyl group (e.g. C<sub>1-6</sub> alkyl-carbonyl such as acetyl, propionyl, etc.), a carboxyl group, a  
15 lower alkoxy-carbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.), a carbamoyl group, a mono-lower alkyl-carbamoyl group (e.g. mono-C<sub>1-6</sub> alkyl-carbamoyl such as methylcarbamoyl,  
20 ethylcarbamoyl, etc.), a di-lower alkyl-carbamoyl group (e.g. di-C<sub>1-6</sub> alkyl-carbamoyl such as dimethylcarbamoyl, diethylcarbamoyl, etc.), an aryl-carbamoyl group (e.g. C<sub>6-10</sub> aryl-carbamoyl such as phenylcarbamoyl, naphthylcarbamoyl, etc.), a sulfo group, a lower  
25 alkylsulfonyl group (e.g. C<sub>1-6</sub> alkylsulfonyl such as methylsulfonyl, ethylsulfonyl, etc.), an aryl group (e.g. C<sub>6-10</sub> aryl such as phenyl, naphthyl, etc.) or an aryloxy group (e.g. C<sub>6-10</sub> aryloxy such as phenyloxy, naphthyloxy, etc.).

30 The "optionally halogenated lower alkyl group" mentioned above includes, for example, a lower alkyl group optionally having 1 to 3 halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.) (e.g. C<sub>1-6</sub> alkyl such as methyl, ethyl, propyl, isopropyl, butyl,  
35 isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.).

Specific examples thereof include methyl, chloromethyl, difluoromethyl, trichloromethyl, trifluoromethyl, ethyl, 2-bromoethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, propyl, 3,3,3-trifluoropropyl, isopropyl, butyl, 4,4,4-trifluorobutyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, 5,5,5-trifluoropentyl, hexyl, 6,6,6-trifluorohexyl, etc.

The "optionally halogenated lower alkenyl group" and "optionally halogenated lower alkynyl group" include, for example, a lower alkenyl group optionally having 1 to 3 halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.) (e.g. C<sub>2-6</sub> alkenyl such as vinyl, propenyl, isopropenyl, 2-buten-1-yl, 4-penten-1-yl, 5-hexen-1-yl, etc.) or a lower alkynyl group optionally having 1 to 3 halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.) (e.g. C<sub>2-6</sub> alkynyl such as 2-butyne-1-yl, 4-pentyne-1-yl, 5-hexyne-1-yl, etc.).

The "optionally halogenated lower alkoxy group" include, for example, a lower alkoxy group optionally having 1 to 3 halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.) (e.g. C<sub>1-6</sub> alkoxy such as methoxy, ethoxy, butoxy, propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, etc.). Specific examples thereof include methoxy, difluoromethoxy, trifluoromethoxy, ethoxy, 2,2,2-trifluoroethoxy, n-propoxy, isopropoxy, n-butoxy, 4,4,4-trifluorobutoxy, isobutoxy, sec-butoxy, pentyloxy, hexyloxy, etc.

The "optionally halogenated lower alkylthio group" include, for example, a lower alkylthio group optionally having 1 to 3 halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.) (e.g. C<sub>1-6</sub> alkylthio such as methylthio, ethylthio, n-propylthio, isopropylthio, n-butylthio, isobutylthio, sec-butylthio, tert-butylthio, etc.). Specific examples thereof include methylthio,

difluoromethylthio, trifluoromethylthio, ethylthio, propylthio, isopropylthio, butylthio, 4,4,4-trifluorobutylthio, pentylthio, hexylthio, etc.

5 The "acylamino group" include, for example, -NHCOOR<sup>3</sup>, -NHCONHR<sup>3</sup>, -NHCOR<sup>3</sup> or -NHSO<sub>2</sub>R<sup>3</sup> (R<sup>3</sup> is an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group, preferably optionally substituted hydrocarbon group).

10 The substituent that may be present on the "optionally substituted aromatic ring in any position" for Ar<sup>1</sup>, Ar<sup>2</sup> and Ar<sup>3</sup> includes, for preferred example, a halogen atom, an optionally halogenated C<sub>1-6</sub> alkyl group, optionally halogenated C<sub>1-6</sub> alkoxy group, a C<sub>1-3</sub> alkylenedioxy group (particularly methylenedioxy), a  
15 cyano group, a hydroxyl group, etc. Among them, a halogen atom, an optionally halogenated C<sub>1-6</sub> alkyl group and an optionally halogenated C<sub>1-6</sub> alkoxy group are particularly preferred, and a halogen atom is commonly used.

20 The preferred one for Ar<sup>1</sup> and Ar<sup>2</sup> include independently, for example, optionally halogenated phenyl (e.g. phenyl, 4-chlorophenyl, 4-fluorophenyl, etc.) 2-pyridyl, 3-pyridyl and 4-pyridyl. Among them, phenyl and 2-pyridyl are more preferred. As Ar<sup>1</sup> and  
25 Ar<sup>2</sup>, phenyl is commonly used independently.

As Ar<sup>3</sup>, a C<sub>1-3</sub> alkyl group optionally substituted with 1 to 3 halogen atoms, a C<sub>1-3</sub> alkoxy group optionally substituted with 1 to 3 halogen atoms or a phenyl group optionally substituted with halogen  
30 (preferably, chlorine, fluorine, etc.) (e.g. 4-chlorophenyl, 4-fluorophenyl, 4-methoxyphenyl, 3,5-dichlorophenyl, 3,5-difluorophenyl, 4-trifluoromethylphenyl, etc.) or 2-pyridyl, 3-pyridyl, 4-pyridyl are preferred. Among them, optionally  
35 halogenated phenyl is preferred and 4-chlorophenyl is

particularly preferred.

The "optionally substituted hydrocarbon group" for  $R^2$  and  $R^3$  represents a group obtained by eliminating one hydrogen from a hydrocarbon compound and examples thereof include acyclic or cyclic hydrocarbon groups such as alkyl, alkenyl, cycloalkyl, aryl, aralkyl, etc. Preferred are acyclic or cyclic hydrocarbon groups having 1 to 16 carbon atoms as described below.

(a) a lower alkyl group (e.g.  $C_{1-6}$  alkyl such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.)

(b) a lower alkenyl group ( $C_{2-6}$  alkenyl such as vinyl, allyl, isopropenyl, butenyl, isobutenyl, sec-butenyl, etc.)

(c) a lower alkynyl group ( $C_{2-6}$  alkynyl such as propargyl, ethynyl, butynyl, 1-hexynyl, etc.)

(d) a lower cycloalkyl group (e.g.  $C_{3-6}$  cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl optionally fused with a benzene ring optionally having 1 to 3 lower alkoxy groups (e.g.  $C_{1-6}$  alkoxy such as methoxy, etc.))

(e) an aryl group (e.g.  $C_{6-17}$  aryl group such as phenyl, 1-naphthyl, 2-naphthyl, biphenyl, 2-indenyl, 2-anthryl, etc., preferably phenyl)

(f) an aralkyl group (e.g.  $C_{7-16}$  aralkyl group such as benzyl, phenethyl, diphenylmethyl, triphenylmethyl, 1-naphthylmethyl, 2-naphthylmethyl, 2-diphenylethyl, 3-phenylpropyl, 4-phenylbutyl, 5-phenylpentyl, etc., preferably benzyl). Among them, a lower alkyl group, an aryl group and an aralkyl group are preferred.

Especially, a lower alkyl group is preferred.

The substituent which may be present on the "optionally substituted hydrocarbon group" for  $R^2$  and  $R^3$  may have 1 to 5, preferably 1 to 3 substituents in



substitutable positions, and where the number of substituents is 2 or more, the substituent groups may be the same or different.

The substituent that may be present on the

5 "optionally substituted hydrocarbon group" includes, for example, a halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), a lower alkylenedioxy group (e.g. C<sub>1-3</sub> alkylenedioxy such as methylenedioxy, ethylenedioxy, etc.), a nitro group, a cyano group, an

10 optionally halogenated lower alkyl group, a lower cycloalkyl group (e.g. C<sub>3-6</sub> cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc.), an optionally halogenated lower alkoxy group, an optionally halogenated lower alkylthio group, a

15 hydroxyl group, an amino group, a mono-lower alkylamino group (e.g. mono-C<sub>1-6</sub> alkylamino such as methylamino, ethylamino, etc.), a di-lower alkylamino group (e.g. di-C<sub>1-6</sub> alkylamino such as dimethylamino, diethylamino, etc.), a lower alkyl-carbonyl group (e.g. C<sub>1-6</sub>

20 alkyl-carbonyl such as acetyl, ethylcarbonyl, etc.), a carboxyl group, a lower alkoxy-carbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.), a carbamoyl group, a mono-lower alkyl-carbamoyl group

25 (e.g. mono-C<sub>1-6</sub> alkyl-carbamoyl such as methylcarbamoyl, ethylcarbamoyl, etc.), a di-lower alkyl-carbamoyl group (e.g. di-C<sub>1-6</sub> alkyl-carbamoyl such as dimethylcarbamoyl, diethylcarbamoyl, etc.), a sulfo group, a lower alkylsulfonyl group (e.g. C<sub>1-6</sub> alkylsulfonyl such as

30 methylsulfonyl, ethylsulfonyl, etc.), an aryl group (e.g. C<sub>6-10</sub> aryl such as phenyl, naphthyl, etc.), an aryloxy group (e.g. C<sub>6-10</sub> aryloxy such as phenyloxy, naphthyloxy, etc.) or a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from a

35 nitrogen atom, an oxygen atom and a sulfur atom in

addition to a carbon atom or a group fused with a benzene ring.

The "optionally halogenated lower alkyl group," "optionally halogenated lower alkoxy group" and  
5 "optionally halogenated lower alkylthio group" include the same substituents as mentioned for the aromatic group.

The "aryl group (preferably phenyl) and aryloxy group (preferably phenyloxy)" may have the same  
10 substituents mentioned for the "optionally substituted aromatic group in any position."

The "5- to 7-membered heterocyclic group or a group fused with a benzene ring" include, for example, 5- to 7-membered (preferably 5- or 6-membered)  
15 heterocyclic group having 1 to 3, preferably 1 or 2 hetero atoms of 1 or 2 kinds selected from a nitrogen atom, an oxygen atom and a sulfur atom in addition to a carbon atom. Specific examples thereof include 1-, 2- or 3-pyrrolidinyl, 2- or 4-imidazoliny, 2-, 3- or  
20 4-pyrazolidinyl, piperidino, 2-, 3- or 4-piperidyl, 1- or 2-piperazinyl, morpholino, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-furyl, 3-furyl, pyrazinyl, 2-pyrimidinyl, 3-pyrrolyl, 3-pyridazinyl, 3-isothiazolyl, 3-isoxazolyl, etc. These groups may be  
25 fused with a benzene ring in any position. Furthermore, the "5- to 7-membered heterocyclic group or a group fused with a benzene ring" may have 1 to 3 substituents in substitutable positions.

The substituent include substituents that may be  
30 present on the "optionally substituted hydrocarbon group" for  $Ar^1$ ,  $Ar^2$  and  $Ar^3$ . The preferred one include, for example, a halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), a lower alkylenedioxy group (e.g.  $C_{1-3}$  alkylenedioxy such as methylenedioxy,  
35 ethylenedioxy, etc.), a nitro group, a cyano group, an optionally halogenated lower alkyl group, a lower

cycloalkyl group (e.g. C<sub>3-6</sub> cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc.), an optionally halogenated lower alkoxy group, an optionally halogenated lower alkylthio group, a hydroxyl group, an amino group, a mono-lower alkylamino group (e.g. mono-C<sub>1-6</sub> alkylamino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), a di-lower alkylamino group (e.g. di-C<sub>1-6</sub> alkylamino such as dimethylamino, diethylamino, dipropylamino, dibutylamino, etc.), a 5- to 7-membered cyclic amino group (e.g. morpholino, piperazin-1-yl, piperidino, pyrrolidin-1-yl, etc.), a lower alkyl-carbonyl group (e.g. C<sub>1-6</sub> alkyl-carbonyl such as acetyl, propionyl, etc.), a carboxyl group, a lower alkoxy-carbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.), a carbamoyl group, a mono-lower alkyl-carbamoyl group (e.g. mono-C<sub>1-6</sub> alkyl-carbamoyl such as methylcarbamoyl, ethylcarbamoyl, etc.), a di-lower alkyl-carbamoyl group (e.g. di-C<sub>1-6</sub> alkyl-carbamoyl such as dimethylcarbamoyl, diethylcarbamoyl, etc.), an aryl-carbamoyl group (e.g. C<sub>6-10</sub> aryl-carbamoyl such as phenylcarbamoyl, naphthylcarbamoyl, etc.), a sulfo group, a lower alkylsulfonyl group (e.g. C<sub>1-6</sub> alkylsulfonyl such as methylsulfonyl, ethylsulfonyl, etc.), an aryl group (e.g. C<sub>6-10</sub> aryl such as phenyl, naphthyl, etc.) or an aryloxy group (e.g. C<sub>6-10</sub> aryloxy such as phenyloxy, naphthyloxy, etc.).

The "optionally halogenated lower alkyl group," "optionally halogenated lower alkoxy group" and "optionally halogenated lower alkylthio group" include the same substituents mentioned for the "optionally substituted aromatic group" for Ar<sup>1</sup>, Ar<sup>2</sup> and Ar<sup>3</sup>.

The preferred "optionally substituted hydrocarbon"

for  $R^2$  is a  $C_{1-6}$  alkyl group which may be substituted with a  $C_{1-6}$  alkoxy-carbonyl group, a carboxyl group, a  $C_{1-6}$  alkyl-carbonyl group, or a formyl group.

The "acyl group" for  $R^2$  includes, for example,  
5  $-(C=O)-R^4$ ,  $-SO_2-R^4$ ,  $-SO-R^4$ ,  $-(C=O)NR^5-R^4$ ,  $-(C=O)O-R^4$ ,  
 $-(C=S)O-R^4$ ,  $-(C=S)NR^5-R^4$  ( $R^4$  is a hydrogen atom, an optionally substituted hydrocarbon group, an optionally substituted heterocyclic group, an optionally substituted lower alkyl-carbonyl group (e.g.  $C_{1-6}$  alkyl-carbonyl such as acetyl, propionyl, butyryl, etc.), a  
10 carboxyl group, an optionally substituted lower alkoxy-carbonyl group (e.g.  $C_{1-6}$  alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.), an optionally substituted  
15 mono-lower alkylaminocarbonyl group (e.g.  $C_{1-6}$  alkyl-carbamoyl such as methylaminocarbonyl, ethylaminocarbonyl, propylaminocarbonyl, butylaminocarbonyl, etc.), an optionally substituted  
20 di-lower alkylaminocarbonyl group (e.g.  $C_{1-6}$  alkyl-carbamoyl such as dimethylaminocarbonyl, diethylaminocarbonyl, dipropylaminocarbonyl, dibutylaminocarbonyl, etc.), an optionally substituted  
5- or 7-membered cyclic amino group (e.g. 2-piperidyl, 3-piperidyl, 4-piperidyl, 1-pyrrolidinyl,  
25 3-pyrrolidinyl, 2-piperazyl, etc.) or an optionally substituted aryloxy group (e.g.  $C_{6-10}$  aryloxy group such as phenyloxy etc.); and  $R^5$  is a hydrogen atom or a lower alkyl group (e.g.  $C_{1-6}$  alkyl such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl,  
30 tert-butyl, pentyl, hexyl, etc., where  $C_{1-3}$  alkyl such as methyl, ethyl, propyl, isopropyl, etc. are particularly preferred)).

Among them,  $-(C=O)-R^4$ ,  $-SO_2-R^4$ ,  $-SO-R^4$ ,  $-(C=O)NR^5-R^4$  and  $-(C=O)O-R^4$  ( $R^4$  and  $R^5$  have the same meanings as  
35 defined above) are preferred, and  $-(C=O)-R^4$ ,  $-SO_2-R^4$ ,

$-(C=O)NR^5-R^4$  and  $-(C=O)O-R^4$  ( $R^4$  and  $R^5$  have the same meanings as defined above) are more preferred. Especially preferred is  $-(C=O)-R^4$  or  $-(C=O)NH-R^4$  ( $R^4$  is the same meanings as defined above).

5        The preferred example of  $R^2$  is (1) a  $C_{1-6}$  alkyl group which may be substituted with a  $C_{1-6}$  alkoxy-carbonyl group or a carboxyl group, a  $C_{1-6}$  alkyl-carbonyl group or a formyl group, or (2) acyl group.

Especially, acyl group is commonly used.

10        The "hydrocarbon group" of the "optionally substituted hydrocarbon group" for  $R^4$  represents a group obtained by eliminating one hydrogen from a hydrocarbon compound, and examples thereof include acyclic or cyclic hydrocarbon groups such as alkyl,  
15        alkenyl, alkynyl, cycloalkyl, aryl, aralkyl, etc. Specific examples thereof include the same substituents mentioned for the "hydrocarbon group" of the "optionally substituted hydrocarbon group" for  $R^2$  and  $R^3$ . Among them, acyclic or cyclic hydrocarbon groups  
20        having 1 to 16 carbon atoms are preferred, particularly lower ( $C_{1-6}$ ) alkyl group, lower ( $C_{2-6}$ ) alkenyl group or lower ( $C_{6-10}$ ) aryl group is preferred. A lower ( $C_{1-6}$ ) alkyl group is commonly used.

25        The preferred substituent which may be present on the "hydrocarbon group", "heterocyclic group", "lower alkyl-carbonyl group", "a carboxyl group", "lower alkoxy-carbonyl group", "mono-lower alkylaminocarbonyl group", "di-lower alkylaminocarbonyl group", "5- or 7-membered cyclic amino group" and "aryloxy group" for  
30         $R^4$  includes, for example, (i) a halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), (ii) a lower alkylenedioxy group (e.g.  $C_{1-3}$  alkylenedioxy such as methylenedioxy, ethylenedioxy, etc.), (iii) a nitro group, (iv) a cyano group, (v) a  $C_{1-6}$  alkyl group  
35        optionally substituted with (1) a halogen atom, (2) a

C<sub>1-3</sub> alkylenedioxy group, (3) a nitro group, (4) a cyano group, (5) a C<sub>3-6</sub> cycloalkyl group, (6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (8) a hydroxyl group, (9) an amino group, (10) a mono-C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub> alkylamino group, (12) a C<sub>1-6</sub> alkyl-carbonyl group, (13) a carboxyl group, (14) a C<sub>1-6</sub> alkoxy-carbonyl group, (15) a carbamoyl group, (16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (18) a C<sub>6-10</sub> aryl-carbamoyl group, (19) a sulfo group, (20) a C<sub>1-6</sub> alkylsulfonyl group, (21) a C<sub>6-10</sub> aryl group, (22) a C<sub>6-10</sub> aryloxy group or (23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (vi) a C<sub>3-6</sub> cycloalkyl group, (vii) an optionally halogenated lower alkoxy group, (viii) an optionally halogenated lower alkylthio group, (ix) a C<sub>7-16</sub> aralkyl group, (x) a hydroxyl group, (xi) an amino group which may be substituted with a C<sub>1-6</sub> alkyl-carbonyl group, (xii) a mono-lower alkylamino group (e.g. C<sub>1-6</sub> alkylamino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), (xiii) a di-lower alkylamino group (e.g. di-lower alkylamino such as dimethylamino, diethylamino, dipropylamino, dibutylamino, etc.), (xiv) a 5- or 7-membered cyclic amino group optionally having hydroxy or oxo (e.g. morpholino, piperazin-1-yl, piperidino, pyrrolidin-1-yl, 2-pyrrolidone-1-yl, 2-pyridone-1-yl, etc.), (xv) a lower alkyl-carbonyl group (e.g. C<sub>1-6</sub> alkyl-carbonyl such as acetyl, propionyl, etc.), whose alkyl portion may be substituted with (1) a halogen atom, (2) a C<sub>1-3</sub> alkylenedioxy group, (3) a nitro group, (4) a cyano group, (5) a C<sub>3-6</sub> cycloalkyl group, (6) a C<sub>1-</sub>

6 alkoxy group optionally having 1 to 3 halogen atoms,  
(7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3  
halogen atoms, (8) a hydroxyl group, (9) an amino  
group, (10) a mono-C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub>  
5 alkylamino group, (12) a C<sub>1-6</sub> alkyl-carbonyl group, (13)  
a carboxyl group, (14) a C<sub>1-6</sub> alkoxy-carbonyl group,  
(15) a carbamoyl group, (16) a mono-C<sub>1-6</sub> alkyl-carbamoyl  
group, (17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (18) a C<sub>6-10</sub>  
aryl-carbamoyl group, (19) a sulfo group, (20) a C<sub>1-6</sub>  
10 alkylsulfonyl group, (21) a C<sub>6-10</sub> aryl group, (22) a C<sub>6-10</sub>  
aryloxy group or (23) a 5- to 7-membered heterocyclic  
group having 1 to 3 hetero atoms selected from  
nitrogen, oxygen and sulfur in addition to carbon  
atoms, said heterocyclic group being optionally fused  
15 with a benzene ring, (xvi) a carboxyl group, (xvii) a  
lower alkoxy-carbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl  
such as methoxycarbonyl, ethoxycarbonyl,  
propoxycarbonyl, butoxycarbonyl, etc.), (xviii) a  
formyl group which may be substituted with a 5- to 7-  
20 membered heterocyclic group having 1 to 3 hetero atoms  
selected from nitrogen, oxygen and sulfur in addition  
to carbon atoms, said heterocyclic group being  
optionally fused with a benzene ring, (xix) a carbamoyl  
group, (xx) a mono-lower alkyl-carbamoyl group (e.g.  
25 mono-C<sub>1-6</sub> alkyl-carbamoyl such as methylcarbamoyl,  
ethylcarbamoyl, etc.) whose alkyl portion may be  
substituted with (1) a halogen atom, (2) a C<sub>1-3</sub>  
alkylenedioxy group, (3) a nitro group, (4) a cyano  
group, (5) a C<sub>3-6</sub> cycloalkyl group, (6) a C<sub>1-6</sub> alkoxy  
30 group optionally having 1 to 3 halogen atoms, (7) a C<sub>1-6</sub>  
alkylthio group optionally having 1 to 3 halogen atoms,  
(8) a hydroxyl group, (9) an amino group, (10) a mono-  
C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub> alkylamino group,  
(12) a C<sub>1-6</sub> alkyl-carbonyl group, (13) a carboxyl group,  
35 (14) a C<sub>1-6</sub> alkoxy-carbonyl group, (15) a carbamoyl

group, (16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (18) a C<sub>6-10</sub> aryl-carbamoyl group, (19) a sulfo group, (20) a C<sub>1-6</sub> alkylsulfonyl group, (21) a C<sub>6-10</sub> aryl group, (22) a C<sub>6-10</sub> aryloxy group or (23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (xxi) a di-lower alkyl-carbamoyl group (e.g. di-C<sub>1-6</sub> alkyl-carbamoyl such as dimethylcarbamoyl, diethylcarbamoyl, etc.) whose alkyl portion may be substituted with (1) a halogen atom, (2) a C<sub>1-3</sub> alkylenedioxy group, (3) a nitro group, (4) a cyano group, (5) a C<sub>3-6</sub> cycloalkyl group, (6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (8) a hydroxyl group, (9) an amino group, (10) a mono-C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub> alkylamino group, (12) a C<sub>1-6</sub> alkyl-carbonyl group, (13) a carboxyl group, (14) a C<sub>1-6</sub> alkoxy-carbonyl group, (15) a carbamoyl group, (16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (18) a C<sub>6-10</sub> aryl-carbamoyl group, (19) a sulfo group, (20) a C<sub>1-6</sub> alkylsulfonyl group, (21) a C<sub>6-10</sub> aryl group, (22) a C<sub>6-10</sub> aryloxy group or (23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (xxii) an optionally halogenated aryl-carbamoyl group (e.g. C<sub>6-10</sub> aryl-carbamoyl such as phenylcarbamoyl, naphthylcarbamoyl, etc.), (xxiii) an optionally halogenated aryl-carbonyl group (e.g. C<sub>6-10</sub> aryl-carbonyl such as phenylcarbonyl, naphthylcarbonyl, etc.), (xxiv) a sulfo group optionally substituted with amino group, (xxv) a lower alkylsulfonyl group (e.g.



C<sub>1-6</sub> alkylsulfonyl such as methylsulfonyl, ethylsulfonyl, etc.), (xxvi) an aryl group (e.g. C<sub>6-10</sub> aryl such as phenyl, naphthyl, etc.), (xxvii) an aryloxy group (e.g. C<sub>6-10</sub> aryloxy such as phenyloxy, naphthyloxy, etc.), (xxviii) a C<sub>2-6</sub> alkenylamino, (xxix) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (xxx) a sulfamoyl group, (xxxi) a mono-lower alkyl-sulfamoyl group (e.g. mono-C<sub>1-6</sub> alkyl-sulfamoyl such as methylsulfamoyl, ethylsulfamoyl, etc.), (xxxii) a di-lower alkyl-sulfamoyl group (e.g. di-C<sub>1-6</sub> alkyl-sulfamoyl such as dimethylsulfamoyl, diethylsulfamoyl, etc.), (xxxiii) a lower alkoxy-carbamoyl group (e.g. C<sub>1-6</sub> alkoxy-carbamoyl such as methoxycarbamoyl, ethoxycarbamoyl, etc.), and (xxxiv) a carbamoyloxy group.

The preferred one includes, for example, a lower alkylenedioxy group (e.g. C<sub>1-3</sub> alkylenedioxy such as methylenedioxy, ethylenedioxy, etc.); a nitro group; a cyano group; a C<sub>1-6</sub> alkyl group optionally substituted with (1) a halogen atom, (2) a C<sub>1-3</sub> alkylenedioxy group, (3) a nitro group, (4) a cyano group, (5) a C<sub>1-6</sub> cycloalkyl group, (6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (8) a hydroxyl group, (9) an amino group, (10) a mono-C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub> alkylamino group, (12) a C<sub>1-6</sub> alkyl-carbonyl group, (13) a carboxyl group, (14) a C<sub>1-6</sub> alkoxy-carbonyl group, (15) a carbamoyl group, (16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (18) a C<sub>6-10</sub> aryl-carbamoyl group, (19) a sulfo group, (20) a C<sub>1-6</sub> alkylsulfonyl group, (21) a C<sub>6-10</sub> aryl group, (22) a C<sub>6-10</sub> aryloxy group or (23) a 5-

to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring; a C<sub>3-6</sub> cycloalkyl group; an optionally halogenated lower alkoxy group; an optionally halogenated lower alkylthio group; a hydroxyl group; a C<sub>7-16</sub> aralkyl group; an amino group optionally substituted with a C<sub>1-6</sub> alkyl-carbonyl group; a mono-lower alkylamino group (e.g. mono-C<sub>1-6</sub> alkylamino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.); a di-lower alkylamino group (e.g. di-C<sub>1-6</sub> alkylamino such as dimethylamino, diethylamino, dipropylamino, dibutylamino, etc.); a 5- or 7-membered cyclic amino group optionally having hydroxy or oxo (e.g. morpholino, piperazin-1-yl, piperidino, pyrrolidin-1-yl, 2-pyrrolidon-1-yl, 2-pyridon-1-yl, etc.); a lower alkyl-carbonyl group (e.g. C<sub>1-6</sub> alkyl-carbonyl such as acetyl, propionyl, etc.) whose alkyl portion may be substituted with (1) a halogen atom, (2) a C<sub>1-3</sub> alkylenedioxy group, (3) a nitro group, (4) a cyano group, (5) a C<sub>3-6</sub> cycloalkyl group, (6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (8) a hydroxyl group, (9) an amino group, (10) a mono-C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub> alkylamino group, (12) a C<sub>1-6</sub> alkyl-carbonyl group, (13) a carboxyl group, (14) a C<sub>1-6</sub> alkoxy-carbonyl group, (15) a carbamoyl group, (16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (18) a C<sub>6-10</sub> aryl-carbamoyl group, (19) a sulfo group, (20) a C<sub>1-6</sub> alkylsulfonyl group, (21) a C<sub>6-10</sub> aryl group, (22) a C<sub>6-10</sub> aryloxy group or (23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon

atoms, said heterocyclic group being optionally fused with a benzene ring; a carboxyl group; a lower alkoxy-carbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.); a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring; a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (1) a halogen atom, (2) a C<sub>1-3</sub> alkylenedioxy group, (3) a nitro group, (4) a cyano group, (5) a C<sub>3-6</sub> cycloalkyl group, (6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (8) a hydroxyl group, (9) an amino group, (10) a mono-C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub> alkylamino group, (12) a C<sub>1-6</sub> alkyl-carbonyl group, (13) a carboxyl group, (14) a C<sub>1-6</sub> alkoxy-carbonyl group, (15) a carbamoyl group, (16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (18) a C<sub>6-10</sub> aryl-carbamoyl group, (19) a sulfo group, (20) a C<sub>1-6</sub> alkylsulfonyl group, (21) a C<sub>6-10</sub> aryl group, (22) a C<sub>6-10</sub> aryloxy group or (23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring; an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group; an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group; a sulfo group which may be substituted with amino group; an aryl group (e.g. C<sub>6-10</sub> aryl such as phenyl, naphthyl, etc.); an aryloxy group (e.g. C<sub>6-10</sub> aryloxy such as phenyloxy, naphthyloxy, etc.); a C<sub>2-6</sub> alkenylamino; a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected

from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring; a sulfamoyl group; a mono-lower alkyl-sulfamoyl group (e.g. C<sub>1-6</sub> alkyl-sulfamoyl such as methylsulfamoyl, ethylsulfamoyl, etc.); a di-lower alkyl-sulfamoyl group (e.g. di-C<sub>1-6</sub> alkyl-sulfamoyl such as dimethylsulfamoyl, diethylsulfamoyl, etc.); a lower alkoxy-carbamoyl group (e.g. C<sub>1-6</sub> alkoxy-carbamoyl such as methoxycarbamoyl, ethoxycarbamoyl, etc.); and a carbamoyloxy group.

The more preferred one includes, for example, (i) a C<sub>1-6</sub> alkyl group optionally substituted with (1) a halogen atom, (2) a C<sub>1-3</sub> alkylenedioxy group, (3) a nitro group, (4) a cyano group, (5) a C<sub>3-6</sub> cycloalkyl group, (6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (8) a hydroxyl group, (9) an amino group, (10) a mono-C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub> alkylamino group, (12) a C<sub>1-6</sub> alkyl-carbonyl group, (13) a carboxyl group, (14) a C<sub>1-6</sub> alkoxy-carbonyl group, (15) a carbamoyl group, (16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (18) a C<sub>6-10</sub> aryl-carbamoyl group, (19) a sulfo group, (20) a C<sub>1-6</sub> alkylsulfonyl group, (21) a C<sub>6-10</sub> aryl group, (22) a C<sub>6-10</sub> aryloxy group or (23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (ii) a C<sub>3-6</sub> cycloalkyl group, (iii) an C<sub>7-16</sub> aralkyl group, (iv) a hydroxyl group, (v) an amino group optionally having a C<sub>1-6</sub> alkoxy, (vi) a mono-lower alkylamino group (e.g. mono-C<sub>1-6</sub> alkylamino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc), (vii) a di-lower alkylamino group (e.g. di-C<sub>1-6</sub> alkylamino such

as dimethylamino, diethylamino, dipropylamino, dibutylamino, etc.), (viii) a 5- or 7-membered cyclic amino group optionally having hydroxyl or oxo (e.g. morpholino, piperazin-1-yl, piperidino, pyrrolidin-1-yl, 2-pyrrolidone-1-yl, 2-pyridone-1-yl, etc.), (ix) a lower alkyl-carbonyl group (e.g. C<sub>1-6</sub> alkyl-carbonyl such as acetyl, propionyl, etc.) whose alkyl portion may be substituted with (1) a halogen atom, (2) a C<sub>1-3</sub> alkylenedioxy group, (3) a nitro group, (4) a cyano group, (5) a C<sub>3-6</sub> cycloalkyl group, (6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (8) a hydroxyl group, (9) an amino group, (10) a mono-C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub> alkylamino group, (12) a C<sub>1-6</sub> alkyl-carbonyl group, (13) a carboxyl group, (14) a C<sub>1-6</sub> alkoxy-carbonyl group, (15) a carbamoyl group, (16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (18) a C<sub>6-10</sub> aryl-carbamoyl group, (19) a sulfo group, (20) a C<sub>1-6</sub> alkylsulfonyl group, (21) a C<sub>6-10</sub> aryl group, (22) a C<sub>6-10</sub> aryloxy group or (23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (x) a carboxyl group, (xi) a lower alkoxy-carbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.), (xii) formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (xiii) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (1) a halogen atom, (2) a C<sub>1-3</sub> alkylenedioxy group, (3) a

nitro group, (4) a cyano group, (5) a C<sub>3-6</sub> cycloalkyl group, (6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (8) a hydroxyl group, (9) an amino group, (10) a mono-C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub> alkylamino group, (12) a C<sub>1-6</sub> alkyl-carbonyl group, (13) a carboxyl group, (14) a C<sub>1-6</sub> alkoxy-carbonyl group, (15) a carbamoyl group, (16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (18) a C<sub>6-10</sub> aryl-carbamoyl group, (19) a sulfo group, (20) a C<sub>1-6</sub> alkylsulfonyl group, (21) a C<sub>6-10</sub> aryl group, (22) a C<sub>6-10</sub> aryloxy group or (23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (xiv) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (xv) an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group, (xvi) a sulfo group which may substituted with amino group, (xvii) an aryl group (e.g. C<sub>6-10</sub> aryl such as phenyl, naphthyl, etc.), (xviii) an aryloxy group (e.g. C<sub>6-10</sub> aryloxy such as phenyloxy, naphthyloxy, etc.), (xix) a C<sub>2-6</sub> alkenylamino, (xx) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring; (xxi) a lower alkoxy-carbamoyl group (e.g. C<sub>1-6</sub> alkoxy-carbamoyl such as methoxycarbamoyl, ethoxycarbamoyl, etc.), and (xxii) a carbamoyloxy group.

The "optionally halogenated lower alkoxy group" and "optionally halogenated lower alkylthio group" includes, for example, the same groups as those mentioned for the substituents of the "optionally substituted aromatic group" for Ar<sup>1</sup>, Ar<sup>2</sup> and Ar<sup>3</sup>.

The "heterocyclic group" of the "optionally substituted heterocyclic group" for  $R^3$  and  $R^4$  include, for example, a 5- to 11-membered (cyclic or bicyclic) heterocyclic group having at least one (e.g. 1 to 4, preferably 1 to 3, more preferably 1 or 2) hetero atoms of 1 or 2 kinds selected from a nitrogen atom, an oxygen atom and a sulfur atom in addition to a carbon atom. Examples thereof include 1-, 2- or 3-pyrrolidinyl, 2- or 4-imidazoliny, 2-, 3- or 4-pyrazolidinyl, piperidino, 2-, 3- or 4-piperidyl, 1- or 2-piperazinyl, morpholinyl, non-aromatic heterocyclic group such as 3- or 4-azepinyl (preferably 5- to 7-membered saturated cyclic amino group such as 1- or 2-piperazinyl) and heteroaromatic groups (e.g. 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-furyl, 3-furyl, 4-quinolyl, 8-quinolyl, 4-isoquinolyl, pyrazinyl, 2-pyrimidinyl, 3-pyrrolyl, 2-imidazolyl, 3-pyridazinyl, 3-isothiazolyl, 3-isoxazolyl, i-indolyl, 2-isoindolyl, etc. Among them, a heteroaromatic group or a 5- to 7-membered saturated cyclic amino group is preferred. The more preferred one includes, for example, a 5- or 7-membered heteroaromatic group having 1 to 3 hetero atoms of 1 or 2 kinds selected from a nitrogen atom, an oxygen atom and a sulfur atom in addition to a carbon atom (e.g. 2-thienyl, 3-thienyl, 2-pyridyl, 4-pyridyl, etc.) and a 5- to 7-membered saturated cyclic amino group.

Especially, 2-, 3- or 4-piperidyl, 1- or 2-piperazinyl or morpholinyl is preferred.

The substituent which may substituted on the "optionally substituted heterocyclic group" includes, for example, the same number of the same substituents as mentioned for the "optionally substituted hydrocarbon group" for  $R^4$ .

Preferred examples of  $R^4$  is (i) a hydrogen atom, (ii) a  $C_{1-6}$  alkyl group which may have 1 to 5

- substituents selected from (a) a hydroxyl group, (b) an amino group which may be substituted with a C<sub>1-6</sub> alkyl-carbonyl group, (c) a mono-C<sub>1-6</sub> alkylamino group, (d) a di-C<sub>1-6</sub> alkylamino group, (e) a carboxyl group, (f) a C<sub>1-6</sub> alkoxy-carbonyl group, (g) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (h) a sulfo group which may be substituted with amino group (i) a 5- to 7-membered cyclic amino group which may have an oxo group or which may be substituted with a hydroxyl group, (j) a C<sub>1-6</sub> alkoxy-carbamoyl group, and (k) a carbamoyloxy group.
- (iii) a C<sub>2-6</sub> alkenyl group,  
(iv) a C<sub>6-10</sub> aryl group,  
(v) a 5- to 11-membered heterocyclic groups having at least one hetero atoms of 1 or 2 kinds selected from nitrogen, oxygen and sulfur in addition to a carbon atom, said heterocyclic group being optionally fused with a benzene ring,  
(vi) a C<sub>1-6</sub> alkyl group which may be substituted with a C<sub>1-6</sub> alkyl-carbonyl group,  
(vii) a carboxyl group which may be substituted with a C<sub>1-6</sub> alkyl group,  
(viii) a 5- to 7-membered cyclic amino group which may be substituted with  
(a) a C<sub>1-6</sub> alkyl group optionally substituted with (a-1) a hydroxyl group, (a-2) a di-C<sub>1-6</sub> alkylamino group, (a-3) a C<sub>1-6</sub> alkoxy-carbonyl group or (a-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,  
(b) a C<sub>7-16</sub> aralkyl group, (c) a C<sub>1-6</sub> alkyl-carbonyl group whose alkyl portion may be substituted with (c-1) a halogen atom, (c-2) a mono-C<sub>1-6</sub> alkylamino group, (c-3) a C<sub>1-6</sub> alkoxy-carbonyl group or (c-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms



selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,

(d) a C<sub>1-6</sub> alkoxy-carbonyl group, (e) a formyl group

5 which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,


10 (f) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with a halogen atom or a C<sub>1-6</sub> alkyl-carbonyl group, (g) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (h) an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group or (i) a C<sub>1-6</sub> alkoxy-carbamoyl group, or

15 (ix) a C<sub>6-10</sub> aryloxy group.

More preferred example of R<sup>4</sup> is a group represented by the formula:

(1)


20



or

(2)

25



wherein R<sup>6</sup> and R<sup>7</sup> independently represent (a) a hydrogen atom, (b) a C<sub>1-6</sub> alkyl group optionally substituted with

30 (b-1) a halogen atom, (b-2) a C<sub>1-3</sub> alkylenedioxy group, (b-3) a nitro group, (b-4) a cyano group, (b-5) a C<sub>3-6</sub> cycloalkyl group, (b-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (b-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (b-8) a hydroxyl group, (b-9) an amino group, (b-10) a mono-C<sub>1-6</sub>

35

alkylamino group, (b-11) a di-C<sub>1-6</sub> alkylamino group, (b-12) a C<sub>1-6</sub> alkyl-carbonyl group, (b-13) a carboxyl group, (b-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (b-15) a carbamoyl group, (b-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (b-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (b-18) a C<sub>6-10</sub> aryl-carbamoyl group, (b-19) a sulfo group, (b-20) a C<sub>1-6</sub> alkylsulfonyl group, (b-21) a C<sub>6-10</sub> aryl group, (b-22) a C<sub>6-10</sub> aryloxy group or (b-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (c) a C<sub>3-6</sub> cycloalkyl group, (d) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (e) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (f) a C<sub>7-16</sub> aralkyl group, (g) a hydroxyl group, (h) an amino group, (i) a mono-C<sub>1-6</sub> alkylamino group, (j) a di-C<sub>1-6</sub> alkylamino group, (k) a C<sub>1-6</sub> alkyl-carbonyl group whose alkyl portion may be substituted with (k-1) a halogen atom, (k-2) a C<sub>1-3</sub> alkylenedioxy group, (k-3) a nitro group, (k-4) a cyano group, (k-5) a C<sub>3-6</sub> cycloalkyl group, (k-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (k-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (k-8) a hydroxyl group, (k-9) an amino group, (k-10) a mono-C<sub>1-6</sub> alkylamino group, (k-11) a di-C<sub>1-6</sub> alkylamino group, (k-12) a C<sub>1-6</sub> alkyl-carbonyl group, (k-13) a carboxyl group, (k-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (k-15) a carbamoyl group, (k-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (k-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (k-18) a C<sub>6-10</sub> aryl-carbamoyl group, (k-19) a sulfo group, (k-20) a C<sub>1-6</sub> alkylsulfonyl group, or (k-21) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said

heterocyclic group being optionally fused with a benzene ring, (l) a carboxyl group, (m) a C<sub>1-6</sub> alkoxy-carbonyl group, (n) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (o) a carbamoyl group, (p) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (p-1) a halogen atom, (p-2) a C<sub>1-3</sub> alkylenedioxy group, (p-3) a nitro group, (p-4) a cyano group, (p-5) a C<sub>3-6</sub> cycloalkyl group, (p-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (p-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (p-8) a hydroxyl group, (p-9) an amino group, (p-10) a mono-C<sub>1-6</sub> alkylamino group, (p-11) a di-C<sub>1-6</sub> alkylamino group, (p-12) a C<sub>1-6</sub> alkyl-carbonyl group, (p-13) a carboxyl group, (p-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (p-15) a carbamoyl group, (p-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (p-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (p-18) a C<sub>6-10</sub> aryl-carbamoyl group, (p-19) a sulfo group, (p-20) a C<sub>1-6</sub> alkylsulfonyl group, (p-21) a C<sub>6-10</sub> aryl group, (p-22) a C<sub>6-10</sub> aryloxy group or (p-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (q) a di-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (q-1) a halogen atom, (q-2) a C<sub>1-3</sub> alkylenedioxy group, (q-3) a nitro group, (q-4) a cyano group, (q-5) a C<sub>3-6</sub> cycloalkyl group, (q-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (q-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (q-8) a hydroxyl group, (q-9) an amino group, (q-10) a mono-C<sub>1-6</sub> alkylamino group, (q-11) a di-C<sub>1-6</sub>

alkylamino group, (q-12) a C<sub>1-6</sub> alkyl-carbonyl group, (q-13) a carboxyl group, (q-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (q-15) a carbamoyl group, (q-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (q-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (q-18) a C<sub>6-10</sub> aryl-carbamoyl group, (q-19) a sulfo group, (q-20) a C<sub>1-6</sub> alkylsulfonyl group, (q-21) a C<sub>6-10</sub> aryl group, (q-22) a C<sub>6-10</sub> aryloxy group or (q-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (r) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (s) an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group, (t) a sulfo group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, (w) a C<sub>6-10</sub> aryloxy group, (x) a C<sub>2-6</sub> alkenylamino group or (y) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring.

Preferred example of R<sup>6</sup> and R<sup>7</sup> is, independently, (a) a hydrogen atom, (b) a C<sub>1-6</sub> alkyl group optionally substituted with

(b-1) a hydroxyl group, (b-2) a di-C<sub>1-6</sub> alkylamino group, (b-3) a C<sub>1-6</sub> alkoxy-carbonyl group, or (b-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (c) a C<sub>7-16</sub> aralkyl group, (d) a C<sub>1-6</sub> alkyl-carbonyl group whose alkyl portion may be substituted with (d-1) a halogen atom, (d-2) a mono-C<sub>1-6</sub> alkylamino group, (d-3) a C<sub>1-6</sub> alkoxy-carbonyl group, or (d-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon

atoms, said heterocyclic group being optionally fused with a benzene ring, (e) a C<sub>1-6</sub> alkoxy-carbonyl group, (f) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (g) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (g-1) a halogen atom, or (g-2) a C<sub>1-6</sub> alkyl-carbonyl group, (h) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (i) an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group, or (j) a C<sub>6-10</sub> aryloxy group.

The "lower alkyl group" of the "optionally substituted lower alkyl group" for R<sup>1</sup> is, for example, a straight-chain or branched lower alkyl group having 1 to 6 carbon atoms, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.

The "lower alkyl-carbonyl group" of the "optionally substituted lower alkyl-carbonyl group" for R<sup>1</sup> is, for example, an C<sub>1-6</sub> alkyl-carbonyl group such as methylcarbonyl, ethylcarbonyl, butylcarbonyl, etc.

The substituent which may be present on the "lower alkyl group" and "lower alkyl-carbonyl group" includes, for example, the same substituents as mentioned for the "optionally substituted hydrocarbon group" for R<sup>2</sup>.

Preferred examples of R<sup>1</sup> include a hydrogen atom or a lower alkyl group (e.g. C<sub>1-6</sub> alkyl such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.). Among them, a hydrogen atom and methyl are particularly preferred.

Especially preferred for R<sup>1</sup> is a hydrogen atom.

The "nitrogen-containing heterocyclic group formed by bonding R<sup>1</sup> and R<sup>2</sup> together with adjacent nitrogen" is, for example, a 4- to 8-membered ring optionally

having at least one nitrogen atom and 1 to 3  
(preferably 1 to 2) ring-constituting atoms such as an  
oxygen atom, a sulfur atom, etc. in addition to a  
carbon atom, or the 4- to 8-membered ring fused with a  
5 benzene ring.

Examples thereof include an aromatic heterocyclic  
group (e.g. 1-pyrrolyl, 1-imidazolyl, 1-indolyl,  
1-pyrazolyl, 2-isoindolyl, 1-indazolyl, etc.), a cyclic  
amino group (e.g. morpholino, piperidino,  
10 1-piperaziny, 1-pyrrolidinyl, 1-pirazolidinyl,  
1-azepinyl, etc.) or the cyclic amino group fused with  
a benzene ring (e.g. 1-indolinyl, 2-isoindolinyl,  
1,2,3,4-tetrahydroquinolin-1-yl, 1,2,3,4-tetrahydro-  
isoquinolin-2-yl, 3-benzazepin-3-yl, etc.) or a lactam  
15 or an imide group (e.g. phthalimide, succinimide,  
2-pyrrolidon-1-yl, 2-pyridon-1-yl, 2-quinolon-1-yl,  
etc.).

The "nitrogen-containing heterocyclic group formed  
by bonding  $R^1$  and  $R^2$  together with adjacent nitrogen"  
20 may have the same substituent as that may be present on  
the "optionally substituted hydrocarbon group" for  $R^2$ .  
The group fused with a benzene ring may have one or  
plurality (preferably 1 to 5, more preferably 1 to 3,  
further more preferably 1 or 2) of substituents  
25 selected from a halogen group (e.g. fluorine, chlorine,  
bromine, iodine, etc.), a lower alkylenedioxy group  
(e.g.  $C_{1-3}$  alkylenedioxy such as methylenedioxy,  
ethylenedioxy, etc.), a nitro group, a cyano group, an  
optionally halogenated lower alkyl group, an optionally  
30 halogenated lower alkoxy group, an optionally  
halogenated lower alkylthio group, a hydroxyl group, an  
amino group, a mono-lower alkylamino group (e.g.  
mono- $C_{1-6}$  alkylamino such as methylamino, ethylamino,  
propylamino, isopropylamino, butylamino, etc.), a  
35 di-lower alkylamino group (e.g. di- $C_{1-6}$  alkylamino such  
as dimethylamino, diethylamino, dipropylamino,

dibutylamino, etc.), a carboxyl group, a lower alkocycarbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.) or a carbamoyl group in any position on the benzene ring.

The "optionally halogenated lower alkyl group," "optionally halogenated lower alkoxy group" and "optionally halogenated lower alkylthio group" include the same groups as mentioned for the substituents of the "optionally substituted aromatic group" for Ar<sup>1</sup>, Ar<sup>2</sup> and Ar<sup>3</sup>.

As the "nitrogen-containing heterocyclic group" of the "nitrogen-containing heterocyclic group formed by bonding R<sup>1</sup> and R<sup>2</sup> together with adjacent nitrogen," "1-piperazinyl" is preferred. The "1-piperazinyl" having a substituent on a nitrogen atom at the 4-position is preferred.

The preferred substituent on the nitrogen atom at the 4-position of the "1-piperazinyl" includes, for example, a lower alkyl group (e.g. C<sub>1-6</sub> alkyl such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.), aryl (e.g. C<sub>6-14</sub> aryl such as phenyl, 1-naphthyl, 2-naphthyl, biphenyl, 2-indenyl, 2-anthryl, etc., preferably phenyl), 2-pyridyl, 3-pyridyl, 4-pyridyl, an aralkyl group (e.g. C<sub>7-16</sub> aralkyl such as benzyl, phenethyl, diphenylmethyl, 1-naphthylmethyl, 2-naphthylmethyl, 2-diphenylethyl, 3-phenylpropyl, 4-phenylbutyl, 5-phenylpentyl, etc., preferably benzyl), a phenacyl group or a nicotinoyl group.

Furthermore, an aryl group, an aralkyl group, a phenacyl group and a nicotinoyl group may have one or plurality (preferably 1 to 5, more preferably 1 to 3, further more preferably 1 or 2) of substituents selected from a halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), a lower alkylenedioxy group

(e.g. C<sub>1-3</sub> alkylenedioxy such as methylenedioxy, ethylenedioxy, etc.), a nitro group, a cyano group, optionally halogenated lower alkyl group, an optionally halogenated lower alkoxy group, an optionally halogenated lower alkylthio group, a hydroxyl group, an amino group, a mono-lower alkylamino group (e.g. mono-C<sub>1-6</sub> alkylamino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), a di-lower alkylamino group (e.g. di-C<sub>1-6</sub> alkylamino such as dimethylamino, diethylamino, dipropylamino, dibutylamino, etc.), a carboxyl group, a lower alkoxy-carbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.) or a carbamoyl group in any position on the benzene ring.

The "optionally halogenated lower alkyl group," "optionally halogenated lower alkoxy group" and "optionally halogenated lower alkylthio group" include the same groups mentioned for the substituents of the "optionally substituted aromatic group" for Ar<sup>1</sup>, Ar<sup>2</sup> and Ar<sup>3</sup>.

The term "divalent aliphatic hydrocarbon group" of the "optionally substituted divalent aliphatic hydrocarbon group optionally having oxygen or sulfur in the carbon chain" for Q<sup>1</sup> and Q<sup>2</sup> means a group obtained by eliminating each one hydrogen (two hydrogens in total) bound to the same or different carbon atoms from the saturated or unsaturated aliphatic hydrocarbon and preferably have not more than 6 carbon atoms. Specific examples thereof include the following:

- (i) a C<sub>1-6</sub> alkylene group (e.g. -CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, -(CH<sub>2</sub>)<sub>4</sub>-, -(CH<sub>2</sub>)<sub>5</sub>-, -(CH<sub>2</sub>)<sub>6</sub>-, etc.)
- (ii) a C<sub>2-6</sub> alkenylene group (e.g. -CH=CH-, -CH=CH-CH<sub>2</sub>-, -CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>-CH=CH-CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>-CH=CH-(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-CH=CH-CH<sub>2</sub>-, etc.)



(iii) a C<sub>2-6</sub> alkynylene group (e.g.  
-C≡C-, -C≡C-CH<sub>2</sub>-, -CH<sub>2</sub>-C≡C-CH<sub>2</sub>-,  
-(CH<sub>2</sub>)<sub>2</sub>-C≡C-CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>-C≡C-(CH<sub>2</sub>)<sub>2</sub>-,  
-(CH<sub>2</sub>)<sub>3</sub>-C≡C-CH<sub>2</sub>-, etc).

5

Preferred one is a C<sub>1-6</sub> alkylene group and particularly preferred one is a C<sub>1-3</sub> alkylene group.

These groups may have an oxygen atom or an optionally oxidized sulfur atom in the carbon atom, or  
10 any carbon atom may be substituted with an oxo group or a thioxo group in the carbon chain.

For example, a group represented by the formula  
-(CH<sub>2</sub>)<sub>a</sub>-T-(CH<sub>2</sub>)<sub>m</sub>- [wherein T is a bond, an oxygen atom or an optionally oxidized sulfur atom; and a and m  
15 independently represent an integer of 0 to 5 and the total of them is 1 to 6].

Preferred examples of Q<sup>1</sup> and Q<sup>2</sup> is a C<sub>1-6</sub> alkylene group optionally having an oxo group, for example,  
-CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, -(CH<sub>2</sub>)<sub>4</sub>-, -(CH<sub>2</sub>)<sub>2</sub>CO-, -CH<sub>2</sub>CO-,  
20 -CO-, etc.


Preferred examples of Q<sup>1</sup> is a C<sub>1-4</sub> alkylene group optionally having an oxo group, for example, -CH<sub>2</sub>-,  
-(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, -(CH<sub>2</sub>)<sub>4</sub>-, -(CH<sub>2</sub>)<sub>2</sub>CO- and -CH<sub>2</sub>CO-. Particularly preferred are -CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-,  
25 -(CH<sub>2</sub>)<sub>2</sub>CO- and -CH<sub>2</sub>CO-. Among them, -(CH<sub>2</sub>)<sub>3</sub>- is commonly used.

Preferred examples of Q<sup>2</sup> are -CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, -(CH<sub>2</sub>)<sub>4</sub>-, -(CH<sub>2</sub>)<sub>2</sub>CO-, -CH<sub>2</sub>CO- and -CO-. Particularly preferred are -CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>- and -(CH<sub>2</sub>)<sub>3</sub>-.  
30 Among them, -CH<sub>2</sub>- is commonly used.

The divalent aliphatic hydrocarbon group may have ether oxygen or sulfur in the carbon chain, and examples thereof include -CH<sub>2</sub>-O-CH<sub>2</sub>-, -CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>-,  
-CH<sub>2</sub>-CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>-CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>-,  
35 (CH<sub>2</sub>)<sub>2</sub>-CH<sub>2</sub>-O-CH<sub>2</sub>-(CH<sub>2</sub>)<sub>2</sub>-, (CH<sub>2</sub>)<sub>3</sub>-CH<sub>2</sub>-O-CH<sub>2</sub>-CH<sub>2</sub>-,

-CH<sub>2</sub>-S-CH<sub>2</sub>-, -CH<sub>2</sub>-S-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-S-CH<sub>2</sub>-CH<sub>2</sub>-,  
 -(CH<sub>2</sub>)<sub>2</sub>-CH<sub>2</sub>-S-CH<sub>2</sub>-CH<sub>2</sub>-, (CH<sub>2</sub>)<sub>2</sub>-CH<sub>2</sub>-S-CH<sub>2</sub>-(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-  
 CH<sub>2</sub>-S-CH<sub>2</sub>-CH<sub>2</sub>-, etc. A sulfur atom may be sulfoxide or  
 sulfon.

- 5 The "optionally substituted monocyclic or fused  
 nitrogen-containing heterocyclic ring" represented by  
 a group of the formula:




10

may have 1 or 2 unsaturated bonds and represents a  
 monocyclic 4- to 9-membered ring or bicyclic 6- to  
 14-membered ring optionally having 1 or 2 substituents  
 in any position other than N and Z.

- 15 The preferred "monocyclic nitrogen-containing  
 heterocyclic ring" of the "optionally substituted  
 monocyclic nitrogen-containing heterocyclic ring"  
 includes, for example, the following:


20



25

(wherein Z has the same meanings as defined above;  
 and ..... represents a single bond or a double bond).  
 Among them,

30



is preferred.

Especially, is preferred.

35

These monocyclic nitrogen-containing heterocyclic

ring may be fused with a 3- to 10-membered cyclic hydrocarbon group, for example, a lower cycloalkane group (e.g. C<sub>3-8</sub> cycloalkane such as cyclopropane, cyclobutane, cyclopentane, cyclohexane, etc.), a lower  
5 cycloalkene group (e.g. C<sub>3-6</sub> cycloalkene such as cyclopropene, cyclopentene, cyclohexene, etc.) or an aryl group (e.g. C<sub>6-10</sub> aryl such as benzene, etc.) to form a bicyclic 6- to 14-membered nitrogen-containing heterocycle. Among them, pyrrolidine, piperidine,  
10 azepine or one of these three groups fused with a benzene ring are preferred. Particularly preferred is piperidine.

Examples of the substituent which may present on the monocyclic or fused nitrogen-containing  
15 heterocyclic ring include an optionally substituted lower alkyl group (e.g. C<sub>1-6</sub> alkyl such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.), an optionally substituted lower alkoxy group (e.g. C<sub>1-6</sub> alkoxy such as  
20 methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, etc.), an optionally substituted lower alkylthio group (e.g. C<sub>1-6</sub> alkylthio such as methylthio, ethylthio, propylthio, isopropylthio, n-butylthio, isobutylthio,  
25 sec-butylthio, tert-butylthio, etc.), a hydroxyl group, an amino group, a mono-lower alkylamino group (e.g. mono-C<sub>1-6</sub> alkylamino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), a di-lower alkylamino group (e.g. di-C<sub>1-6</sub> alkylamino such  
30 as dimethylamino, diethylamino, dipropylamino, dibutylamino, etc.), a lower alkyl-carbonyl group (e.g. C<sub>1-6</sub> alkyl-carbonyl such as acetyl, propionyl, etc.), a carboxyl group, a lower alkoxy-carbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl such as methoxycarbonyl,  
35 ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.),

a carbamoyl group, a mono-lower alkyl-carbamoyl group (e.g. mono-C<sub>1-6</sub> alkyl-carbamoyl such as methylcarbamoyl, ethylcarbamoyl, etc.), a di-lower alkyl-carbamoyl group (e.g. di-C<sub>1-6</sub> alkyl-carbamoyl such as dimethylcarbamoyl, diethylcarbamoyl, etc.), an aryl-carbamoyl group (e.g. C<sub>6-10</sub> aryl-carbamoyl such as phenylcarbamoyl, naphthylcarbamoyl, etc.), a sulfo group, a lower alkylsulfonyl group (e.g. C<sub>1-6</sub> alkylsulfonyl such as methylsulfonyl, ethylsulfonyl, etc.), an aryl group (C<sub>6-10</sub> aryl such as phenyl, naphthyl, etc.) or an aryloxy group (e.g. C<sub>6-10</sub> aryloxy such as phenyloxy, naphthyloxy, etc.).


The substituent which may present on the "optionally substituted lower alkyl group," "optionally substituted lower alkoxy group" and "optionally substituted lower alkylthio group" include, for examples, a lower alkoxy group (e.g. C<sub>1-6</sub> alkoxy such as methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, etc.), a lower alkylthio group (e.g. C<sub>1-6</sub> alkylthio such as methylthio, ethylthio, propylthio, isopropylthio, n-butylthio, isobutylthio, sec-butylthio, tert-butylthio, etc.), a hydroxyl group, an amino group, a mono-lower alkylamino group (e.g. mono-C<sub>1-6</sub> alkylamino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), a di-lower alkylamino group (e.g. di-C<sub>1-6</sub> alkylamino such as dimethylamino, diethylamino, dipropylamino, dibutylamino, etc.), a lower alkyl-carbonyl group (e.g. C<sub>1-6</sub> alkyl-carbonyl such as acetyl, propionyl, etc.), a carboxyl group, a lower alkoxy-carbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.), a carbamoyl group, a mono-lower alkyl-carbamoyl group (e.g. mono-C<sub>1-6</sub> alkyl-carbamoyl such as methylcarbamoyl, ethylcarbamoyl, etc.), a

di-lower alkyl-carbamoyl group (e.g. di-C<sub>1-6</sub> alkyl-carbamoyl such as dimethylcarbamoyl, diethylcarbamoyl, etc.), an aryl-carbamoyl group (e.g. C<sub>6-10</sub> aryl-carbamoyl such as phenylcarbamoyl, naphthylcarbamoyl, etc.), a sulfo group, an alkylsulfonyl group (e.g. C<sub>1-6</sub> alkylsulfonyl such as methylsulfonyl, ethylsulfonyl, etc.), an aryl group (C<sub>6-10</sub> aryl such as phenyl, naphthyl, etc.) or an aryloxy group (e.g. C<sub>6-10</sub> aryloxy such as phenyloxy, naphthyloxy, etc.).

2 is, for example, the following:


[1] An optionally substituted 1, 2-phenylene,

[2] A group of the formula:



[wherein Ar<sup>3</sup> has the same meanings as defined above; and n is an integer of 0 to 3],


[3] A group of the formula:



[wherein Ar<sup>3</sup> and n have the same meanings as defined above; and Y is an hydrogen atom, an optionally halogenated lower alkyl group, an optionally halogenated lower alkoxy group, an optionally halogenated lower alkylthio group, a hydroxyl group, a cyano group, an alkyl-carbonyl group (e.g. C<sub>1-6</sub> alkyl-carbonyl such as acetyl, propionyl, etc.), a lower alkyl-carbonyloxy group (e.g. C<sub>1-6</sub> alkyl-carbonyloxy such as acetyloxy, propionyloxy, etc.), a formylamino group, an amino group, a mono-lower alkylamino group (e.g. mono-C<sub>1-6</sub> alkylamino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), a di-lower alkylamino group (e.g. di-C<sub>1-6</sub> alkylamino such as dimethylamino, diethylamino,


dipropylamino, dibutylamino, etc.), a carboxyl group, a lower alkoxy-carbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.) or a lower alkyl-carbonylamino group (e.g. C<sub>1-6</sub> alkyl-carbonylamino such as acetylamino, propionylamino, etc.) ("optionally halogenated lower alkyl," "optionally halogenated lower alkoxy" and "optionally halogenated lower alkylthio" have the same meanings as mentioned for the substituents of the "optionally substituted aromatic group" for Ar<sup>3</sup>) (Preferred examples of Y include hydrogen atom, a hydroxyl group, a cyano group, a C<sub>1-6</sub> alkoxy group, an amino group and a mono-C<sub>1-6</sub> alkylamino group and, among them, a hydrogen group, a hydroxyl group, an amino group and a mono-C<sub>1-6</sub> alkylamino group are preferred. Particularly preferred are a hydrogen atom and a hydroxyl group. A hydroxyl group is commonly used.)

[4] A group of the formula:



[wherein Ar<sup>3</sup> and n have the same meanings as defined above.], or


[5] A group of the formula:



[wherein Ar<sup>3</sup> and n have the same meanings as defined above.]

Preferred example of n is an integer of 0 to 2. More preferred is 0 or 1. Among them, 0 is particularly preferred.

Among them, preferred example of Z include a group of the formula:




[wherein  $\text{Ar}^3$  and  $n$  have the same meanings as  
 5 defined above; and  $\text{Y}$  is a hydrogen atom or a hydroxyl  
 group, preferably a hydroxyl group].

In the case that  $\text{Z}$  is a 1,2-phenylene group,  
 examples of the ring represented by the formula:




include the following:




20

Among them,



is preferred.

30 In the case that  $\text{Z}$  is a group represented by the  
 formula:




35 [wherein  $\text{Ar}^3$  and  $n$  have the same meanings as  
 defined above; and  $\text{Y}$  is a hydrogen atom or a hydroxyl,

preferably a hydroxyl group}, the most preferred examples of the ring represented by the formula:



include a group represented by the formula:



[wherein  $\text{Ar}^3$  has the same meanings as defined above].

The substituent which may be present on the "1,2-phenylene" includes, for example, the same substituents as mentioned for the substituents of the "optionally substituted aromatic group". Preferred examples thereof include a halogen atom (particularly preferably fluorine, chlorine), a lower alkylendioxy group (e.g.  $\text{C}_{1-3}$  alkylendioxy such as methylenedioxy, ethylenedioxy, etc.), a nitro group, a cyano group, an optionally halogenated lower alkyl group or an optionally halogenated lower alkoxy group.

The "optionally halogenated lower alkyl group" and "optionally halogenated lower alkoxy group" include the same groups as mentioned for the substituents of the "optionally substituted aromatic group" for  $\text{Ar}^1$ ,  $\text{Ar}^2$  and  $\text{Ar}^3$ .

Preferred compound (I) or a salt thereof is one wherein  $\text{Q}^1$  is  $-\text{CH}_2-$ ,  $-(\text{CH}_2)_2-$ ,  $-(\text{CH}_2)_3-$ ,  $-(\text{CH}_2)_4-$  or  $-(\text{CH}_2)_2\text{CO}-$ ;


$\text{Q}^2$  is  $-\text{CH}_2-$ ,  $-(\text{CH}_2)_2-$ ,  $-(\text{CH}_2)_3-$ ,  $-(\text{CH}_2)_4-$ ,  $-\text{CO}-$ ,  $-\text{CH}_2\text{CO}-$  or  $-(\text{CH}_2)_2\text{CO}-$ ;

$\text{Ar}^1$  and  $\text{Ar}^2$  independently represent phenyl, 4-chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl or




4-pyridyl;

a group of the formula:




5

is



10

wherein Z is a group of the formula




15

[wherein  $\text{Ar}^3$  is a  $\text{C}_{1-3}$  alkyl group optionally substituted with 1 to 3 halogen atoms, a  $\text{C}_{1-3}$  alkoxy group substituted with 1 to 3 halogen atoms or a phenyl group optionally substituted with a halogen atom (preferably chlorine, fluorine) (e.g. phenyl, 4-chlorophenyl, 4-fluorophenyl, 4-methoxyphenyl, 3,5-dichlorophenyl, 3,5-difluorophenyl, 4-trifluoromethylphenyl, etc.), 2-pyridyl, 3-pyridyl or 4-pyridyl; and n is an integer of 0 to 3],


20

25



[wherein  $\text{Ar}^3$  and n have the same meanings as defined above; and Y is a hydrogen atom, a hydroxyl group, an amino group or a mono- $\text{C}_{1-6}$  alkylamino group (particularly a hydrogen atom and a hydroxyl group are preferred)] or

30



35

[wherein  $\text{Ar}^3$  has the same meanings as defined above];

$R^1$  is a hydrogen atom or methyl;

$R^2$  is (1) an  $C_{1-6}$  alkyl group which may be substituted with a  $C_{1-6}$  alkoxy-carbonyl group, a carboxyl group, a  $C_{1-6}$  alkyl-carbonyl group or a formyl group, or (2) an acyl group represented by  $-(C=O)-R^4$ ,  
5  $-SO_2-R^4$ ,  $-SO-R^4$ ,  $-(C=O)NR^5R^4$  or  $-(C=O)O-R^4$ ;

$R^3$  is a hydrogen atom or a  $C_{1-3}$  alkyl group such as methyl, ethyl, propyl, isopropyl, etc.; and

$R^4$  is a hydrogen atom, a lower alkyl group (e.g.  $C_{1-6}$  alkyl such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.), a lower alkenyl group (e.g.  $C_{2-6}$  alkenyl such as vinyl, allyl, isopropenyl, etc.), a lower alkyl-carbonyl group (e.g.  $C_{1-6}$  alkyl-carbonyl such as acetyl, propionyl, butyryl, etc.), a carboxyl group, a lower  
10 alkoxy-carbonyl group (e.g.  $C_{1-6}$  alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.), a mono-lower alkylaminocarbonyl group (e.g. mono- $C_{1-6}$  alkylaminocarbonyl such as methylaminocarbonyl, ethylaminocarbonyl,  
15 propylaminocarbonyl, butylaminocarbonyl, etc.), a di-lower alkylaminocarbonyl group (e.g. di- $C_{1-6}$  alkylaminocarbonyl such as dimethylaminocarbonyl, diethylaminocarbonyl, dipropylaminocarbonyl, dibutylaminocarbonyl, etc.), a  $C_{6-10}$  aryl group  
20 (preferably phenyl) or a 5- to 7-membered cyclic amino group (preferably 2-piperidyl, 3-piperidyl, 4-piperidyl, 1-pyrrolidinyl, 3-pyrrolidinyl, 2-piperazinyl, etc.).

30 The "lower alkyl group," "lower alkenyl group," "lower alkyl-carbonyl group," "carboxyl group," "lower alkoxy-carbonyl group," "mono-lower alkylaminocarbonyl group," "di-lower alkylaminocarbonyl group" and "5- to 7-membered cyclic amino group" for  $R^4$  may have 1 to 3  
35 substituents on any carbon atom. The substituent

include, for example, (i) a halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), (ii) a lower alkylenedioxy group (e.g. C<sub>1-3</sub> alkylenedioxy such as methylenedioxy, ethylenedioxy, etc.), (iii) a nitro group, (iv) a cyano group, (v) a C<sub>1-6</sub> alkyl group optionally substituted with (1) a halogen atom, (2) a C<sub>1-3</sub> alkylenedioxy group, (3) a nitro group, (4) a cyano group, (5) a C<sub>3-6</sub> cycloalkyl group, (6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (8) a hydroxyl group, (9) an amino group, (10) a mono-C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub> alkylamino group, (12) a C<sub>1-6</sub> alkyl-carbonyl group, (13) a carboxyl group, (14) a C<sub>1-6</sub> alkoxy-carbonyl group, (15) a carbamoyl group, (16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (18) a C<sub>6-10</sub> aryl-carbamoyl group, (19) a sulfo group, (20) a C<sub>1-6</sub> alkylsulfonyl group, (21) a C<sub>6-10</sub> aryl group, (22) a C<sub>6-10</sub> aryloxy group or (23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (vi) a C<sub>3-6</sub> cycloalkyl group, (vii) an optionally halogenated lower alkoxy group (e.g. optionally halogenated C<sub>1-6</sub> alkoxy such as methoxy, difluoromethoxy, trifluoromethoxy, ethoxy, 2,2,2-trifluoromethoxy, propoxy, isopropoxy, butoxy, 4,4,4-trifluorobutoxy, isobutoxy, sec-butoxy, pentyloxy, hexyloxy, etc.), (viii) an optionally halogenated lower alkylthio group (e.g. optionally halogenated C<sub>1-6</sub> alkylthio such as methylthio, difluoromethylthio, trifluoromethylthio, ethylthio, propylthio, isopropylthio, butylthio, 4,4,4-trifluorobutylthio, pentylthio, hexylthio, etc.), (ix) a C<sub>7-16</sub> aralkyl group, (x) a hydroxyl group, (xi)

an amino group, (xii) a mono-lower alkylamino group (e.g. mono-C<sub>1-6</sub> alkylamino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), (xiii) a di-lower alkylamino group (e.g. di-C<sub>1-6</sub> alkylamino such as dimethylamino, diethylamino, dipropylamino, dibutylamino, etc.), (xiv) 5- to 7-membered cyclic amino group optionally having a hydroxyl group or an oxo group (e.g. morpholino, piperazin-1-yl, piperidino, pyrrolidin-1-yl, 2-pyrrolidone-1-yl, 2-pyridone-1-yl, etc.), (xv) a lower alkyl-carbonyl group (C<sub>1-6</sub> alkyl-carbonyl such as acetyl, propionyl, etc.), whose alkyl portion may be substituted with (1) a halogen atom, (2) a C<sub>1-3</sub> alkylenedioxy group, (3) a nitro group, (4) a cyano group, (5) a C<sub>3-6</sub> cycloalkyl group, (6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (8) a hydroxyl group, (9) an amino group, (10) a mono-C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub> alkylamino group, (12) a C<sub>1-6</sub> alkyl-carbonyl group, (13) a carboxyl group, (14) a C<sub>1-6</sub> alkoxy-carbonyl group, (15) a carbamoyl group, (16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (18) a C<sub>6-10</sub> aryl-carbamoyl group, (19) a sulfo group, (20) a C<sub>1-6</sub> alkylsulfonyl group, (21) a C<sub>6-10</sub> aryl group, (22) a C<sub>6-10</sub> aryloxy group or (23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (xvi) a carboxyl group, (xvii) a lower alkoxy-carbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.), (xviii) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms


selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (xix) a carbamoyl group, (xx) a mono-lower alkyl-carbamoyl group (e.g. mono-C<sub>1-6</sub> alkyl-carbamoyl such as methylcarbamoyl, ethylcarbamoyl, etc.) whose alkyl portion may be substituted with (1) a halogen atom, (2) a C<sub>1-3</sub> alkylenedioxy group, (3) a nitro group, (4) a cyano group, (5) a C<sub>3-6</sub> cycloalkyl group, (6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (8) a hydroxyl group, (9) an amino group, (10) a mono-C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub> alkylamino group, (12) a C<sub>1-6</sub> alkyl-carbonyl group, (13) a carboxyl group, (14) a C<sub>1-6</sub> alkoxy-carbonyl group, (15) a carbamoyl group, (16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (18) a C<sub>6-10</sub> aryl-carbamoyl group, (19) a sulfo group, (20) a C<sub>1-6</sub> alkylsulfonyl group, (21) a C<sub>6-10</sub> aryl group, (22) a C<sub>6-10</sub> aryloxy group or (23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (xxi) a di-lower alkyl-carbamoyl group (e.g. di-C<sub>1-6</sub> alkyl-carbamoyl such as dimethylcarbamoyl, diethylcarbamoyl, etc.) whose alkyl portion may be substituted with (1) a halogen atom, (2) a C<sub>1-3</sub> alkylenedioxy group, (3) a nitro group, (4) a cyano group, (5) a C<sub>3-6</sub> cycloalkyl group, (6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (8) a hydroxyl group, (9) an amino group, (10) a mono-C<sub>1-6</sub> alkylamino group, (11) a di-C<sub>1-6</sub> alkylamino group, (12) a C<sub>1-6</sub> alkyl-carbonyl group, (13) a carboxyl group, (14) a C<sub>1-6</sub> alkoxy-carbonyl group,


(15) a carbamoyl group, (16) a mono- $C_{1-6}$  alkyl-carbamoyl group, (17) a di- $C_{1-6}$  alkyl-carbamoyl group, (18) a  $C_{6-10}$  aryl-carbamoyl group, (19) a sulfo group, (20) a  $C_{1-6}$  alkylsulfonyl group, (21) a  $C_{6-10}$  aryl group, (22) a  $C_{6-10}$  aryloxy group or (23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (xxii) an aryl-carbamoyl group (e.g.  $C_{6-10}$  aryl-carbamoyl such as phenylcarbamoyl, naphthylcarbamoyl, etc.), (xxiii) a sulfo group, (xxiv) a lower alkylsulfonyl group (e.g.  $C_{1-6}$  alkylsulfonyl such as methylsulfonyl, ethylsulfonyl, etc.), (xxv) an aryl group ( $C_{6-10}$  aryl such as phenyl, naphthyl, etc.), (xxvi) an aryloxy group (e.g.  $C_{6-10}$  aryloxy such as phenyloxy, naphthyloxy, etc.), (xxvii) a sulfamoyl group, (xxviii) a mono-lower alkyl-sulfamoyl group (e.g.  $C_{1-6}$  alkyl-sulfamoyl such as methylsulfamoyl, ethylsulfamoyl, etc.), (xxix) a di-lower alkyl-sulfamoyl group (e.g. di- $C_{1-6}$  alkyl-sulfamoyl such as dimethylsulfamoyl, diethylsulfamoyl, etc.), (xxx) a lower alkoxy-carbamoyl group (e.g.  $C_{1-6}$  alkoxy-carbamoyl such as methoxycarbamoyl, ethoxycarbamoyl, etc.), and (xxxi) a carbamoyloxy group.

More preferred is a compound wherein  $Q^1$  is  $-CH_2-$ ,  $-(CH_2)_2-$  or  $-(CH_2)_3-$ ;


$Q^2$  is  $-CH_2-$ ,  $-(CH_2)_2-$ ,  $-(CH_2)_3-$ ,  $-CH_2CO-$  or  $-(CH_2)_2CO-$ ;

$Ar^1$  and  $Ar^2$  independently represent phenyl or 2-pyridyl;  
a group of the formula:






5            wherein Z is  
a group of the formula:




10            [wherein  $Ar^3$  is a phenyl group optionally  
substituted with 1 to 3 (preferably 1 or 2) halogen  
atoms (preferably chlorine, fluorine) (e.g. phenyl,  
4-chlorophenyl, 4-fluorophenyl, 3,5-dichlorophenyl,  
3,5-difluorophenyl, etc.) or 2-pyridyl; and n  
represents 0];

15



20            [wherein  $Ar^3$  and n have the same meanings as  
defined above; and Y is a hydrogen atom or a hydroxyl  
group] or



25            [wherein  $Ar^3$  and n have the same meanings as  
defined above];

$R^1$  is a hydrogen atom or methyl;

$R^2$  is an acyl group represented by  $-(C=O)-R^4$ ,  
 $-(C=O)NR^3-R^4$  or  $-(C=O)O-R^4$ ;

$R^5$  is a hydrogen atom; and

30             $R^4$  is a hydrogen atom, an optionally substituted  
lower alkyl group (e.g.  $C_{1-6}$  alkyl such as methyl,  
ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl,  
tert-butyl, pentyl, hexyl, etc.), a carboxyl group, a  
lower alkoxy-carbonyl group (e.g.  $C_{1-6}$  alkoxy-carbonyl  
35            such as methoxycarbonyl, ethoxycarbonyl,

propoxycarbonyl, butoxycarbonyl, etc.), a phenyl group or 1-piperazinyl.

The "lower alkyl group" for R<sup>4</sup> may have 1 substituent on any carbon atom. The substituent include, for example, a hydroxyl group, an amino group, a di-lower alkylamino group (e.g. di-C<sub>1-6</sub> alkylamino such as dimethylamino, diethylamino, dipropylamino, dibutylamino, etc.), a 5- to 7-membered cyclic amino group optionally having a hydroxyl group or an oxo group (e.g. morpholino, piperazin-1-yl, piperidino, pyrrolidin-1-yl, 2-pyrrolidone-1-yl, 2-pyridone-1-yl, etc.), a lower alkyl-carbonyl group (C<sub>1-6</sub> alkyl-carbonyl such as acetyl, propionyl, etc.), a carboxyl group, a lower alkoxy-carbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.), a sulfamoyl group, a mono-lower alkyl-sulfamoyl group (e.g. mono-C<sub>1-6</sub> alkyl-sulfamoyl such as methylsulfamoyl, ethylsulfamoyl, etc.) or a di-lower alkyl-sulfamoyl group (e.g. di-C<sub>1-6</sub> alkyl-sulfamoyl such as dimethylsulfamoyl, diethylsulfamoyl, etc.).


Particularly preferred is a compound wherein Q<sup>1</sup> is -(CH<sub>2</sub>)<sub>3</sub>-;

Q<sup>2</sup> is -CH<sub>2</sub>- or -(CH<sub>2</sub>)<sub>2</sub>-;

Ar<sup>1</sup> is a phenyl group or 2-pyridyl;

Ar<sup>2</sup> is a phenyl group;

a group of the formula:



30

is


a group of the formula:



35



wherein Z is  
a group of the formula:



5 [wherein Ar<sup>3</sup> is 4-chlorophenyl; n is 0; and Y is  
hydrogen atom or a hydroxyl group];

R<sup>1</sup> is a hydrogen atom;


R<sup>2</sup> is an acyl group represented by -(C=O)-R<sup>4</sup>,  
-(C=O)NR<sup>5</sup>-R<sup>4</sup> or -(C=O)O-R<sup>4</sup>;

10 R<sup>5</sup> is a hydrogen atom; and


R<sup>4</sup> is a (1) hydrogen atom or (2) a lower alkyl  
group (C<sub>1-6</sub> alkyl such as methyl, ethyl, propyl,  
isopropyl, butyl, isobutyl, sec-butyl, tert-butyl,  
pentyl, hexyl, etc.) optionally having one substituent  
15 selected from (a) a hydroxyl group, (b) a 5- to  
7-membered cyclic amino group optionally having (b-1) a  
hydroxyl group or (b-2) an oxo group (e.g. morpholino,  
piperazin-1-yl, piperidino, pyrrolidin-1-yl,  
2-pyrrolidone-1-yl, 2-pyridone-1-yl, etc.) or (c) a  
20 sulfamoyl group.

In addition, preferred compound (I) is one wherein  
Ar<sup>1</sup> and Ar<sup>2</sup> independently represent, phenyl, 4-  
chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl, or  
4-pyridyl;


25 Q<sup>1</sup> is a C<sub>1-4</sub> alkylene group; Q<sup>2</sup> is a methylene group;  
a group of the formula:



30 is




35 wherein Z is a group of the formula:



5 [wherein Ar<sup>3</sup> is a phenyl group optionally substituted with a halogen atom, n is an integer of 0 to 3, and Y is a hydrogen atom or a hydroxyl group];

R<sup>1</sup> is a hydrogen atom or methyl;

R<sup>2</sup> is (1) an alkyl group which may be substituted with a C<sub>1-6</sub> alkoxy-carbonyl group, a carboxyl group, a C<sub>1-6</sub> alkyl-carbonyl group, a formyl group or (2) an acyl group represented by the formula:



[wherein R<sup>4</sup> is

- (i) a hydrogen atom,
- 15 (ii) a C<sub>1-6</sub> alkyl group which may have 1 to 5 substituents selected from (a) a hydroxyl group, (b) an amino group which may be substituted with a C<sub>1-6</sub> alkyl-carbonyl group, (c) a mono-C<sub>1-6</sub> alkylamino group, (d) a di-C<sub>1-6</sub> alkylamino group, (e) a carboxyl group, (f) a C<sub>1-6</sub> alkoxy-carbonyl group, (g) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (h) a sulfo group which may be substituted with amino group (i) a 5- to 7-membered cyclic amino group which may have an oxo group or which may be substituted with a hydroxyl group, (j) a C<sub>1-6</sub> alkoxy-carbamoyl group, and (k) a carbamoyloxy group.

(iii) a C<sub>2-6</sub> alkenyl group,

(iv) a C<sub>6-10</sub> aryl group,

(v) a 5- to 11-membered heterocyclic group having at least one hetero atoms of 1 or 2 kinds selected from nitrogen, oxygen and sulfur in addition to a carbon atom, said heterocyclic group being optionally fused with a benzene ring,

30 (vi) a C<sub>1-6</sub> alkyl group which may be substituted with a C<sub>1-6</sub> alkyl-carbonyl group,

35 (vii) a carboxyl group which may be substituted with a


C<sub>1-6</sub> alkyl group,

(viii) a 5- to 7-membered cyclic amino group which may be substituted with

- (a) a C<sub>1-6</sub> alkyl group optionally substituted with (a-1) a hydroxyl group, (a-2) a di-C<sub>1-6</sub> alkylamino group, (a-3) a C<sub>1-6</sub> alkoxy-carbonyl group or (a-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,
- (b) a C<sub>7-16</sub> aralkyl group, (c) a C<sub>1-6</sub> alkyl-carbonyl group whose alkyl portion may be substituted with (c-1) a halogen atom, (c-2) a mono-C<sub>1-6</sub> alkylamino group, (c-3) a C<sub>1-6</sub> alkoxy-carbonyl group or (c-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,
- (d) a C<sub>1-6</sub> alkoxy-carbonyl group, (e) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,
- (f) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with a halogen atom or a C<sub>1-6</sub> alkyl-carbonyl group, (g) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (h) an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group or (i) a C<sub>1-6</sub> alkoxy-carbamoyl group, or
- (ix) a C<sub>6-10</sub> aryloxy group;
- R<sup>5</sup> is a hydrogen atom or a C<sub>1-6</sub> alkyl group].


More preferred compound (I) is one wherein Ar<sup>1</sup> and Ar<sup>2</sup> independently represent, phenyl, 4-chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl;

$Q^1$  is a  $C_{1-4}$  alkylene group;  $Q^2$  is a methylene group;  
a group of the formula:




5

is



10

wherein Z is a group of the formula:



15 [wherein  $\text{Ar}^3$  is a phenyl group optionally substituted with a halogen atom, n is an integer of 0 to 3, and Y is a hydrogen atom or a hydroxyl group];


$R^1$  is a hydrogen atom or methyl;

$R^2$  is an acyl group represented by the formula:

20  $\text{---(C=O)---R}^4$ ,  $\text{---SO}_2\text{---R}^4$ ,  $\text{---(C=O)NR}^5\text{R}^4$  or  $\text{---(C=O)O---R}^4$

[wherein  $R^4$  is represented by the formula:


(1)



25

or

(2)



30 wherein  $R^6$  and  $R^7$  independently represent (a) a hydrogen atom, (b) a  $C_{1-6}$  alkyl group optionally substituted with  
(b-1) a hydroxyl group, (b-2) a di- $C_{1-6}$  alkylamino group, (b-3) a  $C_{1-6}$  alkoxy-carbonyl group, or (b-4) a 5-  
35 to 7-membered heterocyclic group having 1 to 3 hetero

atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (c) a  $C_{7-16}$  aralkyl group, (d) a  $C_{1-6}$  alkyl-carbonyl group whose alkyl portion may be substituted with (d-1) a halogen atom, (d-2) a mono- $C_{1-6}$  alkylamino group, (d-3) a  $C_{1-6}$  alkoxy-carbonyl group, or (d-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (e) a  $C_{1-6}$  alkoxy-carbonyl group, (f) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (g) a mono- $C_{1-6}$  alkyl-carbamoyl group whose alkyl portion may be substituted with (g-1) a halogen atom, or (g-2) a  $C_{1-6}$  alkyl-carbonyl group, (h) an optionally halogenated  $C_{6-10}$  aryl-carbamoyl group, (i) an optionally halogenated  $C_{6-10}$  aryl-carbonyl group, or (j) a  $C_{6-10}$  aryloxy group;  $R^5$  is a hydrogen atom or a  $C_{1-6}$  alkyl group].

Preferred compound (II) is one wherein  $Q^1$  is  $-CH_2-$ ,  $-(CH_2)_2-$ ,  $-(CH_2)_3-$ ,  $-(CH_2)_4-$  or  $-(CH_2)_2CO-$ ;  
 $Q^2$  is  $-CH_2-$ ,  $-(CH_2)_2-$ ,  $-(CH_2)_3-$ ,  $-(CH_2)_4-$ ,  $-CO-$ ,  $-CH_2CO-$  or  $-(CH_2)_2CO-$ ;

$Ar^1$  and  $Ar^2$  independently represent phenyl, 4-chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl or 4-pyridyl;

$Ar^3$  is (1) a phenyl optionally substituted with (a) a  $C_{1-3}$  alkyl group optionally substituted with 1 to 3 halogen atoms, (b) a  $C_{1-3}$  alkoxy group optionally substituted with 1 to 3 halogen atoms or (c) a halogen atom (preferably chlorine, fluorine) (e.g. phenyl, 4-chlorophenyl, 4-fluorophenyl, 4-methoxyphenyl,

3,5-dichlorophenyl, 3,5-difluorophenyl,  
4-trifluoromethylphenyl, etc.), or (2) 2-pyridyl,  
3-pyridyl or 4-pyridyl;

$R^2$  is an acyl group represented by  $-(C=O)-R^4$ ,  
5  $-SO_2-R^4$ ,  $-SO-R^4$ ,  $-(C=O)NR^5R^4$  or  $-(C=O)O-R^4$ ;

$R^5$  is a hydrogen atom or a  $C_{1-3}$  alkyl group such as  
methyl, ethyl, propyl, isopropyl, etc.; and

$R^4$  is (1) a hydrogen atom, (2) an optionally  
substituted lower alkyl group (e.g.  $C_{1-6}$  alkyl such as  
10 methyl, ethyl, propyl, isopropyl, butyl, isobutyl,  
sec-butyl, tert-butyl, pentyl, hexyl, etc.), (3) an  
optionally substituted lower alkyl-carbonyl group (e.g.  
 $C_{1-6}$  alkyl-carbonyl such as acetyl, propionyl, butyryl,  
etc.), (4) a carboxyl group, (5) an optionally  
15 substituted lower alkoxy-carbonyl group (e.g.  $C_{1-6}$   
alkoxy-carbonyl such as methoxycarbonyl,  
ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.),  
(6) an optionally substituted mono-lower  
alkylaminocarbonyl group (e.g. mono- $C_{1-6}$   
20 alkylaminocarbonyl such as methylaminocarbonyl,  
ethylaminocarbonyl, propylaminocarbonyl,  
butylaminocarbonyl, etc.), (7) an optionally  
substituted a di-lower alkylaminocarbonyl group (e.g.  
di- $C_{1-6}$  alkylaminocarbonyl such as  
25 dimethylaminocarbonyl, diethylaminocarbonyl,  
dipropylaminocarbonyl, dibutylaminocarbonyl, etc.), (8)  
a  $C_{6-10}$  aryl group (preferably phenyl) or (9) an  
optionally substituted 5- to 7-membered cyclic amino  
group (preferably 2-piperidyl, 3-piperidyl,  
30 4-piperidyl, 1-pyrrolidinyl, 3-pyrrolidinyl,  
2-piperazinyl, etc.).

The "lower alkyl group," "lower alkyl-carbonyl  
group," "lower alkoxy-carbonyl group," "mono-lower  
alkylaminocarbonyl group," "di-lower  
35 alkylaminocarbonyl" and "5- to 7-membered cyclic amino

group" for  $R^4$  may have 1 to 3 substituents on any carbon atom. The substituent include, for example, a (1) halogen group (e.g. fluorine, chlorine, bromine, iodine, etc.), (2) a lower alkylenedioxy group (e.g.  $C_{1-3}$  alkylenedioxy such as methylenedioxy, ethylenedioxy, etc.), (3) a nitro group, (4) a cyano group, (5) an optionally halogenated lower alkoxy group (e.g. optionally halogenated  $C_{1-6}$  alkoxy such as methoxy, difluoromethoxy, trifluoromethoxy, ethoxy, 2,2,2-trifluoromethoxy, propoxy, isopropoxy, butoxy, 4,4,4-trifluorobutoxy, isobutoxy, sec-butoxy, pentyloxy, hexyloxy, etc.), (6) an optionally halogenated lower alkylthio group (e.g. optionally halogenated  $C_{1-6}$  alkylthio such as methylthio, difluoromethylthio, trifluoromethylthio, ethylthio, propylthio, isopropylthio, butylthio, 4,4,4-trifluorobutylthio, pentylthio, hexylthio, etc.), (7) a hydroxyl group, (8) an amino group, (9) a mono-lower alkylamino group (e.g. mono- $C_{1-6}$  alkylamino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), (10) a di-lower alkylamino group (e.g. di- $C_{1-6}$  alkylamino such as dimethylamino, diethylamino, dipropylamino, dibutylamino, etc.), (11) a 5- to 7-membered cyclic amino group optionally having a hydroxyl group or an oxo group (e.g. morpholino, piperazin-1-yl, piperidino, pyrrolidin-1-yl, 2-pyrrolidone-1-yl, 2-pyridone-1-yl, etc.), (12) an acylamino group ("acylamino group" include, for example, the same groups as mentioned for the substituents of the "optionally substituted aromatic group" for  $Ar^1$ ,  $Ar^2$  and  $Ar^3$  and preferred examples thereof include  $-NHCOOR^3$ ,  $-NHCONHR^3$  and  $-NHCOR^3$  ( $R^3$  is a lower alkyl group (e.g.  $C_{1-6}$  alkyl such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.) or a lower

alkoxy group (e.g. C<sub>1-6</sub> alkoxy such as methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, etc.)), (13) a lower alkyl-carbonyl group (e.g. C<sub>1-6</sub> alkyl-carbonyl such as acetyl, propionyl, etc.), (14) a carboxyl group, (15) a lower alkoxy-carbonyl group (e.g. C<sub>1-6</sub> alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.), (16) a carbamoyl group, (17) a mono-lower alkyl-carbamoyl group (e.g. mono-C<sub>1-6</sub> alkyl-carbamoyl such as methylcarbamoyl, ethylcarbamoyl, etc.), (18) a di-lower alkyl-carbamoyl group (e.g. di-C<sub>1-6</sub> alkyl-carbamoyl such as dimethylcarbamoyl, diethylcarbamoyl, etc.), (19) an aryl-carbamoyl group (e.g. C<sub>6-10</sub> aryl-carbamoyl such as phenylcarbamoyl, naphthylcarbamoyl, etc.), (20) a sulfo group, (21) a lower alkylsulfonyl group (e.g. C<sub>1-6</sub> alkylsulfonyl such as methylsulfonyl, ethylsulfonyl, etc.), (22) an aryl group (C<sub>6-10</sub> aryl such as phenyl, naphthyl, etc.), (23) an aryloxy group (e.g. C<sub>6-10</sub> aryloxy such as phenyloxy, naphthyloxy, etc.), (24) a sulfamoyl group, (25) a mono-lower alkyl-sulfamoyl group (e.g. mono-C<sub>1-6</sub> alkyl-sulfamoyl such as methylsulfamoyl, ethylsulfamoyl, etc.) or (26) a di-lower alkyl-sulfamoyl group (e.g. di-C<sub>1-6</sub> alkyl-sulfamoyl such as dimethylsulfamoyl, diethylsulfamoyl, etc.).

More preferred is a compound wherein Q<sup>1</sup> is -CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>- or -(CH<sub>2</sub>)<sub>3</sub>-;

Q<sup>2</sup> is -CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, -CH<sub>2</sub>CO- or -(CH<sub>2</sub>)<sub>2</sub>CO-;

Ar<sup>1</sup> and Ar<sup>2</sup> independently represent phenyl or 2-pyridyl;

Ar<sup>3</sup> is a phenyl group optionally substituted with 1 to 3 halogen atoms (preferably chlorine, fluorine) (e.g. phenyl, 4-chlorophenyl, 4-fluorophenyl, 3,5-dichlorophenyl, 3,5-difluorophenyl, etc.) or



2-pyridyl;

$R^2$  is (1) a  $C_{1-6}$  alkyl group which may be substituted with a  $C_{1-6}$  alkoxy-carbonyl group, a carboxyl group, a  $C_{1-6}$  alkyl-carbonyl group or a formyl group, or (2) an acyl group represented by  $-(C=O)-R^4$ ,  
5  $-(C=O)NR^5R^4$  or  $-(C=O)O-R^4$ ;

$R^5$  is an hydrogen atom; and

$R^4$  is a hydrogen atom, an optionally substituted lower alkyl group (e.g.  $C_{1-6}$  alkyl such as methyl,  
10 ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.), a carboxyl group, a lower alkenyl group (e.g.  $C_{2-6}$  alkenyl such as vinyl, allyl, isopropenyl, etc.), a lower alkoxy-carbonyl group (e.g.  $C_{1-6}$  alkoxy-carbonyl such as  
15 methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.), a phenyl group or 1-piperazinyl.

The "lower alkyl group" for  $R^4$  may have 1 to 3 substituents on any carbon atom. The substituent include, for example, a hydroxyl group, an amino group,  
20 a di-lower alkylamino group (e.g. di- $C_{1-6}$  alkylamino such as dimethylamino, diethylamino, dipropylamino, dibutylamino, etc.), a 5- to 7-membered cyclic amino group optionally having a hydroxyl group or an oxo group (e.g. morpholino, piperazin-1-yl, piperidino,  
25 pyrrolidin-1-yl, 2-pyrrolidon-1-yl, 2-pyridone-1-yl, etc.), an acylamino group ("acylamino group" include  $-NHCOOR^3$ ,  $-NHCONHR^3$  and  $-NHCOR^3$  ( $R^3$  is a lower alkyl group (e.g.  $C_{1-6}$  alkyl such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl,  
30 pentyl, hexyl, etc.) or a lower alkoxy group (e.g.  $C_{1-6}$  alkoxy such as methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, etc.)), a lower alkyl-carbonyl group ( $C_{1-6}$  alkyl-carbonyl such as acetyl, propionyl, etc.), a carboxyl group, a lower  
35 alkoxy-carbonyl group (e.g.  $C_{1-6}$  alkoxy-carbonyl such as

methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, etc.), a sulfamoyl group, a mono-lower alkyl-sulfamoyl group (e.g. mono-C<sub>1-6</sub> alkyl-sulfamoyl such as methylsulfamoyl, ethylsulfamoyl, etc.) or a  
5 di-lower alkyl-sulfamoyl group (e.g. di-C<sub>1-6</sub> alkyl-sulfamoyl such as dimethylsulfamoyl, diethylsulfamoyl, etc.).

Particularly preferred is a compound wherein Q<sup>1</sup> is  
-(CH<sub>2</sub>)<sub>3</sub>-;

10 Q<sup>2</sup> is -CH<sub>2</sub>- or -(CH<sub>2</sub>)<sub>2</sub>-;

Ar<sup>1</sup> is phenyl or 2-pyridyl;

Ar<sup>2</sup> is phenyl;

Ar<sup>3</sup> is 4-chlorophenyl; n is 0; and Y is an  
hydrogen atom or a hydroxyl group];

15 R<sup>2</sup> is an acyl group represented by -(C=O)-R<sup>4</sup>,  
-(C=O)NR<sup>5</sup>-R<sup>4</sup> or -(C=O)O-R<sup>4</sup>;

R<sup>5</sup> is a hydrogen atom; and


R<sup>4</sup> is (1) a hydrogen atom or (2) a lower alkyl  
group (C<sub>1-6</sub> alkyl such as methyl, ethyl, propyl,  
20 isopropyl, butyl, isobutyl, sec-butyl, tert-butyl,  
pentyl, hexyl, etc.) an optionally having one  
substituent selected from a (a) hydroxyl group, (b) a  
5- to 7-membered cyclic amino group optionally having a  
hydroxyl group or an oxo group (e.g. morpholino,  
25 piperzin-1-yl, piperidino, pyrrolidin-1-yl,  
2-pyrrolidon-1-yl, 2-pyridon-1-yl, etc.) and (c) a  
sulfamoyl group.

In addition, preferred Compound (II) is one  
wherein Ar<sup>1</sup> and Ar<sup>2</sup> independently represent, phenyl, 4-  
30 chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl, or  
4-pyridyl;

Q<sup>1</sup> is a C<sub>1-4</sub> alkylene group; Q<sup>2</sup> is a methylene group;

R<sup>2</sup> is (1) an alkyl group which may be substituted with  
a C<sub>1-6</sub> alkoxy-carbonyl group, a carboxyl group, a C<sub>1-6</sub>  
35 alkyl-carbonyl group, a formyl group or (2) an acyl

group represented by the formula:



[wherein  $R^4$  is


- (i) a hydrogen atom,
- 5 (ii) a  $C_{1-6}$  alkyl group which may have 1 to 5 substituents selected from (a) a hydroxyl group, (b) an amino group which may be substituted with a  $C_{1-6}$  alkyl-carbonyl group, (c) a mono- $C_{1-6}$  alkylamino group, (d) a di- $C_{1-6}$  alkylamino group, (e) a carboxyl group, (f) a  $C_{1-6}$  alkoxy-carbonyl group, (g) a mono- $C_{1-6}$  alkyl-carbamoyl group, (h) a sulfo group which may be substituted with amino group (i) a 5- to 7-membered cyclic amino group which may have an oxo group or which may be substituted with a hydroxyl group, (j) a  $C_{1-6}$  alkoxy-carbamoyl group, and (k) a carbamoyloxy group.
- 15 (iii) a  $C_{2-6}$  alkenyl group,
- (iv) a  $C_{6-10}$  aryl group,
- (v) a 5- to 11-membered heterocyclic groups having at least one hetero atoms of 1 or 2 kinds selected from nitrogen, oxygen and sulfur in addition to a carbon atom, said heterocyclic group being optionally fused with a benzene ring,
- 20 (vi) a  $C_{1-6}$  alkyl group which may be substituted with a  $C_{1-6}$  alkyl-carbonyl group,
- 25 (vii) a carboxyl group which may be substituted with a  $C_{1-6}$  alkyl group,
- (viii) a 5- to 7-membered cyclic amino group which may be substituted with
  - (a) a  $C_{1-6}$  alkyl group optionally substituted with (a-1) a hydroxyl group, (a-2) a di- $C_{1-6}$  alkylamino group, (a-3) a  $C_{1-6}$  alkoxy-carbonyl group or (a-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being
  - 30 optionally fused with a benzene ring,
  - 35

- (b) a C<sub>7-16</sub> aralkyl group, (c) a C<sub>1-6</sub> alkyl-carbonyl group whose alkyl portion may be substituted with (c-1) a halogen atom, (c-2) a mono-C<sub>1-6</sub> alkylamino group, (c-3) a C<sub>1-6</sub> alkoxy-carbonyl group or (c-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,
- (d) a C<sub>1-6</sub> alkoxy-carbonyl group, (e) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,
- (f) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with a halogen atom or a C<sub>1-6</sub> alkyl-carbonyl group, (g) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (h) an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group or (i) a C<sub>1-6</sub> alkoxy-carbamoyl group, or
- (ix) a C<sub>6-10</sub> aryloxy group;
- R<sup>5</sup> is a hydrogen atom or a C<sub>1-6</sub> alkyl group]; and
- Ar<sup>3</sup> is a phenyl group optionally substituted with a halogen atom.

In addition, preferred Compound (II) is one wherein Ar<sup>1</sup> and Ar<sup>2</sup> independently represent, phenyl, 4-chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl;


Q<sup>1</sup> is a C<sub>1-4</sub> alkylene group; Q<sup>2</sup> is a methylene group;

R<sup>2</sup> is (1) an alkyl group which may be substituted with a C<sub>1-6</sub> alkoxy-carbonyl group, a carboxyl group, a C<sub>1-6</sub> alkyl-carbonyl group, a formyl group or (2) an acyl group represented by the formula:




[wherein R<sup>4</sup> is a group represented by the formula:

(1)



or

5 (2)



wherein  $R^6$  and  $R^7$  independently represent (a) a  
 10 hydrogen atom, (b) a  $C_{1-6}$  alkyl group optionally  
 substituted with  
 (b-1) a hydroxyl group, (b-2) a di- $C_{1-6}$  alkylamino  
 group, (b-3) a  $C_{1-6}$  alkoxy-carbonyl group, or (b-4) a 5-  
 to 7-membered heterocyclic group having 1 to 3 hetero  
 15 atoms selected from nitrogen, oxygen and sulfur in  
 addition to carbon atoms, said heterocyclic group being  
 optionally fused with a benzene ring, (c) a  $C_{7-16}$   
 aralkyl group, (d) a  $C_{1-6}$  alkyl-carbonyl group whose  
 alkyl portion may be substituted with (d-1) a halogen  
 20 atom, (d-2) a mono- $C_{1-6}$  alkylamino group, (d-3) a  $C_{1-6}$   
 alkoxy-carbonyl group, or (d-4) a 5- to 7-membered  
 heterocyclic group having 1 to 3 hetero atoms selected  
 from nitrogen, oxygen and sulfur in addition to carbon  
 atoms, said heterocyclic group being optionally fused  
 25 with a benzene ring, (e) a  $C_{1-6}$  alkoxy-carbonyl group,  
 (f) a formyl group which may be substituted with a 5-  
 to 7-membered heterocyclic group having 1 to 3 hetero  
 atoms selected from nitrogen, oxygen and sulfur in  
 addition to carbon atoms, said heterocyclic group being  
 30 optionally fused with a benzene ring, (g) a mono- $C_{1-6}$   
 alkyl-carbamoyl group whose alkyl portion may be  
 substituted with (g-1) a halogen atom, or (g-2) a  $C_{1-6}$   
 alkyl-carbonyl group, (h) an optionally halogenated  $C_{6-10}$   
 10 aryl-carbamoyl group, (i) an optionally halogenated  
 35  $C_{6-10}$  aryl-carbonyl group, or (j) a  $C_{6-10}$  aryloxy group;

and

R<sup>5</sup> is a hydrogen atom or a C<sub>1-6</sub> alkyl group]; and  
Ar<sup>3</sup> is a phenyl group optionally substituted with a  
halogen atom.

5           Examples of the preferred compound include the  
following.

5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-1-formylamin  
o-2,2-diphenylpentane hydrochloride,  
5-[4-(4-Fluorophenyl)-piperadin-1-yl]-1-formylamino-  
10 2,2-diphenylpentane dihydrochloride,  
7-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-1-  
formylamino-2,2-diphenylheptane hydrochloride,  
1-[5-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-  
diphenylpentyl]-3-(3-hydroxypropyl)urea hydrochloride,  
15 1-[5-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-  
diphenylpentyl]-3-(4-hydroxybutyl)urea hydrochloride,  
1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-  
diphenylpentyl]-3-(3-diethylaminopropyl)urea,  
2-[3-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-  
20 diphenylpentyl]ureido]ethanesulfonamide hydrochloride,  
N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphe  
nyl]pentylmalonamic acid,  
N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphe  
nyl]pentylglutamic acid,  
25 N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2-phenyl-  
(2-pyridyl)]pentylsuccinamic acid,  
1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-3-(piperidin-4-yl)urea,  
Ethyl 4-[4-[5-[4-(4-chlorophenyl)-4-hydroxy-  
30 piperidino]-2,2-diphenylpentyl]aminocarbonylamino]  
piperidino-4-oxobutyrate,  
N-Ethyl-4-[5-[4-(4-chlorophenyl)-4-hydroxy-  
piperidino]-2,2-diphenylpentyl]aminocarbonylamino-1-  
piperidinecarboxamide,  
35 N-Ethoxycarbonylmethyl-4-[5-[4-(4-chlorophenyl)-  
4-hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonyl-

amino-1-piperidinecarboxamide,  
Ethyl 3-[4-[5-[4-(4-chlorophenyl)-4-hydroxy-  
piperidino]-2,2-diphenylpentyl]aminocarbonylamino  
piperidino]-3-oxopropionate,  
5 1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-3-(1-ethylpiperidin-4-yl)urea,  
1-[(Piperidin-4-yl)carboxamido]-5-[4-(4-chlorophenyl)-4-  
-hydroxypiperidino]-2,2-diphenylpentane,  
1-[[N-Ethylcarbamoyl]piperidin-4-yl]carboxamido]-5-[4-  
10 (4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpenta  
ne,  
1-[[N-(Ethoxycarbonylacetyl)piperidin-4-yl]carboamido]-  
5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl  
pentane,  
15 1-[[N-(3-Methoxycarbonylpropionyl)piperidin-4-yl]carbox  
amido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-  
diphenylpentane, or a salt thereof.

Among them, especially,

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
20 2,2-diphenylpentyl]-3-(piperidin-4-yl)urea,  
Ethyl 4-[4-[5-[4-(4-chlorophenyl)-4-hydroxy-  
piperidino]-2,2-diphenylpentyl]aminocarbonylamino]  
piperidino-4-oxobutyrate,  
N-Ethyl-4-[5-[4-(4-chlorophenyl)-4-hydroxy-  
25 piperidino]-2,2-diphenylpentyl]aminocarbonylamino-1-  
piperidinecarboxamide,  
N-Ethoxycarbonylmethyl-4-[5-[4-(4-chlorophenyl)-  
4-hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonyl-  
amino-1-piperidinecarboxamide,  
30 Ethyl 3-[4-[5-[4-(4-chlorophenyl)-4-hydroxy-  
piperidino]-2,2-diphenylpentylaminocarbonylamino]  
piperidino]-3-oxopropionate,  
1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-3-(1-ethylpiperidin-4-yl)urea,  
35 1-[(Piperidin-4-yl)carboamido]-5-[4-(4-chlorophenyl)-4-  
hydroxypiperidino]-2,2-diphenylpentane,

1-[[N-Ethylcarbamoyl]piperidin-4-yl]carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane,

1-[[N-(Ethoxycarbonylacetyl)piperidin-4-yl]carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane,


1-[[N-(3-Methoxycarbonylpropionyl)piperidin-4-yl]carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane or a salt thereof, is preferred.

The above mentioned compounds may be in any form of free compound, salt, hydrate, etc.

While a plurality of synthetic technologies are contemplatable for producing the compound (I) and a salt thereof (hereinafter abbreviated to a "compound (I)," merely) and compound (II) and a salt thereof (hereinafter abbreviated to a "compound (II)," merely), typical production technology will be illustrated as follows:

In the explanation of the following processes, starting materials and reaction products may form a salt thereof which does not inhibit the reaction.

#### Process 1



(In the above schema, L is a leaving group and the other symbols have the same meanings as defined above).

The reaction is carried out applying a usual



alkylation of an amino group [e.g. procedure described in Organic Functional Group Preparations, Vol. 2, Academic Press Inc.]. Examples of the leaving group include a halogen atom (preferably chloro, bromo, iodo, etc.), a methanesulfonyloxy group, a p-toluenesulfonyloxy group, a benzenesulfonyloxy group, etc.

The reaction is carried out by stirring in an inert solvent within the range from room temperature to 100°C (preferably room temperature to 50°C) for 0.5 to 1 day. Usually, 1 to 3 equivalents of a base is added to the reaction system but is not essential. As the inert solvent, alcoholic solvent, etheral solvent, halogenated solvent, aromatic solvent, acetonitrile, N,N-dimethylformylamido (DMF), acetone, methyl ethyl ketone and dimethyl sulfoxide can be used alone or in combination thereof. Among them, acetonitrile, DMF, acetone and ethanol are preferred.


The base include strong bases (1) such as hydrides of alkaline or alkaline earth metals (e.g. lithium hydride, sodium hydride, potassium hydride, calcium hydride, etc.), amides of alkaline or alkaline earth metals (e.g. lithium amide, sodium amide, lithium diisopropylamide, lithium dicyclohexylamide, lithium hexamethylsilazide, sodium hexamethylsilazide, potassium hexamethylsilazide, etc.) and lower (C<sub>1-4</sub>) alkoxides of alkaline or alkaline earth metals (e.g. sodium methoxide, sodium ethoxide, potassium t-butoxide, etc.); inorganic salts (2) such as hydroxides of alkaline or alkaline earth metals (e.g. sodium hydroxide, potassium hydroxide, lithium hydroxide, barium hydroxide, etc.), carbonates of alkaline or alkaline earth metals (e.g. sodium carbonate, potassium carbonate, cesium carbonate, etc.) and hydrogencarbonates of alkaline or alkaline earth metals (e.g. sodium hydrogencarbonate, potassium

hydrogencarbonate, etc.); and organic bases (3) such as amines (e.g. triethylamine, diisopropylethylamine, N-methylmorpholine, 4-dimethylaminopyridine, DBU (1,8-diazabicyclo[5.4.0]-7-undecene), DBN (1,5-diazabicyclo[4.3.0]non-5-ene), etc.) and basic heterocyclic compounds (e.g. pyridine, imidazole, 2,6-lutidine, etc.).

The compound (I) or (II) obtained by the above process can be further converted to the objective product of this invention by a usual organic synthesis reaction such as hydrolysis, halogenation, oxidation, reduction, alkylation, acylation, ring formation etc. The reaction examples include the following process.

When the compound has carbonyl in the molecule, it can be converted to the following compound having a hydroxyl group by the Grignard reaction.

## Process 2



(wherein M is a metal (e.g. lithium, sodium, bromomagnesium, etc.) used for so-called Grignard reaction; and the other symbols have the same meanings as defined above).

The Grignard reaction is conducted by reacting 1

to 10 equivalents of a so-called Grignard reagent prepared separately or alkyl lithium or alkyl sodium with the compound (VII) or (VII') in an ethereal solvent at room temperature to 80°C (preferably 30 to 60°C) for 1 to 24 hours. The reaction is preferably conducted under the condition of deoxidation in the absence of water. It is preferred to conduct the reaction in the presence of anhydrous cerium chloride (catalytic amount to 2 equivalent, preferably 1 equivalent).

When  $R^1$  and  $R^2$  independently represent an acyl group or an alkyl-carbonyl group, the group can be converted into an alkyl group by the reduction.

The reduction can be conducted by the procedure using metal hydrides or catalytic reduction process. The catalytic reduction process can be conducted by reacting with a catalytic amount of a metal catalyst such as Raney-nickel, platinum oxide, palladium metal, palladium-on-carbon, etc. in an inert solvent (e.g. alcoholic solvent) at room temperature to 100°C under a hydrogen pressure of 1 to 100 atm for 1 to 48 hours.

The reduction using the metal hydride can be easily conducted in an inert solvent using a metal hydride (e.g. lithium aluminum hydride, sodium borohydride, lithium borohydride, sodium cyanoborohydride, diborane, dibutylaluminum hydride, etc.) or a metal (e.g. zinc, iron, sodium, potassium, etc.). The inert solvent include ethereal solvents (e.g. diethyl ether, tetrahydrofuran, dioxane, etc.), alcoholic solvents (e.g. methanol, ethanol, tert-butanol, etc.), toluene and hexane. The preferred metal hydride include lithium aluminum hydride. The amount of the metal hydride to be used is from 4 to 20 equivalents, more preferably from 6 to 12 equivalents. The reaction is conducted at the reaction temperature of -70 to 100°C for 30 minutes to 18 hours. The preferred reaction temperature varies depending on the

kind of a reducing agent to be used, but is usually from 30 to 70°C. It is also possible to selectively reduce only a cyano or ester group.


5 The conversion from a ketone represented by the compound (VII) or (VII') to an alcohol of -CH(OH) can be easily accomplished by reacting with the metal hydride (e.g. lithium aluminum hydride, sodium borohydride, lithium borohydride, sodium  
10 cyanoborohydride, diborane, dibutylaluminum hydride, etc.) in an inert solvent. The inert solvent include etheral solvents (e.g. ethyl ether, tetrahydrofura, dioxane, etc.) and alcoholic solvents (e.g. methanol, ethanol, tert-butanol, etc), toluene and hexane. The amount of the metal hydride to be used is from 1 to 20  
15 equivalents, more preferably from 3 to 12 equivalents. The reaction temperature is from -70 to 100°C. The preferred reaction temperature and reaction time vary depending on the kind of a reducing agent to be used. In case of the metal hydride, the reduction is  
20 preferably conducted at 0 to 30°C for 30 minutes to 18 hours.

When R<sup>1</sup> or R<sup>2</sup> independently represents an acyl group, the group can also be converted into another acyl group, directly or through hydrolysis. The  
25 hydrolysis includes an alkali hydrolysis and an acid hydrolysis. In case of the "alkali hydrolysis," a compound is reacted with an alkali (e.g. inorganic hydroxides such as lithium hydroxide, sodium hydroxide, potassium hydroxide, barium hydroxide, etc.) in a  
30 solvent (e.g. water, alcohols, ethers alone or a mixed solvent using two or more kinds of them). As the solvent, a mixed solvent of water and methanol is preferred. As the alkali, sodium hydroxide is preferred. The usage amount of the alkali is about 2  
35 to 100 equivalents, preferably about 5 to 100 equivalents, relative to the compound. The reaction

temperature is from about 10 to 120°C, preferably from about 50 to 120°C. The reaction time is from about 5 minutes to 100 hours, preferably from about 10 to 50 hours. In the preferred reaction example, the solvent  
5 is a mixed solvent of water and methanol and the reaction temperature is from about 50 to 120°C and, the reaction time is from about 10 to 50 hours.

Regarding the "acid hydrolysis process," a compound may be heated with stirring in water, acetic  
10 acid or an alcoholic solvent in the presence of an excess amount of mineral acid (e.g. hydrochloric acid, sulfuric acid, phosphoric acid, etc.) at room temperature to 120°C for 0.5 to 18 hours. Preferably, the heating is conducted in the presence of dilute  
15 hydrochloric acid alone or in combination with acetic acid at room temperature to 60°C.

When  $R^1$  and  $R^2$  independently represent a "protective group of an amino group," there can be sometimes used reduction process, ultraviolet  
20 irradiation process, hydrazine process, etc. in addition to the hydrolysis process. Typical examples of the "reduction process" include a catalytic reduction process. For example, starting materials are stirred in an inert solvent (e.g. water, alcoholic  
25 solvent, ethyl acetate, etheral solvent, etc.) in the presence of metal catalysts (catalytic amount to one equivalent) such as palladium catalyst (e.g. palladium acetate, palladium-carbon, palladium black, palladium-barium carbonate, etc.), platinum oxide and  
30 Ranney-nickel, etc. at room temperature to 100°C under a hydrogen pressure of 1 to 100 atm (preferably from 1 to 10 atm) for 0.5 to 24 hours. If necessary, a catalytic amount to 2 equivalents of a mineral acid such as hydrochloric acid or an organic acid such as  
35 acetic acid is sometimes added.



[In the above schema,  $\text{R}^2$  is an acyl group; and the other symbols have the same meanings as defined above].

The acylation can be conducted according to the per se known procedure described in Organic Functional Group Preparations, Vol. 2, Academic Press Inc.

For example, when an acyl group represented by  $\text{R}^2$  is  $-(\text{C}=\text{O})-\text{R}^4$ ,  $-\text{SO}_2-\text{R}^4$ ,  $-\text{SO}-\text{R}^4$  or  $-(\text{C}=\text{O})\text{O}-\text{R}^4$  ( $\text{R}^4$  is an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group), the acylation reaction is conducted by reacting 1 to 5 equivalents, preferably 1 to 3 equivalents, of a reactive derivative of the corresponding organic acid with the compound (IX) or (IX') in an inert solvent at the reaction temperature of  $-20$  to  $50^\circ\text{C}$  (preferably  $0^\circ\text{C}$  to room temperature) for 5 minutes to 100 hours. As the inert solvent, there can be used etheral solvent, halogenated solvent, aromatic solvent, acetonitrile, N,N-dimethylformulamido (DMF), acetone, methyl ethyl ketone, dimethylsulfoxide (DMSO), water, etc. alone or in combination thereof. Among them, acetonitrile, dichloromethane and chloroform are preferred. The reaction sometimes proceed more smoothly in the presence of 1 to 10 equivalents, preferably 1 to 3 equivalents of a base. As the base, both inorganic and organic bases are effective. The inorganic base



includes hydroxides, hydrides, carbonates, hydrogencarbonates, organic acid salts of alkaline or alkaline earth metals. Among them, potassium carbonate, sodium carbonate, sodium hydroxide, potassium hydroxide, sodium hydrogencarbonate and potassium hydrogencarbonate are preferred. As the organic base, tertiary amines such as triethylamine is preferred. The reactive derivative includes acid anhydride, acid halide (e.g. acid chloride, acid bromide, etc.) and active ester. Among them, acid halide is preferred.

Acylation using carboxylic acid can be used a procedure of reacting 1 to 1.5 equivalents of carboxylic acid in an inert solvent (e.g. halogenated solvent, acetonitrile, etc.) with a dehydration condensing agent such as dicyclohexylcarbodiimide (DCC) (1 to 1.5 equivalents) at room temperature for 0.5 to 24 hours.

When an acyl group represented by  $R^2$  is  $-(C=O)NH-R^4$ ,  $-(C=S)NH-R^4$ ,  $-(C=S)O-R^4$  or  $-(C=O)O-R^4$  ( $R^4$  has the same meanings as defined above), the acylation is conducted in an inert solvent (e.g. halogenated solvent, acetonitrile, etc.) at the reaction temperature of  $-20$  to  $50^\circ\text{C}$  (preferably  $0^\circ\text{C}$  to room temperature) for 5 minutes to 100 hours, using one equivalent or excess amount of the corresponding isocyanate ( $OCN-R^4$  ( $R^4$  has the same meanings as defined above) and isothiocyanate ( $SCN-R^4$  ( $R^4$  has the same meanings as defined above)). In order to accelerate the reaction, the reaction is sometimes conducted in the presence of 1 to 10 equivalents of an organic base such as triethylamine. When the acyl group represented by  $R^2$  is  $-CONR^5-R^4$  ( $R^4$  and  $R^5$  have the same meanings as defined above) (hydrogen is preferred as  $R^5$ ), it is also possible to produce by the following exchange reaction (process 4).




## Process 4



15 [wherein Ph is a phenyl group; and the other symbols have the same meanings as defined above].

The reaction proceeds by reacting one equivalent to excess amount of amine (HN-R<sup>4</sup>-R<sup>5</sup> (R<sup>4</sup> and R<sup>5</sup> have the same meanings as defined above)) with the compound (X) or (X') in an inert solvent such as acetonitrile, DMF, water, etc. in the presence of 1 to 10 equivalents of an inorganic base (e.g. potassium carbonate, sodium carbonate, etc.) at room temperature to 50°C for 1 to 24 hours.

## Process 5



[wherein, the symbols have the same meanings as defined above; and L' is a leaving group].

15 In the process 5, the objective product can be obtained by reacting 1 equivalent to excess amount of:



20 with the compound (XII) or (XII'). The reaction conditions are the same as those of the alkylation reaction of the amino group in the "process 1." As the base, the above strong base, inorganic base or organic base is used.

25 The leaving group L' used in the "process 5" includes the same one for L. Among them, bromo and iodo are preferred. When "R<sup>1</sup> and R<sup>2</sup> bonds together with the adjacent nitrogen to form an optionally substituted nitrogen-containing heterocyclic group,"

30 the objective product can be synthesized by introducing the corresponding nitrogen-containing heterocycle according to the "process 5." For example, morpholino, piperazino, 1-piperaziny, 1-imidazolyl, phthalimide, etc. can be easily introduced.

35 The compound used as the starting material in the above "process 1" and "process 2" can be synthesized by

using the synthesis procedures which are known in references in combination. For example, the following compound used in the above "process 1" is easily available or synthesized.

5




Among them, the compound wherein Z is -C(OH)-(CH<sub>2</sub>)<sub>n</sub>-Ar<sup>3</sup> can be produced from the corresponding ketone according to the same manner as that described in "process 2."


The compound (III) or (III') as the starting material can be synthesized by the per se known procedure, and examples thereof include the following schema 1.

15

Schema 1



20



25

[wherein J<sup>1</sup> is a cyano group, a carboxyl group, a lower (C<sub>1-3</sub>) alkoxy-carbonyl group or a formyl group; J<sup>2</sup> is a group capable of converting into a leaving group (e.g. cyano, carboxyl, lower (C<sub>1-3</sub>) alkoxy-carbonyl, protected hydroxyl group, etc.); L'' is the same meanings as defined in L; and the other symbols have the same meanings as defined above.]

30

It is possible to convert to the compound (XVII) by reacting the compound (XV) with one equivalent to excess amount of the compound (XVI) in any inert solvent (e.g. etheral solvent, DMF, DMSO, alcoholic

35

solvent, acetonitrile, acetone, etc.) or mixed solvent thereof in the presence of a base (usually 1 to 3 equivalents) at -20 to 120°C for 5 minutes to 24 hours. The compound (XVII) can also be obtained by heating the  
5 compound (XV) and excess acrylonitrile or lower alkyl acrylate (2 to 10 equivalents) in the presence of a base catalyst.




The base include strong bases (1) such as hydrides of alkaline or alkaline earth metals (e.g. lithium  
10 hydride, sodium hydride, potassium hydride, calcium hydride, etc.), amides of alkaline or alkaline earth metals (e.g. lithium amide, sodium amide, lithium diisopropylamide, lithium dicyclohexylamide, lithium hexamethylsilazide, sodium hexamethylsilazide,  
15 potassium hexamethylsilazide, etc.) and lower (C<sub>1-4</sub>) alkoxides of alkaline or alkaline earth metals (e.g. sodium methoxide, sodium ethoxide, potassium t-butoxide, etc.); inorganic salts (2) such as hydroxides of alkaline or alkaline earth metals (e.g.  
20 sodium hydroxide, potassium hydroxide, lithium hydroxide, barium hydroxide, etc.), carbonates of alkaline or alkaline earth metals (e.g. sodium carbonate, potassium carbonate, cesium carbonate, etc.) and hydrogencarbonates of alkaline or alkaline earth  
25 metals (e.g. sodium hydrogencarbonate, potassium hydrogencarbonate, etc.); and organic bases (3) such as amines (e.g. triethylamine, diisopropylethylamine, N-methylmorpholine, 4-dimethylaminopyridine, DBU (1,8-diazabicyclo[5.4.0]-7-undecene), DBN  
30 (1,5-diazabicyclo[4.3.0]non-5-ene), etc.) and basic heterocyclic compounds (e.g. pyridine, imidazol, 2,6-lutidine, etc.).

The compound (XVII) can be converted to the compound (III) or (III') by appropriately combining per  
35 se known processes, for example, general organic synthesis reactions such as hydrolysis, halogenation,




oxidation, reduction, alkylation, acylation, ring formation etc.

The reaction example include the following process.

5 Method 1



Method 2



[wherein T is a bond, an oxygen atom or an

optionally oxidized sulfur atom; L and L" independently represent a leaving group; a and m independently represent an integer of 0 to 5 and the total of them is 1 to 6; h is an integer of 0 to 2; and Q' is a group  
5 obtained by removing one methylene group from Q<sup>2</sup>.]

The reduction reaction of the compound (XX) and the compound (XXIV) can be conducted by the process using metal hydrides or catalytic reduction process. The catalytic reduction process can be conducted by  
10 reacting with a catalytic amount of a metal catalyst such as Ranney-nickel, platinum oxide, metallic palladium, palladium-carbon, etc. in an inert solvent (e.g. alcoholic solvent) at room temperature to 100°C under a hydrogen pressure of 1 to 100 atm for 1 to 48  
15 hours.

The reduction reaction using the metal hydride can be easily conducted by reacting in an inert solvent using a metal hydride (e.g. lithium aluminum hydride, sodium borohydride, lithium borohydride, sodium  
20 cyanoborohydride, diborane, dibutylaluminum hydride, etc.) or a metal (e.g. zinc, iron, sodium, potassium, etc.). The inert solvent include ethereal solvents (e.g. diethyl ether, tetrahydrofuran, dioxane, etc.), alcoholic solvents (e.g. methanol, ethanol,  
25 tert-butanol, etc.), toluene and hexane. The preferred metal hydride include lithium aluminum hydride. The amount of the metal hydride to be used is from 4 to 20 equivalents, more preferably from 6 to 12 equivalents. The reaction temperature is from -70 to 100°C. The  
30 preferred reaction temperature varies depending on the kind of a reducing agent to be used, but is normally from 30 to 70°C. The reaction time is from 30 minutes to 18 hours. It is also possible to selectively reduce only a cyano or ester group.


35 The conversion from a hydroxyl group to a leaving group or introduction of a protective group of an amino

group can be conducted according to the procedure described in Comprehensive Organic Transformations, VCH Publishers Inc.

5 The compound (XXII) can be converted to the compound (XXIII) by the Wittig reaction. The reaction can be conducted in an inert solvent (e.g. alcoholic solvent, etheral solvent, etc.), if necessary, in the presence of a base at 20 to 60°C for 5 minutes to 18 hours, using 1 equivalent to excess amount of a Wittig reagent (e.g. ethyl triphenylphosphoranilidene-acetate, 10 ethyl diethylphosphonoacetate, etc.). The base include strong bases (1) such as sodium hydride, t-butoxy potassium, etc.); inorganic bases (2) such as hydroxides of alkaline or alkaline earth metals (e.g. 15 sodium hydride, potassium hydroxide, lithium hydroxide, barium hydroxide, etc.), carbonates of alkaline or alkaline earth metals (e.g. sodium carbonate, potassium carbonate, cesium carbonate, etc.) and hydrogencarbonates of alkaline or alkaline earth metals 20 (e.g. sodium hydrogencarbonate, potassium hydrogencarbonate, etc.); and organic bases (3) amines (e.g. triethylamine, DBU, etc.).

In case of reducing the double bond, the catalytic reduction process mentioned above can be used.

25



30 as one of starting materials can be synthesized from aryl acetonitrile or diaryl ketone according to the per se known procedure (e.g. Synthesis, page 172, 1977).

Moreover, in any of the aforementioned reactions and any of the reactions for synthesizing the starting 35 compounds, when the raw materials have amino, carboxyl or hydroxyl group as a substituent, these functional

groups may be protected with protective groups which are commonly used in peptide chemistry or related art. The desired compounds can be then be obtained by eliminating such protective groups when needed.

5           The amino-protective group that can be used includes, for example, C<sub>1-6</sub> alkyl-carbonyl (e.g. formyl, acetyl, ethylcarbonyl, etc.), C<sub>1-6</sub> alkyloxycarbonyl (e.g. methoxycarbonyl, ethoxycarbonyl, etc.), benzoyl group, C<sub>7-10</sub> aralkyl-carbonyl (e.g. benzylcarbonyl, 10           etc.), trityl, phthaloyl and N,N-dimethylaminomethylene. These groups may respectively have 1 to 3 substituents, for example, halogen (e.g. fluorine, chlorine, bromine, iodine, etc.) and nitro.

15           The carboxyl-protective group which can be used includes, for example, C<sub>1-6</sub> alkyl (e.g. methyl, ethyl, ethyl, n-propyl, isopropyl, butyl, tert-butyl, etc.), phenyl, trityl and silyl. These groups may respectively have 1 to 3 substitutes, for example, 20           halogen (e.g. fluorine, chlorine, bromine, iodine, etc.), C<sub>1-6</sub> alkyl-carbonyl (e.g. formyl, acetyl, propionyl, butylcarbonyl, etc.) and nitro.

          The hydroxyl-protective group which can be used includes, for example, C<sub>1-6</sub> alkyl (e.g. methyl, ethyl, 25           n-propyl, isopropyl, butyl, tert-butyl, etc.), phenyl, C<sub>7-10</sub> aralkyl group (e.g. benzyl, etc.), formyl, C<sub>1-6</sub> alkyl-carbonyl group (e.g. acetyl, propionyl, etc.), benzoyl, C<sub>7-10</sub> aralkyl-carbonyl (e.g. benzylcarbonyl, etc.), tetrahydropyranyl, tetrahydrofuranyl, and silyl. 30           These groups may respectively have 1 to 3 substituents, for example, halogen (e.g. fluorine, chlorine, bromine, iodine, etc.), C<sub>1-6</sub> alkyl (methyl, ethyl, n-propyl, etc.), phenyl, C<sub>7-10</sub> aralkyl (e.g. benzyl, etc.) and nitro.

35           These protective groups can be removed by the per



se known procedures or any procedures analogous thereto. For example, a process using an acid, a base, a reducing agent, an ultraviolet light, hydrazine, phenylhydrazine, N-methyldithiocarbamate, 5 tetrabutylammonium fluoride or palladium acetate can be utilized.

The salt of the compound (I) or (II) include, for example, salts with inorganic bases, salts with organic bases, salts with inorganic acids and salts with basic 10 or acidic amino acids. The preferred salts with inorganic bases include, for example, alkaline metal salt (e.g. sodium salt, potassium salt, etc.), alkaline earth metal salt (e.g. calcium salt, magnesium salt, etc.) and aluminum salt and ammonium salt. The 15 preferred salts with organic bases include, for example, salts with trimethylamine, triethylamine, pyridine, picoline, ethanolamine, diethanolamine, triethanolamine, dicyclohexylamine, N,N'-dibenzylethylenediamine, etc. The preferred salts 20 with inorganic acids include, for example, salts with hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid, etc. The preferred salts with organic acids include, for example, salts with formic acid, acetic acid, trifluoroacetic acid, 25 fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, etc. The preferred salts with basic amino acids include, for example, salts with arginine, lysine, 30 ornithine, etc. The preferred salts with acidic amino acids include, for example, salts with aspartic acid, glutamic acid, etc.

Among them, pharmaceutically acceptable salts are particularly preferred. In case the compound has a 35 basic functional group in its molecule, the pharmaceutically acceptable salts include, for example,

inorganic salt such as hydrochloride, sulfate, phosphate, hydrobromide, etc., or organic salt such as acetate, maleate, fumarate, succinate, methanesulfonate, p-toluenesulfonate, citrate, tartrate, etc. In case of having an acidic functional group, the pharmaceutically acceptable salts include, for example, inorganic salt such as alkaline metal salt (e.g. sodium salt, potassium salt, etc.) or alkaline metal salt (e.g. calcium salt, magnesium salt, etc.) and ammonium salt.

The compounds (I) and (II) of this invention and their salts can be separated and purified by known procedures such as solvent extraction, pH change, redistribution, crystallization, recrystallization, chromatography, etc. The starting compounds of the compounds (I) and (II) of this invention and their salts can be separated and purified by the same known procedures as those described above, but the reaction mixture containing them may be respectively be submitted to the next reaction steps.

When the compounds (I) and (II) of this invention and their salts include optical isomers, stereoisomers, position isomers or rotational isomers, these are also included as the compounds of this invention and can be obtained by the per se known synthesis and isolation procedures. For example, when optical isomers exist in the compounds of this invention, optical isomers resolved from the compounds can also be included in this invention.

The optical isomers can be produced by the per se known method. Specifically, a desired optically active isomer can be obtained by using an optically active intermediate, or by optically resolving a mixture of racemic modifications as a final product according to a usual procedure.

As an optical resolution procedure, for example,

there can be used the following fractional recrystallization process, chiral column process, diastereomer process, etc.,.

(1) Fractional recrystallization process

5 A process comprising reacting a racemic modification with an optically active compound to form a salt and separating the salt according to a fractional recrystallization method and optionally producing a free optical isomer through a neutralizing  
10 step.

(2) Chiral column process

A process of separating a racemic modification or a salt thereof using a column for separating an optical isomer (chiral column). In case of a liquid  
15 chromatography, for example, the optical isomer is separated by adding a mixture of optical isomers to a chiral column such as ENANTIO-OVM (manufactured by Toso Co.) and developing with water, various buffers (e.g. phosphate buffer, etc.) and an organic solvent (e.g.  
20 ethanol, methanol, acetonitrile, etc.) alone or in combination thereof. In case of a gas chromatography, it is separated by using a chiral column such as CP-Chirasil-DeX CB (manufactured by GL Science Co.).

(3) Diastereomer process

25 A process comprising reacting a mixture of racemic modifications with an optically active reagent to form a mixture of diastereomers, separating the mixture into a single substance through normal means (e.g. fractional recrystallization, chromatography, etc.) and  
30 cleaving the optically active reagent site due to a chemical treatment such as hydrolysis reaction. For example, when the compound of this invention has a hydroxyl group or a primary or secondary amino group in the molecule, a diastereomer as an ester or amide can  
35 be obtained by subjecting the compound and an optically active organic acid (e.g.

MPTA[ $\alpha$ -methoxy- $\alpha$ -(trifluoromethyl)phenylacetic acid, (-)-menthoxyacetic acid, etc.) to a condensation reaction. On the other hand, when the compound of this invention has a carboxylic group, the diastereomer as the ester or amide can be obtained by subjecting the compound and an optically active amine or an alcohol reagent to a condensation reaction. The separated diastereomer is converted into an optical isomer of the original compound by subjecting to an acid hydrolysis or basic hydrolysis reaction.

The compounds (I) and (II) of this invention and their salts can be safely administered as they are or as a pharmaceutical composition containing a medicinally acceptable carrier in various dosage forms such as tablest (inclusive of dragees and film-coated tablets), powders, granules, capsules (inclusive of soft capsules), solutions, injections, suppositories, controlled-release preparations, etc. by the oral route or parenteral route (e.g. local, rectal or intravenous administration) according to the per se known method. An amount of the compound (I) or a salt thereof contained in the preparation of this invention is from 0.1 to 100% by weight based on the total weight. The dosage is dependent on the subject, route of administration, administration route, diseases, etc., but for the treatment of viral encephalitis, etc., for instance, the recommend oral regimen for an adult patient (b.wt. 60 kg) is about 0.1 to 500 mg/day, preferably about 1 to 100 mg/day, more preferably about 5 to 100 mg/day, to be administered once a day or in a few divided doses daily.

The pharmaceutically acceptable carrier includes a variety of organic and inorganic carriers or vehicles which are commonly used in the pharmaceutical field. Here, excipients, lubricants, binders, disintegrators, etc. are all subsumed in the concept of carrier for

solid preparations, while solvents, solubilizers, suspending agents, isotonizing agents, buffers, analgesics, etc. can be used in the formulation of liquid preparations. Where necessary, various additives such as preservatives, antioxidants, coloring agents, sweeteners, absorbents, moistening agents, etc. can also be added. The preferred excipient includes lactose, sucrose, D-mannitol, starch, corn starch, crystalline cellulose, and light silicic anhydride. The lubricant include magnesium stearate, calcium stearate, talc, and colloidal silica.

The binder includes crystalline cellulose, saccharose, D-mannitol, dextrin, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyvinylpyrrolidone, starch, sucrose, gelatin, methylcellulose, and carboxymethylcellulose.

The disintegrator includes starch, carboxymethylcellulose, carboxymethylcellulose calcium, croscarmellose sodium, carboxymethylstarch sodium, and L-hydroxypropylcellulose. The solvent include water for injection, alcohol, propylene glycol, macrogols, sesame oil, and corn oil.

The solubilizer includes polyethylene glycol, propylene glycol, D-mannitol, benzyl benzoate, ethanol, trisaminomethane, cholesteryl, triethanolamine, sodium carbonate, and sodium citrate.

The suspending agent includes surfactants such as stearyl triethanolamine, sodium lauryl sulfate, laurylaminopropionic acid, lecithin, benzalkonium chloride, benzethonium chloride, and glycerin monostearate, etc. and hydrophilic macromolecular substances such as polyvinyl alcohol, polyvinylpyrrolidone, carboxymethylcellulose sodium, methylcellulose, hydroxymethylcellulose, hydroxyethylcellulose, and hydroxypropylcellulose.

The isotonizing agent includes glucose,

D-sorbitol, sodium chloride, glycerin, D-mannitol, etc.

The buffer includes various buffer solutions such as phosphate, acetate, carbonate, and citrate.

The anallgesic includes benzyl alcohol.

5       The preservative includes paraoxybenzoate, chlorobutanol, benzyl alcohol, phenethyl alcohol, dehydroacetic acid, and sorbic acid.

The antioxidant includes sulfite, and ascorbic acid.

10       The drug comprising the diphenylmethane derivative of this invention and it's medicinally acceptable salt have an excellent MIP-1 $\alpha$ /RANTES receptor antagonism and, therefore, they are useful as an medicament for mammals (e.g. humans, dogs, cats, rats, mice, bovines,  
15       etc.) for preventing or treating viral diseases or infectionary diseases (e.g. acute viral encephalitis, acute bacterial meningitis, Hericobacter pirolli infectious disease, pneumonia, hapatitis A, hepatitis B, hepatitis C, herpes simplex virus infectious  
20       disease, vesicle-strip blister virus infectious disease, HIV infectious disease (AIDS), influenza infectious disease, invasive staphylococcosis, tuberculosis, etc.), tumors (e.g. bladder cancer, mammary cancer, cervical carcinoma, chronic lymphatic  
25       leukemia, chronic myelocytic leukemia, colon cancer, multiple myeloma, malignant myeloma, prostatic cancer, lung cancer, stomach cancer, Hodgkin's disease, etc.), allergic diseases (e.g. bronchial asthma, atopic dermatitis, allergic rhinitis, etc.), inflammatory  
30       disease (e.g. arteriosclerosis, arterial sclerosis broken out after heart transplantation, (chronic) rheumatism, etc.), diabetic diseases (e.g. diabetes, diabetic nephropathy, diabetic complication, diabetic retinopathy, diabetic microangiopathy, etc.), central  
35       diseases (e.g. Alzheimer's disease, epilepsy, fever, ache, dementia, etc.), hyperlipemia,

hyperchlosterolemia, thrombocytopenia due to dialysis, spinal cord injury, osteoporosis, ulcerative colitis, peptic ulcer, sepsis (shock), reperfusion disorder of lung and heart, unstable angina pectoris, transient  
5 ischemic attack, valvular disease of heart, rejection after organ transplantation, retinosis after angioplasty, systematic lupus erythematosus, multiple sclerosis, renal failure, endometriosis, fibroid lung, adult respiratory distress syndrome, cardiac  
10 dysrhythmia, etc. Particularly, they are useful for preventing or treating allergic diseases, inflammatory diseases or multiple sclerosis.

The compound used for MIP-1 $\alpha$ /RANTES receptor antagonism of this invention is low toxic and has a low  
15 risk of side effect. The oral acute toxicity (LD<sub>50</sub>) of the compound of this invention in rats is not less than 100 mg/kg.

[Mode of Working the Invention]

The following reference, working, formulation and  
20 test examples are intended to describe this invention in further detail, but they are mere examples and should by no means be construed as defining the scope of the invention. Thus, various modifications can be made without departing from the scope of the invention.

25 In the following reference and working examples, the term "room temperature" means any temperature within the range of 0 to 30°C. The organic solvents were dried over anhydrous magnesium sulfate or anhydrous sodium sulfate. "%" means percent by weight  
30 otherwise specified. The other symbols have the following meanings.

s:	singlet
d:	doublet
t:	triplet
35 q:	quartet
m:	multiplet

br: broad  
J: coupling constant  
Hz: Herz  
CDCl<sub>3</sub>: deuteriochloroform  
5 THF: tetrahydrofuran  
DMF: N,N-dimethylformamide  
DMSO: dimethyl sulfoxide  
10 <sup>1</sup>H-NMR: proton nuclear magnetic resonance (The  
sample was measured in a free form and  
when a conformational isomer existed,  
the only main peak was read.)  
DMEM: Dulbecco's modified Eagle's medium  
PBS: phosphate buffered saline



## [Examples]

## Reference Example 1-1:

## 3,3-Diphenyl-3-formylpropionitrile

To a solution of diphenylacetaldehyde (1 g) in  
5 tetrahydrofuran (10 ml) was added dropwise slowly a  
suspension of 60% sodium hydride (0.25 g) in  
tetrahydrofuran (5 ml) under ice-cooling and stirring.  
After completion of dropwise addition, the mixture was  
further stirred for 20 minutes. Then,  
10 bromoacetonitrile (0.41 ml) was added and the mixture  
was further stirred for 30 minutes. The reaction  
mixture was poured into ice-water and the oil that had  
separated out was extracted with ethyl acetate. The  
organic layer was taken, washed with water, dried over  
15 anhydrous sodium sulfate, and concentrated to dryness.  
The residue was purified by silica gel column  
chromatography to provide the titled compound (0.85 g)  
as colorless oil.

## Reference Example 1-2:

## 20 4,4-Diphenyl-4-formylbutyronitrile

Diphenylacetaldehyde (25.6 g), acrylonitrile (12.5  
ml) and DBU (2.5 g) were stirred in isopropyl alcohol  
(250 ml) with warming at 70°C for 6 hours. The  
reaction mixture was concentrated to dryness and the  
25 residue was purified by silica gel column  
chromatography. The crude crystal crop obtained was  
washed with isopropyl ether to provide the titled  
compound (19.8 g) as colorless prisms.

The structural formulas and NMR spectra of the  
30 respective compounds are shown in Table 1.

## Reference Example 2-1:

## Ethyl 5-cyano-4,4-diphenyl-2-pentenoate

3,3-Diphenyl-3-formylpropionitrile (0.85 g) and  
(carboethoxymethylene)triphenylphosphorane (1.46 g)  
35 were heated in chloroform (20 ml) under reflux for 7  
hours. The reaction mixture was then concentrated to

dryness and the residue was purified by silica gel column chromatography to provide the titled compound (0.7 g) as colorless oil.

5 The compound of Reference Example 2-2 was synthesized in the same manner as Reference Example 2-1.

Reference Example 2-2:

Ethyl 6-cyano-4,4-diphenyl-2-hexenoate

10 The structural formulas and NMR spectra of the above compounds are shown in Table 2.

Reference Example 3-1:

(4-Chlorophenyl)phenylacetonitrile

15 To a mixture of mandelonitrile (5 g) and chlorobenzene (15.7 g) was added sulfuric acid (9.8 ml) dropwise while the temperature of the mixture was maintained at 5°C - 10°C. After completion of dropwise addition, the mixture was stirred for another 1.5 hours. The reaction mixture was poured into ice-water and the syrup that had separated out was extracted with  
20 ethyl acetate. The extract was washed with saturated aqueous sodium hydrogen carbonate solution, dried over anhydrous sodium sulfate, and concentrated to dryness. The residue was purified by silica gel column chromatography to provide the titled compound (3.6 g)  
25 as pale yellow syrup.

The compounds of Reference Examples 3-2 and 3-3 were synthesized in the same manner as Reference Example 3-1.

Reference Example 3-2:

30 (4-Methoxyphenyl)phenylacetonitrile

Reference Example 3-3:

Bis(4-chlorophenyl)acetonitrile

The structural formulas and NMR spectra of the respective compounds are shown in Table 3.

35 Reference Example 4-1:

Ethyl 4-cyano-4,4-diphenylbutyrate

To a solution of diphenylacetonitrile (28 g) in ethanol (100 ml) were added DBU (6 ml) and ethyl acrylate (30 ml). The mixture was heated and stirred at 80°C for 16 hours. After cooling, 2N-hydrochloric acid (200 ml) was added and the mixture was extracted with isopropyl ether. The organic extract was washed with water, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The crude crystal crop was recrystallized from hexane-isopropyl ether to provide the titled compound (34 g).

The compounds of Reference Example 4-2 - 4 were synthesized in the same manner as Reference Example 4-1.

Reference Example 4-2:

Ethyl 4-(4-chlorophenyl)-4-cyano-4-phenylbutyrate

Reference Example 4-3:

Ethyl 4-cyano-4-(4-methoxyphenyl)-4-phenylbutyrate

Reference Example 4-4:

Ethyl 4,4-bis(4-chlorophenyl)-4-cyanobutyrate

Reference Example 4-5:

Ethyl 5-cyano-5,5-diphenylpentanoate

To a stirring solution of diphenylacetonitrile (1 g) in tetrahydrofuran (10 ml) was added 60% sodium hydride (0.25 g) in small portion under ice-cooling. After completion of dropwise addition, the mixture was stirred for 20 minutes. Then, ethyl 4-bromobutyrate (0.94 ml) was added dropwise under ice-cooling and the mixture was further stirred at room temperature for 15 minutes. The reaction mixture was poured into ice-water and the organic layer that had separated out was extracted with ethyl acetate. The extract was washed with water, dried over anhydrous sodium sulfate, and concentrated to dryness. The residue was purified by silica gel column chromatography to provide the titled compound (0.87 g) as colorless oil.

Reference Example 4-6:

#### Ethyl 5-cyano-4,4-diphenylpentanoate

To a solution of ethyl 5-cyano-4,4-diphenyl-2-pentenoate (0.7 g) in ethanol (20 ml) was added 10% palladium-on-carbon (0.24 g), and the mixture was  
5 reduced by catalytic hydrogenation at atmospheric pressure and at room temperature. The catalyst in the reaction mixture was filtered off and the filtrate was concentrated to dryness. The residue was purified by silica gel column chromatography to provide the titled  
10 compound (0.6 g) as colorless oil.

The compound of Reference Example 4-7 was synthesized in same manner as Reference Example 4-6. Reference Example 4-7:

#### Ethyl 6-cyano-4,4-diphenylhexanoate

15 The structural formulas and NMR spectra of the respective compounds are shown in Table 4. Reference Example 5-1:

#### 5-Amino-4,4-diphenylpentanol

To a stirred solution of ethyl 4-cyano-4,4-diphenylbutyrate (1.2 g) in tetrahydrofuran (30 ml) was  
20 added lithium aluminum hydride (0.44 g) in small portion under ice-cooling. After completion of dropwise addition, the mixture was heated and stirred at 60°C for 3 hours. The reaction mixture was then  
25 cooled with ice again, water (1 ml) and 15% aqueous sodium hydroxide (3 ml) were added in succession. The insoluble matter that had separated out was filtered off and the filtrate was extracted with ethyl acetate and saturated aqueous sodium hydrogen carbonate. The  
30 organic layer was taken, washed with water, dried over anhydrous sodium sulfate, and concentrated to dryness. The residue was washed with isopropyl ether to provide the titled compound (0.82 g) as colorless powder.

The compounds of Reference Examples 5-2 - 7 were  
35 synthesized in the same manner as Reference Example 5-1.

Reference Example 5-2:

5-Amino-4-(4-chlorophenyl)-4-phenylpentanol

Reference Example 5-3:

5-Amino-4-(4-methoxyphenyl)-4-phenylpentanol

5 Reference Example 5-4:

5-Amino-4,4-bis(4-chlorophenyl)pentanol

Reference Example 5-5:

6-Amino-5,5-diphenylhexanol

Reference Example 5-6:

10 6-Amino-4,4-diphenylhexanol

Reference Example 5-7:

7-Amino-4,4-diphenylheptanol

The structural formulas and NMR spectra of the respective compounds are shown in Table 5.

15 Reference Example 6-1:

5-Formylamino-4,4-diphenylpentanol

5-Amino-4,4-diphenylpentanol (10 g) was dissolved in formic acid (80 ml) followed by addition of acetic anhydride (13 ml). The mixture was stirred at room temperature for 4 hours and concentrated to dryness. The residue was partitioned between chloroform and water. The water layer was made basic with aqueous ammonia and extracted with chloroform. The extracts were dried over anhydrous sodium sulfate and concentrated to dryness. The residue was dissolved in ethanol (30 ml) and the solution was stirred in 1N-aqueous sodium hydroxide (20 ml) at room temperature for 20 minutes. The reaction mixture was diluted with water and the crystals that separated out were collected by filtration. The crystal was washed serially with water and ethyl acetate to provide the titled compound (9 g) as colorless powder.

25 The compounds of Reference Example 6-2 - 7 were synthesized in the same manner as Reference Example 6-1.

35

Reference Example 6-2:

4-(4-Chlorophenyl)-5-formylamino-4-phenylpentanol

Reference Example 6-3:

5-Formylamino-4-(4-methoxyphenyl)-4-phenylpentanol

5 Reference Example 6-4:

4,4-Bis(4-chlorophenyl)-5-(formylamino)pentanol

Reference Example 6-5:

6-Formylamino-5,5-diphenylhexanol

Reference Example 6-6:

10 6-Formylamino-4,4-diphenylhexanol

Reference Example 6-7:

7-Acetylamino-4,4-diphenylheptanol

The structural formulas, physical properties, and  
NMR spectra of the above compounds are shown in Table  
15 6.

Reference Example 7-1:

5-Formylamino-1-iodo-4,4-diphenylpentane

To a solution of 5-formylamino-4,4-  
diphenylpentanol (38.3 g) in methylene chloride (600  
20 ml) were added p-toluenesulfonyl chloride (29.2 g),  
triethylamine (15 g), and 4-(dimethylamino)pyridine  
(catalytic amount). The mixture was stirred at room  
temperature for 4 hours and concentrated to dryness.  
The residue was stirred with Sodium Iodide (46.6 g) in  
25 acetone (600 ml) for 2 hours at 50°C. The reaction  
mixture was concentrated to dryness and the residue was  
diluted with ethyl acetate and water. The organic  
layer was taken, washed with an aqueous solution of  
sodium thiosulfate, dried over anhydrous sodium  
30 sulfate, and concentrated to dryness. The residue was  
purified by silica gel column chromatography to provide  
the titled compound (46.5 g) as yellow syrup.

The compounds of Reference Example 7-3 - 7 and 7-9  
were respectively synthesized in the same manner as

35 Reference Example 7-1.

Reference Example 7-2:

1-Iodo-4,4-diphenyl-5-(tosylamino)pentane

A mixture of 5-amino-4,4-diphenylpentanol (1 g),  
p-toluenesulfonyl chloride (1.65 g), triethylamine (1.2  
ml), and 4-(dimethylamino)pyridine (catalytic amount)  
5 in methylene chloride (20 ml) were stirred at room  
temperature for overnight. The reaction mixture was  
concentrated to dryness and the residue was stirred  
with sodium iodide (0.7 g) in acetone (25 ml) at 50°C  
for 24 hours. The reaction mixture was concentrated to  
10 dryness and the residue was diluted with ethyl acetate  
and water. The separated organic layer, was dried over  
anhydrous sodium sulfate and concentrated to dryness to  
provide the titled compound (1 g) as light-yellow  
powder.

15 The compound of Reference Example 7-8 was  
synthesized in the same manner as Reference Example 7-  
2.

Reference Example 7-3:

4-(4-Chlorophenyl)-5-formylamino-1-iodo-4-phenyl-  
20 pentane

Reference Example 7-4:

5-Formylamino-1-iodo-4-(4-methoxyphenyl)-4-  
phenylpentane

Reference Example 7-5:

25 4,4-bis(4-chlorophenyl)-5-formylaminopentyl-1-  
tosylate

Reference Example 7-6:

6-Formylamino-1-iodo-5,5-diphenylhexane

Reference Example 7-7:

30 6-Formylamino-1-iodo-4,4-diphenylhexane

Reference Example 7-8:

1-Iodo-4,4-diphenyl-6-(tosylamino)hexane

The structural formulas, physical properties, and  
NMR spectra of the respective compounds are shown in  
35 Table 7.

Reference Example 7-9:

## 7-Acetylamino-1-iodo-4,4-diphenylheptane

## Reference Example 8:

## 7-(2-Tetrahydropyranyloxy)-4,4-diphenylheptanenitrile

5 A solution of 6-cyano-4,4-diphenyl-1-hexanoic acid (12.5 g) in THF (85 ml) was added to a suspension of sodium borohydride (1.97 g) in THF (85 ml) at room temperature and stirred for 10 minutes. To the reaction mixture was added a solution of iodine (5.46 g) in THF (85 ml) under ice-cooling and the mixture was stirred for 1 hour. 3N-hydrochloric acid (20 ml) was added and the reaction mixture was concentrated under reduced pressure. The obtained residue was dissolved in ethyl acetate-water. The organic layer was separated, washed serially with water and a saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure to provide 6-cyano-4,4-diphenyl-1-hexanol (13 g). To a solution of the obtained alcohol (13 g) in dichloromethane (150 ml) were added p-toluenesulfonic acid monohydrate (catalytic amount) and 3,4-dihydro-2H-pyran (4.98 g) under ice-cooling and stirred for 15 hours. The reaction mixture was concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography eluting with hexane-ethyl acetate (4:1) to give the titled compound (10 g) as an oil.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.20-1.40(2H,m), 1.41-1.90(6H,m), 1.91-2.08(2H,m), 2.08-2.20(2H,m), 2.40-2.57(2H,m), 3.26-3.39(1H,m), 3.40-3.52(1H,m), 3.60-3.73(1H,m), 3.74-3.88(1H,m), 4.49(1H,br s), 7.02-7.40(10H,m)

## Reference Example 9:

## 1-Formylamino-7-(2-tetrahydropyranyloxy)-4,4-diphenylheptane

35 A solution of 7-(2-tetrahydropyranyloxy)-4,4-diphenylheptanenitrile (16.8 g) in THF (100 ml) was



added to a suspension of lithium aluminum hydride (4.3 g) in THF (150 ml) under ice-cooling and stirred at 60°C for 8 hours. To the reaction mixture was added an aqueous 1N-sodium hydroxide solution and the precipitate that separated out was filtered off and the filtrate was concentrated under reduced pressure. The residue was dissolved in ethyl acetate-water and separated. The organic layer was washed serially with water and a saturated aqueous sodium chloride solution. After drying, the solvent was distilled off under reduced pressure to provide 7-(2-tetrahydropyranyloxy)-4,4-diphenylheptanamine (17 g).

A solution of the obtained amine (3.7 g) in pyridine (25 ml) was added to a solution of formic acid in chloroform (2M, 20 ml) followed by addition of 1,3-dicyclohexylcarbodiimide (4, 12 g) in chloroform (25 ml) with stirring under ice-cooling and the mixture was stirred for 4 hours. The reaction mixture was concentrated under reduced pressure and the precipitate that had separated out was filtered off. The filtrate was concentrated under reduced pressure. The obtained residue was dissolved in ethyl acetate-water and the organic layer was separated, washed serially with water and a saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography eluting with hexane-ethyl acetate (1:2) to give the titled compound (1.4 g) as an oil.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.10-1.37(2H,m), 1.40-1.90(8H,m), 2.05-2.20(4H,m), 3.05-3.40(3H,m), 3.40-3.53(1H,m), 3.56-3.75(1H,m), 3.75-3.90(1H,m), 4.49(1H,br s), 5.20-5.60(1H,br), 7.05-7.33(10H,m), 8.12(1H,d)

Reference Example 10:

1-Formylamino-7-iodo-4,4-diphenylheptane

To a solution of 1-formylamino-7-(2-

tetrahydropyranyloxy)-4,4-diphenylheptane (1.4 g) in methanol (20 ml) was added p-toluenesulfonic acid monohydrate (catalytic amount) at room temperature and the mixture was stirred for 3 hours. The reaction mixture was concentrated under reduced pressure to provide 1-formylamino-7-hydroxy-4,4-diphenylheptane (1.2 g) as an oil.

The obtained oily substance was dissolved in dichloromethane (20 ml). To the solution were added a mixture of triethylamine (1 ml), 4-dimethylaminopyridine (catalytic amount), and p-toluenesulfonylchloride (687 ml) and the mixture was stirred for 3 hours. The reaction mixture was concentrated under reduced pressure and the obtained residue was dissolved in ethyl acetate-1N hydrochloric acid. The organic layer was separated, washed serially with water and saturated aqueous sodium chloride, and dried. The solvent was distilled off under reduced pressure to give 7-formylamino-4,4-diphenylheptyl 7-p-toluenesulfonate (1.3 g) as an oil.

To a solution of the obtained tosylate (1.3 g) in acetone (20 ml) was added sodium iodide (66 mg) and the mixture was stirred at 50°C for 4 hours. The reaction mixture was concentrated under reduced pressure and the obtained residue was dissolved in ethyl acetate-water. The organic layer was separated, washed serially with water and saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography eluting with hexane-ethyl acetate (1:1) to give the titled compound (1.4 g). Melting point: 119°C - 121°C.

Reference Example 11-1:

1-Benzyl-4-[3,5-bis(trifluoromethyl)phenyl]-4-hydroxypiperidine

To a solution of 3,5-bis(trifluoromethyl)-

bromobenzene (1.17 g) in THF (10 ml) was added  
magnesium (97 mg) and stirred under a argon stream at  
60°C for 2 hours. To the thus prepared Grignard  
reagent was added 1-benzyl-4-piperidone (379 mg) in THF  
5 (2 ml) and the mixture was stirred for 30 minutes.  
Saturated aqueous ammonium chloride solution was added  
to the reaction mixture and the mixture was extracted  
with ethyl acetate. The organic extract was washed  
serially with water and saturated aqueous sodium  
10 chloride and dried. The solvent was distilled off  
under reduced pressure. The obtained residue was  
purified by silica gel column chromatography eluting  
with ethyl acetate-hexane (1:4). The solvent was  
distilled off to give the titled compound (620 mg).  
15 Melting point: 89°C - 90°C

The compounds of Reference Examples 11-2 and 11-3  
were synthesized in a manner similar to that described  
above.

Reference Example 11-2:

20 1-Benzyl-4-(4-trifluoromethylphenyl)-4-  
hydroxypiperidine  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.64-1.85(3H,m), 2.16(2H,dt),  
2.46(2H,dt), 2.71(2H,d), 3.59(2H,s), 7.20-7.39(5H,m),  
7.62(4H,ABq)

25 Reference Example 11-3:

1-Benzyl-4-(3,5-dichlorophenyl)-4-  
hydroxypiperidine

Melting point: 75°C - 77°C

Reference Example 12-1:

30 4-[3,5-Bis(trifluoromethyl)phenyl]-4-  
hydroxypiperidine

To a solution of 1-benzyl-4-[3,5-  
bis(trifluoromethyl)phenyl]-4-hydroxypiperidine (1 g)  
in methanol (5 ml) was added 10% palladium-on-carbon  
35 (100 mg) and the mixture was stirred under a hydrogen  
atmosphere at room temperature for 2 hours. The

catalyst was filtered off and the filtrate was concentrated under reduced pressure to give the titled compound (600 mg).

Melting point: 209°C - 210°C

5 Reference Example 12-2:

In a manner similar to Reference Example 12-1, 4-(4-trifluoromethylphenyl)-4-hydroxypiperidine was synthesized.

Melting point: 115°C - 116°C

10 Reference Example 13

4-(3,5-Dichlorophenyl)-4-hydroxypiperidine

To a mixture of 1-benzyl-4-(3,5-dichlorophenyl)-4-hydroxypiperidine (200 ml) and potassium carbonate (276 mg) in toluene (5 ml) was added chloroethyl carbonate (217 mg) and the mixture was stirred at 60°C for 2 days. Water was added to the reaction mixture and extracted with ethyl acetate. The organic extract was washed serially with water and a saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography eluting with hexane-ethyl acetate (4:1) to give [4-(3,5-dichlorophenyl)-1-ethoxycarbonylpiperidin-4-yl] ethyl carbonate (190 mg) as an oil.

To a solution of the carbonate (190 mg) in ethanol (5 ml) was added 4N-potassium hydroxide solution (5 ml) and the mixture was heated under reflux for 15 hours. The solvent was distilled off under reduced pressure. To the residue were added water and ethyl acetate and stirred well. The organic layer was separated, washed serially with water and a saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure to give the titled compound (135 mg).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.60-1.75(4H,m), 1.97(2H,dt), 2.91-

3.16(4H,m), 7.26(1H,d), 7.40(2H,d)

Reference Example 14:

4-(4-Chlorophenyl)piperidine

To a solution of 4-(4-chlorophenyl)-4-  
5 hydroxypiperidine (478 mg) in acetic acid (5 ml) was  
added sulfuric acid (0.5 ml) and the mixture was  
stirred at room temperature for 3 hours. The reaction  
mixture was made basic with 4N-sodium hydroxide and  
extracted with ethyl acetate. The organic layer was  
10 washed serially with water and a saturated aqueous  
sodium chloride solution and dried. The solvent was  
distilled off under reduced pressure to give 4-(4-  
chlorophenyl-1,2,5,6-tetrahydropyridine (350 mg). To a  
solution of 4-(4-chlorophenyl)-1,2,5,6-  
15 tetrahydropyridine (250 mg) in methanol (5 ml) and 4N-  
hydrochloric acid (2 ml) was added 10% palladium-on-  
carbon (150 mg) and the mixture was stirred under a  
hydrogen atmosphere at room temperature for 2.5 hours.  
The catalyst was filtered off and the filtrate was made  
20 basic with a 4N-aqueous sodium hydroxide solution  
followed by extraction with ethyl acetate. The organic  
extract was washed serially with water and a saturated  
aqueous sodium chloride solution and dried. The  
solvent was distilled off under reduced pressure to  
25 give the titled compound (180 mg) as an oily substance.  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.60(2H,dq), 1.70-2.05(3H,m), 2.50-  
2.83(3H,m), 3.19(2H,dr d), 7.15(2H,d), 7.26(2H,d)

Reference Example 15:

4-(4-Chlorophenyl)-4-hydroxyhexamethyleneimine

30 1) 1-Benzyl-4-(4-chlorophenyl)-4-  
hydroxyhexamethyleneimine

In a similar manner to Reference Example 11-1, the  
titled compound was synthesized from 1-  
benzylhexamethylenimin-4-one and 4-chlorobromobenzene.

35 <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.49-2.20(6H,m), 2.32-2.59(2H,m),  
2.60-2.73(1H,m), 2.82-3.13(2H,m), 3.66(2H,ABq), 7.21-

7.43(9H,m)

2) 4-(4-Chlorophenyl-4-hydroxyhexamethyleneimine

To a mixture of 1-benzyl-4-(4-chlorophenyl)-4-hydroxyhexamethyleneimine (472 mg) and potassium carbonate (414 mg) in toluene (3 ml) was added ethyl vinylcarbonate (426 mg) under ice-cooling and stirred at room temperature for 30 minutes. The reaction mixture was diluted with water and extracted with ethyl acetate. The organic extract was washed serially with water and a saturated aqueous sodium chloride solution and dried. The solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography eluting with hexane-ethyl acetate (6:1) to give [4-(4-chlorophenyl)-1-vinylloxycarbonyl-hexamethyleneimin-4-yl] vinyl carbonate (400 mg). To a solution of the product in ethanol (5 ml) was added 4N-aqueous potassium hydroxide solution (5 ml) and the mixture was stirred at 60°C for 4 hours. The solvent was distilled off under reduced pressure. To the residue were added water and ethyl acetate and stirred well. The organic layer was separated washed serially with water and a saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure to give the titled compound (150 mg).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)δ: 1.50-2.85(9H,m), 2.85-3.01(1H,m), 3.17-3.30(1H,m), 3.30-3.50(1H,m), 7.23-7.45(4H,m)

Reference Example 16:

Ethyl 4-cyano-4-phenyl-4-(2-pyridyl)butanoate

To a suspension of 60% sodium hydride (13.2 g) in DMF (400 ml) was added a solution of phenylacetonitrile (35.1 g) in DMF (20 ml) under ice-cooling and the mixture was stirred for 30 minutes. A solution of 2-bromopyridine (47.4 g) in DMF (20 ml) was added to the mixture under ice-cooling and stirred at room temperature for 2 hours. The reaction mixture was

diluted with water and extracted with ethyl acetate. The organic extract was washed serially with water and a saturated aqueous sodium chloride solution and dried. The solvent was distilled off under reduced pressure to give 22 g of phenyl(2-pyridyl)acetonitrile  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 5.32(1H,s), 7.20-7.50(7H,m), 7.70(1H,dt), 8.60(1H,dd)

To a solution of phenyl(2-pyridyl)acetonitrile (19.4 g) in ethanol (250 ml) were added ethyl acrylate (13 g) and 1,8-diazabicyclo[5,4,0]-7-undecene (1.5 ml) and the mixture was heated to reflux for 5 hours. The solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography eluting with hexane-ethyl acetate (4:1 - 3:1) to give the titled compound as an oily substance.  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.23(3H,t), 2.40-2.52(2H,m), 2.70-2.92(1H,m), 2.93-3.15(1H,m), 4.10(2H,q), 7.19-7.55(8H,m), 7.68(1H,dt), 8.63(1H,d)

Reference Example 17:

20 4-Cyano-4-phenyl-4-(2-pyridyl) butanoic acid

To a solution of ethyl 4-cyano-4-phenyl-4-(2-pyridyl)butanoate (2.9 mg) in ethanol (10 ml) was added 1N-aqueous sodium hydroxide solution (15 ml) and the mixture was stirred at 60°C for 30 minutes. The reaction mixture was concentrated under reduced pressure and neutralized with 1N-hydrochloric acid. The aqueous layer was extracted with ethyl acetate. The organic extract was washed serially with water and a saturated aqueous sodium chloride solution and dried. The solvent was distilled off under reduced pressure to give the titled compound (2.7 g).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 2.45-2.60(2H,m), 2.78(1H,ddd), 3.06(1H,ddd), 7.20-7.55(7H,m), 7.68(1H,dt), 8.63(1H,d)

Reference Example 18:

35 N-[4-Cyano-4-phenyl-4-(2-pyridyl)butyl]-4-(4-chlorophenyl)-4-hydroxypiperidine

To a solution of 4-cyano-4-phenyl-4-(2-pyridyl) butyric acid (1.33 g), 4-(4-chlorophenyl)-4-hydroxypiperidine (1.3 g), and diethyl phosphorocyanidate (982 mg) in DMF (20 ml) was added triethylamine (606 mg) at room temperature and the mixture was stirred for 3 hours. The reaction mixture was diluted with water and extracted with ethyl acetate. The organic extract was washed serially with water and a saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography eluting with ethyl acetate-hexane (1:1) to give the titled compound (1.5 g) as a noncrystalline powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.62-2.00(4H,m), 2.21(1H,s), 2.35-2.57(2H,m), 2.70-2.93(1H,m), 2.94-3.14(2H,m), 3.37-3.74(2H,m), 4.53(1H,br d), 7.19-7.53(12H,m), 7.68(1H,dt), 8.62(1H,d)

Reference Example 19:

5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2-phenyl-2-(2-pyridyl)pentylamine 2hydrochloride

To a suspension of lithium aluminum hydride (380 mg) and aluminum chloride (1.3 g) in ether (20 ml) was added N-[4-cyano-4-phenyl-4-(2-pyridyl)butyl]-4-(4-chlorophenyl)-4-hydroxypiperidine (465 mg) under ice-cooling and the mixture was stirred for 20 minutes. To the reaction mixture was added 1N-aqueous sodium hydroxide solution and the resulting solution was extracted with ethyl acetate. The organic extract was washed serially with water and a saturated aqueous sodium chloride solution and dried. The solvent was distilled off under reduced pressure. The obtained residue was purified silica gel column chromatography eluting with ethyl acetate-hexane (1:1). The solvent was distilled off and the residue was treated with 4N-hydrochloric acid/ethyl acetate to give the titled



compound (250 mg) as a noncrystalline powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05-1.42(2H,m), 1.45-1.90(5H,m),  
1.90-2.13(2H,m), 2.14-2.40(6H,m), 2.66(2H,br d),  
3.39(2H,ABq), 6.98-7.47(11H,m), 7.55(1H,dt), 8.57(1H,d)

5 Reference Example 20:

4-Cyano-4,4-diphenyl butanoic acid

To a solution of ethyl 4-cyano-4,4-diphenylbutanoate (16.1 g) in THF (6 ml) was added 1N-sodium hydroxide solution (60.5 ml) and the mixture was  
10 stirred at room temperature for 16 hours. The solution was made acidic with concentrated hydrochloric acid, extracted with ethyl acetate and dried. The solvent was distilled off to give an oily residue. The residue was crystallized from isopropyl ether to give the  
15 titled compound (12.0 g).

Melting point: 164°C - 165°C

Reference Example 21:

4-(4-Chlorophenyl)-1-(4-cyano-4,4-diphenylbutyryl)-4-hydroxypiperidine

To a solution of 4-cyano-4,4-diphenylbutanoic acid  
20 (8.0 g), 4-(4-chlorophenyl)-4-hydroxypiperidine (6.4 g), and diethylphosphoro cyanidate (4.6 ml) in DMF (75 ml) was added triethylamine (8.4 ml) at 0°C and the mixture was stirred at room temperature for 2 hours.  
25 The reaction mixture was poured into pure water (500 ml) and the solid that separated out was collected by filtration. The solid was dissolved in ethyl acetate, washed with a saturated aqueous sodium chloride solution, and dried. The solvent was distilled off  
30 under reduced pressure. The solid residue was suspended with ether and collected by filtration to give the titled compound (12.6 g). Melting point: 205°C - 206°C

Reference Example 22:

35 1-(5-Amino-4,4-diphenylpentanoyl)-4-(4-chlorophenyl)-4-hydroxypiperidine

To the suspension of 4-(4-chlorophenyl)-1-(4-cyano-4,4-diphenylbutyryl)-4-hydroxypiperidine (6.9 g) in saturated ammonia ethanol solution (500 ml) was added Reney-Cobalt catalyst (7 g) and the mixture was  
5 reacted under 5 atmospheric pressure of hydrogen for 8 hours. The catalyst was filtered off and the filtrate was concentrated under reduced pressure to give an oily residue (5.4 g).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.57-1.89(4H,m), 1.99-2.08(2H,m),  
10 2.43-2.53(2H,m), 3.01(1H,dt), 3.25(2H,s), 3.22-3.35(2H,m), 4.51(1H,br d), 7.15-7.37(14H,m)

Reference Example 23:

4-Cyano-4,4-diphenyl-1-butanol

To a solution of ethyl 4-cyano-4,4-diphenylbutanoate (44.0 g) in THF (440 ml) was added  
15 carefully lithium tetrahydroborate (3.9 g) at 0°C. The reaction mixture was allowed to warm to room temperature and stirred for 2 days. The reaction mixture was poured into a cold 1N-hydrochloric acid  
20 (440 ml) and extracted with ethyl acetate. The organic extract was washed with saturated aqueous sodium chloride solution and dried. The solvent was distilled off under reduced pressure. The residue was purified  
25 by silica gel chromatography eluting with ethyl acetate-hexane (1:2) to give the titled compound (34.6 g) as an oil.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.62-1.76(2H,m), 2.47-2.55(2H,m),  
3.69(2H,t), 7.28-7.42(10H,m)

Reference Example 24:

30 1-Bromo-4-cyano-4,4-diphenylbutane

To a suspension of triphenylphosphine (27.5 g) in acetonitrile (100 ml) was added bromine (5.2 ml) dropwise at 0°C. After completion of dropwise addition, a solution of 4-cyano-4,4-diphenyl-1-butanol  
35 (25.1 g) in acetonitrile (40 ml) was added to the reaction mixture at 0°C and the mixture was stirred at

room temperature for 1 hour. The solvent was distilled off under reduced pressure and ether was added to the obtained residue. Triphenylphosphineoxide was filtered off and the filtrate was washed with a saturated aqueous sodium chloride solution, dried, and concentrated under reduced pressure. The residue was purified by silica gel chromatography eluting with ether and crystallized from IPE to give the titled compound (25.6 g)

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.93-2.07(2H,m), 2.527-2.61(2H,m), 3.44(2H,t), 7.26-7.42(10H,m).

Reference Example 25:

4-(4-Chlorophenyl)-1-(4-cyano-4,4-diphenylbutyl)-4-hydroxypiperidine

To a solution of 1-bromo-4-cyano-4,4-diphenylbutane (22.0 g) in acetonitrile (500 ml) were added 4-(4-chlorophenyl)-4-hydroxypiperidine (17.8 g), potassium carbonate (29.0 g), and potassium iodide (1.2 g) and the mixture was stirred at room temperature for 16 hours. The solvent was distilled off under reduced pressure. The residue was dissolved in ethyl acetate and washed with pure water. The organic layer was washed with a saturated aqueous sodium chloride solution and dried. The solvent was distilled off under reduced pressure. The residue was purified by silica gel chromatography eluting with ethyl acetate to give the titled compound (31.4 g) as a noncrystalline power.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.60-1.83(5H,m), 2.06(2H,dt), 2.30-2.49(6H,m), 2.71(2H,br d), 7.27-7.45(14H,m).

Reference Example 26:

1-Amino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane

To a solution of 4-(4-chlorophenyl)-1-(4-cyano-4,4-diphenylbutyl)-4-hydroxypiperidine (31.3 g) in saturated ammonium-ethanol solution (500 ml) was added

Raney-Co catalyst (20 g) and the mixture was stirred under 5 atmospheric pressure of hydrogen gas for 8 hours at 70°C. The solvent was distilled off under reduced pressure. The obtained residue was  
5 crystallized from ethyl acetate to give the titled compound (17.8 g).

Melting point: 116°C - 117°C

Reference Example 27:

2-Benzoylthiophenecyanohydrin

10 A mixture of 2-benzoylthiophene (10 g), trimethylcyanide (6 g), and zinc iodide (0.15 g) acetonitrile (50 ml) was stirred at 50°C for 16 hours. The solvent was distilled off under reduced pressure. 1N-Hydrochloric acid (60 ml) and ethanol (30 ml) were  
15 added to the residue and the mixture was stirred at 55°C for 2 hours. The reaction mixture was extracted with isopropyl ether and the organic extract was washed with saturated aqueous sodium hydrogen carbonate and saturated aqueous sodium chloride solution,  
20 respectively and dried. The solvent was distilled off under reduced pressure to give the titled compound (11.5 g).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 3.68(1H,br s), 6.98(1H,dd), 7.19(1H,dd), 7.34-7.46(4H,m), 7.59(2H,m).

25 Reference Example 28:

Phenyl-2-thienylacetonitrile

A solution of 2-benzoylthiophenecyanhydrin (250 mg) in ether (1 ml) and sodium borohydride (430 mg) were added to a solution of trifluoroacetic acid (5 ml)  
30 at 0°C and the mixture was stirred 15 hours at room temperature. The solvent was distilled off under reduced pressure. The residue was dissolved in 1N-aqueous sodium hydroxide and the water layer was extracted with ethyl acetate. The organic extract was  
35 washed with saturated aqueous sodium chloride and dried. The solvent was distilled off under reduced

pressure and the obtained residue was purified by silica gel column chromatography eluting with hexane-ethyl acetate (8:1) to give the titled compound (110 mg).

5 <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 5.35(1H,s), 6.97(1H,dd), 7.05-7.09(1H,m), 7.27(1H,dd), 7.32-7.44(5H,m).

Reference Example 29:

Ethyl 4-cyano-4-phenyl-4-(2-thienyl)butyrate

10 In a similar manner to Reference Example 4-1, the titled compound was synthesized from phenyl-2-thienylacetonitrile.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.23(3H,t), 2.41(1H,dd), 2.56(1H,dd), 2.80(2H,dt), 4.10(2H,q), 6.96(1H,dd), 7.00(1H,dd), 7.25-7.52(6H,m).

15 Reference Example 30:

5-Formylamino-4-phenyl-4-(2-thienyl)pentanol

In a similar manner to Reference Example 4-1, ethyl 4-cyano-4-phenyl-4-(2-thienyl)butyrate was reduced to obtain

20 5-amino-4-phenyl-4-(2-thienyl)pentanol. Then in a similar manner to Reference Example 6-1, the titled compound was obtained from

5-amino-4-phenyl-4-(2-thienyl)pentanol.

25 <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.21-1.59(2H,m), 1.81(1H,br s), 2.21(2H,t), 3.56(2H,t), 3.98(2H,dd), 4.12(2H,dd), 5.43(2H,br s), 6.85-7.00(2H,m), 7.10-7.40(6H,m), 8.11(1H,s).

Reference Example 31:

5-Formylamino-1-iodo-4-phenyl-4-(2-thienyl)pentane

30 By using iodination according to Reference Example 7-1, the titled compound was obtained from

5-formylamino-4-phenyl-4-(2-thienyl)pentanol.

35 <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.48-1.75(2H,m), 2.08-2.28(2H,m), 3.10(2H,t), 4.03(2H,dd), 5.18-5.65(1H,br m), 6.84-7.00(2H,m), 7.15-7.40(6H,m), 8.12(1H,d).

Reference Example 32:


## 4-Chloromandelonitorile

To a aqueous sodium hydrogensulfite (53.2 g) (400 ml) was added 4-chlorobenzaldehyde (60 g) and the mixture was stirred at 40°C for 1 hour, cooled by 0°C, and ether (250 ml) was added. Sodium cyanide (22.6 g) in water (100 ml) was added to the mixture, and the mixture was stirred at 0°C for 2 hours. The organic layer was separated and washed with saturated aqueous sodium chloride, and dried. The solvent was distilled off under reduced pressure to give the titled compound (65g).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 3.06(1H,br d), 5.53(1H,d), 7.38-7.52(4H,m).

## Example 1-1

5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane hydrochloride



To a solution of 5-formylamino-1-iodo-4,4-diphenylpentane (5 g) and 4-(4-chlorophenyl)-4-hydroxypiperidine (3.9 g) in acetonitrile (150 ml) was added potassium carbonate (7.7 g) and the mixture was stirred at 60°C for 15 hours. The solvent was distilled off under reduced pressure. Water and ethyl acetate were added to the obtained residue and stirred well. The organic layer was separated, washed serially with water and a saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography eluting with ethyl acetate. The solvent was distilled off and the residue was treated with 4N-hydrochloric acid/ethyl


acetate to give the titled compound (5.6 g) as a noncrystalline powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.18-1.40(2H,m), 1.58-1.92(3H,m),  
1.93-2.22(4H,m), 2.23-2.42(4H,m), 2.65(2H,br d),  
4.05(2H,d), 5.13(1H,br t), 7.10-7.28(14H,m),  
8.09(1H,d).

The compounds of 1-2 to 1-10 were synthesized in a manner similar to Example 1-1.

#### Example 1-2


5-[4-(4-Fluorophenyl)piperadin-1-yl]-1-formylamino-2,2-diphenylpentane dihydrochloride



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.20-1.40(2H,m), 2.10-2.40(4H,m),  
2.45(4H,t), 3.05(4H,t), 4.06(2H,d), 5.10(1H,br s),  
6.80-7.00(4H,m), 7.10-7.40(10H,m), 8.10(1H,d).

#### Example 1-3


1-Formylamino-5-(4-hydroxy-4-phenylpiperidino)-2,2-diphenylpentane hydrochloride



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.22-1.45(2H,m), 1.72(2H,br d), 1.80-  
2.28(7H,m), 2.30-2.50(4H,m), 2.65-2.80(2H,m),  
4.05(2H,d), 5.15-5.26(1H,br), 7.13-7.55(15H,m),  
8.10(1H,d).

#### Example 1-4


5-[4-(4-Trifluoromethylphenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane hydrochloride



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.26-1.42(2H,m), 1.67(2H,br d), 1.81-2.26(4H,m), 2.27-2.45(4H,m), 2.73(2H,br d), 4.05(2H,d), 5.09-5.20(1H,br t), 7.13-7.37(10H,m), 7.55-7.70, 8.09(1H,d).

#### Example 1-5


5-[4-[3,5-Bis (trifluoromethyl)phenyl]-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane hydrochloride



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.15-1.40(2H,m), 1.67(2H,br d), 1.90-2.24(4H,m), 2.25-2.45(4H,m), 2.69(3H,br d), 4.03(2H,d), 5.22(1H,br t), 7.05-7.40(10H,m), 7.75(1H,s), 7.97(2H,s), 8.04(1H,d).

#### Example 1-6

5-[4-(3,5-Dichlorophenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane hydrochloride




<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.15-1.40(2H,m), 1.64(2H,br d), 1.94-2.42(9H,m), 2.62-2.76(2H,m), 4.05(2H,d), 5.16(1H,br t), 7.10-7.43(13H,m), 8.08(1H,d).



## Example 1-7

5-[4-(4-Chlorophenyl)-1,2,3,6-tetrahydropyridin-1-yl]-1-formylamino-2,2-diphenylpentane hydrochloride




Recrystallization solvent: ethyl acetate/isopropyl ether

Melting point: 123°C - 125°C

Recrystallization solvent: ethyl acetate/isopropyl ether

## Example 1-8

1-Formylamino-2,2-diphenyl-5-(4-phenylpiperidino)pentane




Recrystallization solvent: ethyl ether/hexane

Melting point: 133°C - 135°C

## Example 1-9


5-[4-(4-Chlorophenyl)piperidino]-1-formylamino-2,2-diphenylpentane hydrochloride



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.15-1.40(2H,m), 1.50-2.50(11H,m), 2.87(2H,br d), 4.05(2H,d), 5.00-5.25(1H,br), 7.00-7.40(14H,m), 8.09(1H,d).

## Example 1-10


7-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-1-formylamino 4,4-diphenylheptane hydrochloride



$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.12-1.30(4H,m), 1.66(2H,br d), 1.77-2.22(7H,m), 2.22-2.43(4H,m), 2.56-2.72(2H,m), 3.22(2H,q), 5.40-5.64(1H,br), 7.10-7.35(12H,m), 7.42(2H,d), 8.08(1H,d).

Example 2-1

5-[4-(4-Fluorophenyl)-4-hydroxypiperidino]-1-formylamino-2,2-diphenylpentane hydrochloride



To a solution of 1-formylamino-5-iodo-2,2-diphenylpentane (1 g), 4-piperidone hydrochloride mono hydrate (450 mg) in acetonitrile (10 ml) was added potassium carbonate (845 g) and the mixture was stirred at 45°C for 2 days. The solvent was distilled off under reduced pressure. Water and ethyl acetate were added to the residue, and the mixture was stirred well. The organic layer was separated, washed with a saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography eluting with ethyl acetate-methanol (9:1) to give 1-formylamino-2,2-diphenyl-5-(4-piperidon-1-yl)pentane (570 mg) as a noncrystalline powder.

To a solution of 4-bromofluorobenzene (298 mg) in THF (5 ml) was added dropwise 1.6 M of n-butyl lithium hexane solution (1.25 ml) under an argon atmosphere at -78°C and the mixture was stirred for 20 minutes.


5 Anhydrous cerium chloride (520 mg) was added to the reaction mixture, and the mixture was stirred for another 45 minutes followed by addition of a solution of 1-formylamino-2,2-diphenyl-5-(4-piperidon-1-yl)pentane (125 mg) in THF (1 ml). The reaction  
10 temperature was raised to -10°C gradually. After 1.5 hours, water and 1N-sodium hydroxide solution were added to the reaction mixture and extracted with ethyl acetate. The organic extract was washed serially with water and a saturated aqueous sodium chloride solution,  
15 and dried. The solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography eluting with ethyl acetate-methanol (9:1). The solvent was distilled off and the obtained residue was treated with 4N-hydrochloric  
20 acid/ethyl acetate to give the titled compound (85 mg) as noncrystalline powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.18-1.42(2H,m), 1.50-1.95(3H,m), 2.00-2.23(4H,m), 2.24-2.44(4H,m), 2.70(2H,br d), 4.06(2H,d), 5.10-5.23(1H,br), 7.01(2H,t), 7.10-  
25 7.38(10H,m), 7.39-7.52(2H,m), 8.09(1H,d).

The compounds of Examples 2-2 and 2-3 were synthesized in a manner similar to Example 2-1.

#### Example 2-2


30 1-Formylamino-5-[4-hydroxy-4-(4-methoxyphenyl)piperidino]-2,2-diphenylpentane hydrochloride



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.15-1.45(2H,m), 1.50-2.20(7H,m),  
2.21-2.40(4H,m), 2.53-2.71(2H,m), 3.80(3H,s),  
4.05(2H,d), 5.12-5.22(1H,br), 6.87(2H,d), 7.10-  
7.45(12H,m), 8.09(1H,d).

5 Example 2-3


1-Formylamino-5-[4-hydroxy-4-(2-  
pyridyl)piperidino]-2,2-diphenylpentane dihydrochloride



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.20-1.53(2H,m), 1.63(2H,br d), 2.16-  
15 3.06(11H,m), 4.06(2H,d), 5.32-5.42(1H,br), 7.03-  
7.40(11H,m), 7.48(1H,t), 7.73(1H,dt), 8.11(1H,d),  
8.50(1H,d).

Example 3-1

20 1-Acetylamino-5-[4-(4-chlorophenyl)-4-  
hydroxypiperidino]-2,2-diphenylpentane hydrochloride




To a mixture of 1-amino-5-[4-(4-chlorophenyl)-4-  
hydroxypiperidino]-2,2-diphenylpentane (112 mg) (in  
ethyl acetate (3 ml)) was added a saturated aqueous  
sodium carbonate solution followed by addition of  
30 anhydrous acetic acid (24 mg) under vigorously stirring  
at 0°C and the mixture was stirred for 5 minutes. The  
organic layer was separated, washed serially with water  
and a saturated aqueous sodium chloride solution, and  
dried. The solvent was distilled off under reduced  
35 pressure. The obtained residue was purified by silica  
gel column chromatography eluting with ethyl acetate-

methanol (19:1). The solvent was distilled off and the residue was treated with 4N-hydrochloric acid/ethyl acetate to give the titled compound (45 mg) as a noncrystalline powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.20-1.42(2H,m), 1.68(2H,br d), 1.85(3H,s), 2.00-2.20(3H,m), 2.26-2.28(4H,m), 2.72(2H,br d), 3.98(2H,d), 5.02(1H,br t), 7.13-7.38(12H,m), 7.42(2H,d).

The compounds of Examples 3-2 to 3-12 were synthesized in a manner similar to Example 3-1.  
Example 3-2


1-Acetoacetyl-amino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane hydrochloride



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.20-1.40(2H,m), 1.64(2H,br d), 1.80-2.20(8H,m), 2.20-2.40(4H,m), 2.65(2H,br d), 3.26(2H,s), 4.02(2H,d), 6.40-6.53(1H,br), 7.14-7.44(14H,m).

Example 3-3


Ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]succinamate hydrochloride



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.17-1.38(5H,m), 1.65(2H,br d), 1.92-2.14(5H,m), 2.20-2.37(6H,m), 2.55-2.72(4H,m), 4.01(2H,d), 4.10(2H,q), 5.18(1H,br t), 7.16-7.38(12H,m), 7.43(2H,d).

Example 3-4


N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]succinamic acid



$^1\text{H-NMR}$  (DMSO- $d_6$ )  $\delta$ : 1.08-1.29(2H,m), 1.53(2H,br d),  
1.80-2.28(9H,m), 2.29-2.48(4H,m), 2.53-2.68(2H,m),  
3.89(2H,br d), 7.10-7.39(12H,m), 7.48(2H,d).

Example 3-5

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-ethylurea




Recrystallization solvent: ethyl acetate/hexane

Melting point: 142°C - 144°C

Example 3-6


N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]methanesulfonamide hydrochloride



$^1\text{H-NMR}$  (DCCl $_3$ )  $\delta$ : 1.20-1.36(2H,m), 1.60-1.80(3H,m),  
2.00-2.43(8H,m), 2.48(3H,s), 2.71(2H,br d), 3.82(2H,d),  
4.78-4.92(1H,br), 7.13-7.40(12H,m), 7.45(2H,d).

Example 3-7


Phenyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]carbamate



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.22-1.42(2H,m), 1.53-1.74(2H,m),  
1.96-2.40(9H,m), 2.19(2H,br d), 4.02(2H,d), 4.89(1H,br  
t), 6.95-7.08(2H,m), 7.10-7.46(17H,m).

#### Example 3-8


1-Acetylamino-5-[4-(4-chlorophenyl)-4-  
hydroxypiperidino]-2-phenyl-2-(2-pyridyl)pentane  
dihydrochloride



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.15-1.50(2H,m), 1.67(2H,br d),  
1.85(3H,s), 1.94-2.48(8H,m), 2.50-2.76(3H,m),  
3.87(1H,dd), 4.13(1H,dd), 6.58(1H,br t), 6.95-  
7.52(11H,m), 7.60(1H,dt), 8.57(1H,dt).

#### Example 3-9

Ethyl N-[5-[4-(4-chlorophenyl)-4-  
hydroxypiperidino]-2,2-diphenylpentyl]oxamate  
hydrochloride




<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.15-1.40(5H,m), 1.64(2H,br d), 1.71-  
2.18(5H,m), 2.19-2.38(4H,m), 2.56-2.69(2H,m),

4.05(2H,d), 4.26(2H,q), 6.72(1H,br t), 7.14-7.44(14H,m).

Example 3-10

5 Ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]malonamate hydrochloride

10




15

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.15-1.38(5H,m), 1.64(2H,d), 1.95-2.19(5H,m), 2.20-2.38(4H,m), 2.57-2.70(2H,m), 3.17(2H,s), 3.98-4.15(4H,m), 6.58(1H,br t), 7.16-7.45(14H,m).

Example 3-11

20 Ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]glutaramate

25




30

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.13-1.40(5H,m), 1.58-1.94(5H,m), 1.95-2.16(6H,m), 2.17-2.39(6H,m), 2.66(2H,br d), 4.01(2H,d), 4.09(2H,q), 5.05(1H,br t), 7.15-7.38(12H,m), 7.43(2H,d).

Example 3-12

Ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2-phenyl-2-(2-pyridyl)pentyl]succinamate dihydrochloride






<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.18-1.50(5H,m), 1.88-2.10(3H,m),  
2.10-2.48(8H,m), 2.49-2.74(6H,m), 3.89(1H,dd), 4.05-  
4.20(3H,m), 6.66(1H,br t), 7.05-7.37(11H,m),  
7.60(1H,dt), 8.56-8.62(1H,m).

#### Example 4-1

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-pentamethyleneurea hydrochloride




To a solution of phenyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl] carbamate (86 mg) and piperidine (43 mg) in DMF (1 ml) was added potassium carbonate (69 mg) and the mixture was stirred at room temperature for 1 hour. The reaction mixture was diluted with water and extracted with ethyl acetate. The organic extract was washed serially with water and a saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography eluting with ethyl acetate-methanol (20:1). The solvent was distilled off and the obtained residue was treated with 4N-hydrochloric acid/ethyl acetate to give the titled compound (80 mg) as a noncrystalline powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.20-1.60(2H,m), 1.66(2H,br d), 1.80-

2.20(5H,m), 2.20-2.40(4H,m), 2.67(2H,br d), 3.06-  
3.13(4H,m), 3.95(2H,s), 7.17-7.38(12H,m), 7.43(2H,d).

Example 4-2

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-  
5 diphenylpentyl]-3-(3-hydroxypropyl)urea hydrochloride



10


To a solution of phenyl N-[5-[4-(4-chlorophenyl)-  
4-hydroxypiperidino]-2,2-diphenylpentyl] carbamate (569  
mg) and 3-amino-1-propanol (113 mg) in DMF (2 ml) was  
added potassium carbonate (267 mg) and the mixture was  
15 stirred at room temperature for 16 hours. The reaction  
mixture was diluted with water and extracted with ethyl  
acetate. The organic extract was washed serially with  
water and a saturated aqueous sodium chloride solution,  
and dried. The solvent was distilled off under reduced  
20 pressure. The obtained residue was purified by silica  
gel column chromatography eluting with ethyl acetate-  
methanol (9:1). The solvent was distilled off and the  
residue was treated with 4N-hydrochloric acid/ethyl  
acetate to give the titled compound (600 mg) as a  
25 noncrystalline powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.15-1.40(2H,m), 1.40-1.72(4H,m),  
1.75-2.18(6H,m), 2.23-2.43(4H,m), 2.70(2H,br d),  
3.24(2H,q), 3.56(2H,t), 3.92(2H,d), 4.18(1H,br t),  
4.48(1H,br t), 7.18-7.48(14H,m).

30

Example 4-3

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-  
diphenylpentyl]-3-(4-hydroxybutyl)urea hydrochloride



5

To a solution of phenyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]carbamate (235 mg) and 4-amino-1-butanol (67 mg) in DMF (1 ml) was added potassium carbonate (138 mg) and the mixture was stirred at room temperature for 4 hours. The reaction mixture was diluted with water and extracted with ethyl acetate. The organic extract was washed serially with water and a saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography eluting with ethyl acetate-methanol (9:1). The solvent was distilled off and the residue was treated with 4N-hydrochloric acid/ethyl acetate to give the titled compound (205 mg) as a noncrystalline powder.

15

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.10-1.43(2H,m), 1.45-1.56(2H,m), 1.68(2H,d), 1.90-2.52(12H,m), 2.76(2H,br d), 3.07(2H,q), 3.61(2H,t), 3.94(2H,d), 4.08(1H,br t), 4.53(1H,br t), 7.14-7.48(14H,m).


20

The compounds of Examples 4-4 to 4-10 were synthesized in the same manner as Example 4-1.

#### Example 4-4

Ethyl 3-[3-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]ureido]propionate

25



30


Recrystallization solvent: ethyl acetate/hexane  
Melting point: 108°C - 110°C

35

## Example 4-5

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(2-dimethylaminoethyl)urea

5




Recrystallization solvent: ethyl acetate/ether

10 Melting point: 104°C - 105°C

## Example 4-6

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(3-diethylaminopropyl)urea

15




Recrystallization solvent: ethyl acetate/ether

20 Melting point: 122°C - 124°C

## Example 4-7

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-[3-(2-pyrrolidono-1-yl)propyl]urea

25




30

Recrystallization solvent: ether

Melting point: 115°C - 116°C

## Example 4-8

35 1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(2-piperidinoethyl)urea




Recrystallization solvent: ether/hexane

Melting point: 122°C - 123°C

Example 4-9

2-[3-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]ureido]ethanesulfonamide hydrochloride




Recrystallization solvent: ethyl ether/isopropyl ether

Melting point: 142°C - 145°C

Example 4-10

2-[3-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]ureido]ethanesulfonic acid




Recrystallization solvent: methanol/isopropyl ether

Melting point: 221°C - 224°C

Example 5-1

N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]succinamic acid



5

To a solution of ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl] succinamate (1.05 g) in ethanol (10 ml) was added 1N-aqueous sodium hydroxide solution (3 ml) and the mixture was stirred at 60°C for 2 hours. The reaction mixture was concentrated, diluted with water, neutralized with 1N-hydrochloric acid and extracted with ethyl acetate. The organic extract was dried and the solvent was distilled off under reduced pressure to give the titled compound (900 mg).

10

15


Melting point: 180°C - 182°C

The compounds of Examples 5-2 to 5-5 were synthesized in the same manner as Example 5-1.

20

#### Example 5-2

N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]oxamic acid




25

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.08-1.40(2H,m), 1.40-1.65(2H,m), 1.75-2.90(9H,m), 3.10-3.50(2H,m), 3.90(2H,br d), 4.80-5.40(1H,br), 7.12-7.50(14H,m).

30

#### Example 5-3


N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]malonamic acid



<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.17-1.30(2H,m), 1.46-1.63(2H,m),  
1.80-2.10(3H,m), 2.10-2.60(6H,m), 2.70-3.20(4H,m),  
3.80-3.92(2H,m), 6.64-6.90(1H,br), 7.00-7.33(14H,m).

Example 5-4


N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]glutamic acid



<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.05-1.30(2H,m), 1.40-1.66(4H,m),  
1.70-2.15(9H,m), 2.20-2.26(4H,m), 2.52-2.66(2H,m),  
3.90(2H,d), 4.30-5.70(2H,br), 7.07-7.39(12H,m),  
7.46(2H,d).

Example 5-5


N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2-phenyl-2-(2-pyridyl)pentyl]succinamic acid



<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.05-1.52(2H,m), 1.55-1.72(2H,m),  
1.90-2.50(9H,m), 2.60-3.13(6H,m), 3.83-4.20(2H,m),  
5.00-5.60(1H,br), 7.03-7.50(11H,m), 7.62-7.73(1H,m),  
8.53(1H,d).

Example 6-1

N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]glycine ethyl ester dihydrochloride



To a solution of 1-amino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane (340 mg) and potassium carbonate (414 mg) in acetonitrile (5 ml) was added ethyl bromoacetate (134 mg) and the mixture was stirred at 60°C for 2.5 hours. The reaction mixture was diluted with water and extracted with ethyl acetate. The organic extract was washed serially with water and a saturated aqueous sodium chloride solution and dried. The solvent was distilled off under reduced pressure. The obtained residue was purified by silica gel column chromatography eluting with ethyl acetate-methanol (20:1). The solvent was distilled off and residue was treated with 4N-hydrochloric acid/ethyl acetate to give the titled compound (200 mg) as a noncrystalline powder.


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.15-1.36(5H,m), 1.50-1.85(4H,m), 2.06(2H,dt), 2.16-2.42(6H,m), 2.67(2H,br d), 3.26(2H,s), 3.31(2H,s), 4.12(2H,q), 7.14-7.35(12H,m), 7.43(2H,d).

The compound of Example 6-2 was synthesized in the manner similar to Example 6-1.

#### Example 6-2

Ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-4-aminobutanoate dihydrochloride






<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.03-1.34(5H,m), 1.35-1.77(6H,m),  
2.04(2H,dt), 2.15-2.40(8H,m), 2.55(2H,t), 2.66(2H,br  
d), 3.21(2H,s), 4.08(2H,q), 7.12-7.34(12H,m),  
7.42(2H,d).

The compounds of Examples 7-1 and 7-2 were  
synthesized in a manner similar to Example 5-1.

#### Example 7-1


N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-  
diphenylpentyl]glycine



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05-1.40(2H,m), 1.45-1.73(2H,m),  
1.83-2.27(4H,m), 2.28-2.80(6H,m), 2.97(2H,s),  
3.17(2H,s), 3.50-4.50(3H,br), 6.90-7.50(14H,m).

#### Example 7-2


N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-  
diphenylpentyl]-4-aminobutyric acid



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.18-1.43(2H,m), 1.52-1.83(4H,m),  
2.05-2.34(7H,m), 2.40-2.80(6H,m), 2.81-3.04(2H,m),  
3.28(2H,s), 4.10-4.80(2H,br), 7.08-7.50(14H,m).

## Example 8-1

N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(3-hydroxypyrrolidin-1-yl)propanamide



1) 5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-(3-chloropropionylamino)-2,2-diphenylpentane

To a solution of 1-amino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane (0.9 g) in THF (20 ml) was added a saturated aqueous sodium hydrogen carbonate solution (20 ml) and the mixture was stirred vigorously under ice-cooling. 3-

Chloropropionylchloride (0.21 ml) was added and the mixture was stirred for 2 hours. The reaction mixture was diluted with ethyl acetate and organic layer was separated, washed with pure water, and dried. The solvent was distilled off under reduced pressure. The residue was purified by silica gel column chromatography eluting with ethyl acetate-methanol (7:3) to give 5-[4-(4-chlorophenyl)-4-

hydroxypiperidino]-1-(3-chloro-propionylamino)-2,2-diphenylpentane (0.82 g) as a noncrystalline powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.25-1.43(2H,m), 1.63-1.75(2H,m), 2.10-2.59(10H,m), 2.75-2.97(2H,m), 3.74(2H,t), 4.05(2H,d), 5.21(1H,br s), 7.14-7.38 (12H,m), 7.43(2H,d).

To a solution of 5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-(3-chloropropionylamino)-2,2-diphenylpentane (0.19 g) in ethanol were added potassium carbonate (0.10 g) and 3-hydroxypyrrolidine (0.045 ml) and the mixture was stirred at room temperature for 3 hours. The reaction mixture was


diluted with ethyl acetate and organic layer was separated, washed with pure water, and dried. The solvent was distilled off under reduced pressure. The residue was purified by silica gel chromatography eluting with ethyl acetate-methanol (1:1) to give the titled compound (0.08 g) as a noncrystalline powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.32-1.47(2H,m), 1.64-1.75(2H,m), 1.87-2.38 (12H,m), 2.41-2.90(8H,m), 4.02-4.18(2H,m), 4.22-4.28(1H,m), 7.18-7.36(12H,m), 7.43(2H,d), 7.93(1H,br s).

The compounds of Examples 8-2 and 8-3 were synthesized in a manner similar to Example 8-1.

#### Example 8-2


5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl-1-(3-pyrrolidin-1-yl-propionylamino)pentane



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.23-1.40(2H,m), 1.49-1.72(8H,m), 1.94-2.15(4H,m), 2.22-2.37(8H,m), 2.50(2H,t), 2.65(2H,br d), 3.83(2H,d), 4.04(2H,d), 7.10-7.36(12H,m), 7.43(2H,d), 8.22(1H,br).

#### Example 8-3

5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-1-[3-(dimethylamino)propionylamino]-2,2-diphenylpentane




<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.25-1.43(2H,m), 1.52-1.79(4H,m), 1.92(6H,s), 2.03-2.52(10H,m), 2.62-2.82(2H,br),

4.03(2H,d), 7.10-7.46(14H,m), 8.22(1H,br).

Example 9

N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(t-butoxycarbonyl)aminopropanamide




To a solution of 1-(5-amino-4,4-diphenylpentyl)-4-(4-chlorophenyl)-4-hydroxypiperidine (0.8 g) in DMF (5 ml) were added N-Boc- $\beta$ -alanine (0.3 g), triethyl amine (0.56 ml), and diethylphosphoro cyanidate (0.28 ml) at 0°C and the mixture was stirred at room temperature for 1 hour. The reaction mixture was poured into pure water (20 ml) and extracted with ethyl acetate. The extract was washed with a saturated aqueous sodium chloride solution and dried. The solvent was distilled off under reduced pressure. The residue was purified by silica gel chromatography eluting with ethyl acetate to give the titled compound (0.85 g) as a noncrystalline powder.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.20-1.39(2H,m), 1.47(9H,s), 1.61-1.75(2H,m), 1.98-2.16(4H,m), 2.18-2.38(10H,m), 2.67-2.81(2H,m), 3.30-3.42(2H,m), 4.01(2H,d), 5.08(1H,brs), 5.79(1H,br s), 7.16-7.35(12H,m), 7.42(2H,d).

Example 10

N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-aminopropanamide dihydrochloride




To a solution of N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(t-butoxycarbonyl)aminopropanamide (0.8 g) in ethyl acetate (10 ml) was added 4N-hydrochloric acid-ethyl acetate solution (2.5 ml) and stirred 60°C for 3 hours. The solvent was distilled off under reduced pressure. The residue was suspended in ethyl acetate and the solid was collected by filtration to give the titled compound (0.74 g) as a noncrystalline powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.22-1.42(2H,m), 1.60-1.73(2H,m), 1.96-2.19(6H,m), 2.23-2.39(4H,m), 2.59-2.71(2H,m), 2.83(2H,t), 4.03(2H,d), 6.52-6.62(1H,m), 7.18-7.33(12H,m), 7.42(2H,d).

#### Example 11

N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(acetylamino)propanamide




N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-aminopropanamide dihydrochloride (0.16 g) was added to the mixture of THF (3 ml) and saturated aqueous sodium hydrogen carbonate solution (3 ml). Anhydrous acetic acid (0.03 ml) was added and the mixture was stirred at room temperature for 1 hour. The reaction mixture was extracted with ethyl acetate and the organic extract was washed with a saturated

aqueous sodium chloride solution and dried. The solvent was distilled off under reduced pressure. The residue was purified by silica gel chromatography eluting with ethyl acetate. The residue was  
5 crystallized from isopropylether to give the titled compound (0.07 g).

Melting point: 128°C - 130°C

Example 12

N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-  
10 diphenylpentyl]-3-(propionylamino)propanamide




The titled compound (0.02 g) was obtained in a manner similar to Example 11.

20 Recrystallization solvent: isopropyl ether

Melting point: 128°C - 130°C

Example 13

1-[4,4-Diphenyl-5-  
(phenyloxycarbonylamino)pentanoyl]-4-(4-chlorophenyl)-  
25 4-hydroxypiperidine




To a solution of 1-(5-amino-4,4-  
diphenylpentanoyl)-4-(4-chlorophenyl)-4-  
35 hydroxypiperidine (2.32 g) obtained in Reference Example 22 in THF (50 ml) were added triethylamine

(1.39 ml) and phenyl chlorocarbonate (0.69 ml) at 0°C. The reaction mixture was stirred for 1 hour, diluted with ethyl acetate, washed with pure water, and a saturated aqueous sodium chloride solution. The organic layer was dried and the solvent was distilled off under reduced pressure. The residue was purified by silica gel chromatography eluting with ethyl acetate-hexane (3:7) to give the titled compound (2.90 g) as a noncrystalline powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.57-2.34(8H,m), 2.56(2H,t), 2.91-3.04(2H,m), 3.25-3.50(2H,m), 3.87-4.17(2H,m), 4.43-4.57(2H,m), 4.83-4.92(2H,m), 7.00(2H,d), 7.14-7.42(12H,m).

#### Example 14

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl-5-oxopentyl]-3-[3-(hydroxy)propyl]urea




In a manner similar to Example 11, the titled compound (0.32 g) was obtained from 1-[4,4-diphenyl-5-phenyloxycarbonylamino)pentanoyl]-4-(4-chlorophenyl)-4-hydroxypiperidine (0.14 g).

Recrystallization solvent: ethyl ether

Melting point: 192°C - 194°C

#### Example 15

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl-5-oxopentyl]-3-[3-(dimethylamino)ethyl]urea




The titled compound was obtained in a similar manner to Example 11.

Recrystallization solvent: ethyl ether/ether

Melting point: 223°C - 225°C

Example 16

1-(5-Acetyl-amino-4,4-diphenylpentanoyl)-4-(4-chlorophenyl)-4-hydroxypiperidine




To a solution of 1-(5-amino-4,4-diphenylpentanoyl)-4-(4-chlorophenyl)-4-hydroxypiperidine (0.46 g) in THF (10 ml) were added triethylamine (0.28 ml) and anhydrous acetic acid (0.1 ml) and the mixture was stirred at room temperature for 1 hour. The reaction mixture was diluted with ethyl acetate and washed serially with pure water and a saturated aqueous sodium chloride solution. The organic layer was dried. The solvent was distilled off under reduced pressure. The residue was purified by silica gel chromatography eluting with ethyl acetate-methanol (9:1) and crystallized from ethyl acetate-ether to give the titled compound (0.36 g).

Melting point: 191°C - 192°C

Example 17

Ethyl N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl-5-oxopentyl]succinamate






In a similar manner to Example 16, 1-(5-amino-4,4-diphenylpentanoyl)-4-(4-chlorophenyl)-4-hydroxypiperidine (0.56 g) was acylated with ethylsuccinylchloride and the desired product was crystallized from ether to give the titled compound (0.53 g).

Melting point: 94°C - 96°C

#### Example 18

N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl-5-oxopentyl]succinamic acid




To a solution of ethyl 4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl-5-oxopentyl]succinamate (0.3 g) in THF (1 ml) was added 1N-aqueous sodium hydroxide solution (1 ml) and the mixture was stirred at room temperature for 2 hours. The reaction mixture was made acidic with 1N-hydrochloric acid and extracted with ethyl acetate. The solvent was distilled off under reduced pressure. The residue was purified by silica gel chromatography eluting with ethyl acetate-methanol (9:1). The desired product was crystallized from ethyl acetate to give the titled compound (0.36 g).

Melting point: 177°C - 180°C

#### Example 19

1-[5-[4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenyl-5-oxopentyl]-3-[3-(2-oxo-1-pyrrolidino)propyl]urea




In a similar manner to Example 16, the titled compound was obtained.

Recrystallization solvent: ethyl ether

Melting point: 194°C - 197°C

#### Example 20

5-[3-(4-Chlorophenyl)-3-hydroxypyrrolidin-1-yl]-2,2-diphenyl-1-formylpentanamine




In a manner similar to Example 1-1, the titled compound was obtained from 3-(4-chlorophenyl)-3-hydroxypyrrolidine (described in Medicinal Chemistry Research 3, 459-467 (1993)).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.20-1.38(2H,m), 1.95(1H,br), 2.13-2.55(8H,m), 2.91(1H,d), 3.01-3.14(1H,m), 3.86-4.08(2H,m), 5.12(1H,br s), 7.16-7.44(14H,m), 8.10(1H,s).

#### Example 21

1-[5-[4-(4-Chlorophenyl)-3-hydroxypiperidine]-2,2-diphenylpentyl]-3-[3-hydroxy)propyl]urea



To a solution of 5-[3-(4-chlorophenyl)-3-hydroxypyrrolidin-1-yl]-2,2-diphenyl-1-formylpentanamine (0.92 g) in ethanol (5 ml) was added 4N aqueous sodium hydroxide solution (5 ml) and the mixture was stirred at 90°C for 16 hours. The reaction mixture was extracted with ethyl acetate and the extract was washed with a saturated aqueous sodium chloride solution and dried. The solvent was distilled off under reduced pressure to provide the deformylated compound (0.81 g). The obtained deformylated compound (0.65 g) was dissolved in THF (15 ml), and triethylamine (0.42 ml) was added. To the resulting mixture was added phenyl chlorocarbonate (0.21 ml) at 0°C and the mixture was stirred at the same temperature for 1 hour. The reaction mixture was extracted with ethyl acetate and the extract was washed with a saturated aqueous sodium chloride solution and dried. The solvent was distilled off under reduced pressure. The residue was purified by silica gel chromatography eluting with ethyl acetate-hexane (4:1). The solvent was distilled off under reduced pressure to provide the phenyl carbamate compound as an oily residue. In a manner similar to Example 4-1, the titled compound (0.2 g) was obtained.


Recrystallization solvent: ether/hexane

Melting point: 150°C - 153°C

Example 22

1-Formylamino-[5-[4-hydroxy-4-(4-chlorophenyl)hexamethylenimin-1-yl]-2,2-diphenylpentane

hydrochloride




In a manner similar to Example 1-1, the titled compound was obtained from 4-(4-chlorophenyl)-4-hydroxyhexamethylenimine.

10  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.17-1.40(2H,m), 1.50-2.23(9H,m), 2.25-2.97(6H,m), 4.06(2H,d), 5.20-5.35(1H,br), 7.05-7.42(14H,m), 8.08(1H,d).

Example 23:

15 5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-1-formylamino-2-phenyl-2-(2-thienyl)pentane hydrochloride




20

In a similar manner to Example 1-1, the titled compound was synthesized as a noncrystalline powder from 5-formylamino-1-iodo-4-phenyl-4-(2-thienyl)pentane described in Reference Example 31.

25  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.20-1.58(2H,m), 1.66(2H,d), 1.97-2.50(9H,m), 2.61-2.77(2H,m), 4.03(2H,dd), 5.40-5.51(1H,br), 6.83-6.99(2H,m), 7.13-7.48(10H,m), 8.08(1H,d).

Example 24:

30 2,2-Bis(4-chlorophenyl)-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-formylaminopentane hydrochloride




In a similar manner to Example 1-1, the titled compound was synthesized as a noncrystalline powder from 4,4-bis(4-chlorophenyl)-5-formylamino-

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05-1.38(2H,m), 1.60-1.80(2H,m), 1.85-2.15(5H,m), 2.23-2.40(4H,m), 2.58-2.75(2H,m), 4.00(2H,d), 5.08-5.20(1H,br), 7.00-7.20(4H,m), 7.29(6H,d), 7.42(2H,d), 8.10(1H,s).

Example 25:

Ethyl N-[2,2-bis(4-chlorophenyl)-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]]pentylsuccinamate hydrochloride



To a solution of 2,2-bis(4-chlorophenyl)-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-1-formylaminopentane (1.7g) in ethanol (30ml) was added 6N-aqueous sodium hydroxide solution (10ml) and the mixture was stirred at 100°C for 14 hours. The solvent was distilled off under reduced pressure and the residue was dissolved in ethyl acetate. The organic layer was washed with water and saturated aqueous sodium chloride solution, respectively and dried. The solvent was distilled off under reduced pressure and 4N-hydrogen chlorolide-ethyl acetate was added to give

2,2-bis(4-chlorophenyl)-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]pentylamine dihydrochloride (1.6g) as a noncrystalline powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.10-1.55(4H,m), 1.58-1.73(2H,m),  
1.94-2.16(5H,m), 2.20-2.40(4H,m), 2.59-2.72(2H,m),  
3.26(2H,s), 7.03-7.18(4H,m), 7.20-7.35(6H,m), 7.35-7.45(2H,m).


In a similar acylation in Example 3-1, the titled compound was synthesized as a noncrystalline powder from

2,2-bis(4-chlorophenyl)-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]pentylamine.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.15-1.40(5H,m), 1.59-1.74(2H,m),  
1.84-2.15(5H,m), 2.22-2.40(6H,m), 2.53-2.72(4H,m),  
3.94(2H,d), 4.09(2H,q), 5.24(1H,br t), 7.05-7.20(4H,m),  
7.20-7.34(6H,m), 7.42(2H,d).

Example 26:

N-[2,2-Bis(4-chlorophenyl)-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]]pentylsuccinamic acid



By a hydrolysis described in Example 5-1, the titled compound was synthesized as a noncrystalline powder from


Ethyl N-[2,2-bis(4-chlorophenyl)-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]]pentylsuccinamate.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.20-1.50(2H,m), 1.60-1.76(2H,m),  
1.97-2.42(9H,m), 2.75-3.20(6H,m), 3.75-4.00(2H,m),  
5.25-5.60(1H,br), 7.17(4H,d), 7.30-7.55(8H,m).

The compound 27-1 to 27-23 were synthesized in the same manner as Example 4-1.

## Example 27-1:

1-[5-[4-(4-Chlorophenyl)-4-hydroxy-  
piperidino]-2,2-diphenylpentyl]-3-[(1-ethoxycarbonyl)  
piperidin-4-yl]urea




Recrystallization solvent: ethyl ether

Melting point : 223°C to 226°C

## Example 27-2:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-3-[2-(1-pyrrolidino)ethyl]urea




Recrystallization solvent : ethyl acetate/ethyl ether

Melting point : 132°C to 133°C

## Example 27-3:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-3-[2-(diethylamino)ethyl]urea




Recrystallization solvent : ethyl acetate/ethyl ether

Melting point : 134°C to 136°C

## Example 27-4:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-3-(3-aminopropyl)-3-methylurea




Recrystallization solvent : ethyl acetate

Melting point : 92°C to 94°C

## Example 27-5:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-3-(5-hydroxypentyl)urea




Recrystallization solvent : ethyl acetate/ethyl ether

Melting point : 149°C to 151°C

## Example 27-6:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-3-[2-(dimethylamino)ethyl]-3-methyl  
urea




Noncrystalline powder



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.22-1.42(2H,m), 1.58-1.71(2H,m),  
1.99(6H,s), 2.00-2.17(4H,m), 2.22-2.38(6H,m), 2.61-  
2.75(2H,m), 2.73(3H,s), 3.12(2H,t), 3.97(2H,d),  
5.23(1H,br s), 7.17-7.33(12H,m), 7.43(2H,d).

5 Example 27-7:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-3-[2-(methylamino)ethyl]-3-methyl-  
urea




15

Noncrystalline powder

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.22-1.38(2H,m), 1.65(2H,brd), 1.95-  
2.42(8H,m), 2.20(3H,s), 2.53-2.78(4H,m), 2.72(3H,s),  
3.12-3.24(2H,m), 3.95(2H,d), 5.20(1H,brs), 6.82-  
20 6.93(1H,m), 7.15-7.34(11H,m), 7.42(2H,d).

Example 27-8:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-3-(2-hydroxyethyl)-3-methylurea



25


30

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.27-1.39(2H,m), 1.67(2H,brd), 2.02-  
2.18(2H,m), 2.25-2.43(2H,m), 2.60-2.78(2H,m),  
2.68(3H,s), 3.32(2H,t), 3.97(2H,t), 4.28(1H,brs), 6.82-  
6.94(1H,m), 7.16-7.34(11H,m), 7.42(2H,d).

35 Example 27-9:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-

2,2-diphenylpentyl]-3-[2-(acetylamino)ethyl]urea




Recrystallization solvent : ethyl acetate/ethyl ether

Melting point : 210°C to 213°C

Example 27-10:

Ethyl 4-[5-[4-(4-Chlorophenyl)-4-hydroxy-piperidino]-2,2-diphenylpentyl]ureido butyrate




Recrystallization solvent : ethyl acetate/ethyl ether

Melting point : 121°C to 123°C

Example 27-11:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(3-hydroxypropyl)urea




Recrystallization solvent : ethyl acetate/ethyl ether

Melting point : 101°C to 102°C

Example 27-12:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(1-benzylpiperidin-4-yl)urea




Recrystallization solvent : isopropyl ether/ethyl ether

Melting point : 176°C to 178°C

Example 27-13:

N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-4-methylpiperidine-1-carboxamide




Recrystallization solvent : isopropyl ether/ethyl ether

Melting point : 156°C to 157°C

Example 27-14:

N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-4-benzylpiperidine-1-carboxamide




Recrystallization solvent : isopropyl ether/ethyl ether

Melting point : 142°C to 143°C

Example 27-15:

1N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-1,2,4,5-tetrahydro-3-benzazepine-3-carboxamide




Noncrystalline powder

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.23-1.44(2H,m), 1.57-1.80(2H,m), 1.98-2.20(4H,m), 2.23-2.47(4H,m), 2.62-2.82(6H,m), 3.32-3.37(4H,m), 3.97-4.06(3H,m), 7.02-7.19(4H,m), 7.21-7.43(14H,m).

Example 27-16:

1N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(trifluoroacetylaminopyrrolidine-1-carboxamide




Noncrystalline powder

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.30(2H,br), 1.61-1.75(2H,m), 1.96-2.42(11H,m), 2.70(2H,br), 3.12-3.24(2H,m), 3.27-3.48(2H,m), 3.72-3.80(1H,m), 3.83-4.07(2H,m), 4.42(1H,br), 7.17-7.42(14H,m).

Example 27-17:

1N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-4-(t-butoxycarboxamido)piperidine-1-carboxamide




Noncrystalline powder

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.15-1.37(2H,m), 1.43(9H,s), 1.60-1.91(6H,m), 1.97-2.16(4H,m), 2.22-2.39(4H,m), 2.57-2.79(4H,m), 3.43-3.65(3H,m), 3.95(3H,br), 4.42(1H,br), 7.15-7.37(12H,m), 7.40-7.45(2H,m).

Example 27-18:

Ethyl [4-[3-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]ureido]piperidino] acetate




Recrystallization solvent : isopropyl ether/ethyl ether

Melting point : 116°C to 118°C

Example 27-19:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-[1-(trifluoroacetyl)piperidin-4-yl]urea




Recrystallization solvent : ethyl ether

Melting point : 192°C to 193°C

Example 27-20:

1N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-4-formyl-1-piperazinecarboxamide




Recrystallization solvent : ethyl acetate/ethyl ether

Melting point : 191°C to 192°C

Example 27-21:

1N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-4-(3-hydroxypropyl)-1-piperazine-  
carboxamide




Recrystallization solvent : ethyl acetate/ethyl ether

Melting point : 125°C to 127°C

Example 27-22:

1N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-4-(ethoxycarbonyl)-1-piperazine-  
carboxamide




Noncrystalline powder

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.20-1.37(2H,m), 1.25(3H,t), 1.61-1.68(2H,m), 1.97-2.17(4H,m), 2.21-2.38(4H,m), 2.60-2.72(2H,m), 3.03-3.20(4H,m), 3.34-3.41(4H,m), 3.92-4.00(1H,br), 3.96(2H,s), 4.12(2H,q), 7.17-7.44(14H).

Example 27-23:

1N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-4-(morpholinocarbonylmethyl)-1-piperadinecarboxamide




Recrystallization solvent : ethyl acetate/ethyl ether

Melting point : 163°C to 165°C

The compound 28-1 and 28-2 were synthesized in the same manner as Example 5-1.

Example 28-1:

3-[3-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]ureido]propionic acid




Noncrystalline powder

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.46(2H,br), 1.71(2H,brd), 2.17-2.60(6H,m), 2.72-3.03(4H,m), 3.15-3.45(4H,m), 3.92(2H,brd), 4.87(1H,br), 5.71(1H,br), 7.12-7.45(14H,m).

Example 28-2:

4-[3-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]ureido]butyric acid




10 Recrystallization solvent : water

Melting point : 137°C to 139°C

Example 29:

1N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]vinylsulfonamide



20

To a solution of 1-amino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane (0.9g) in THF (20ml) were added triethylamine (0.84ml) and 2-chloroethanesulfonylchloride (0.21ml) at room-  
25 temperature. The reaction mixture was stirred for 2 hours, diluted with ethyl acetate, washed with pure water, and dried. The solvent was distilled off under reduced pressure. The residue was purified by silica gel chromatography eluting with ethyl acetate-methanol  
30 (7:3) to give the titled compound as noncrystalline powder (0.82g).


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.21-1.35(2H,m), 1.60-1.73(2H,m), 2.07-2.45(8H,m), 2.65-2.79(2H,m), 3.70(2H,br s), 5.55(1H,dd), 6.11(1H,d), 6.12(1H,d), 7.14-7.48(14H,m).

35 Example 30:

1N-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-



2,2-diphenylpentyl]-2-(pyrrolidino)ethylsulfonamide




To a solution of 1N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]vinylsulfonamide in ethanol was added pyrrolidine (0.062ml) and the mixture was stirred at room temperature for 16 hours. The reaction mixture was diluted with ethyl acetate, washed with saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure. The residue was purified by silica gel chromatography eluting with ethyl acetate-methanol (7:3) to give the titled compound (0.10g).

Recrystallization solvent : ethyl acetate/ethyl ether

Melting point : 140°C to 142°C

Example 31:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-[3-carbamoyloxy)propyl]urea



To a solution of 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(3-hydroxypropyl)urea described in Example 27-11 in THF (5ml) was added chlorosulfonylisocyanate (0.044ml) and the mixture was stirred at room temperature for 2 hours, followed by saturated aqueous sodium hydrogen carbonate (5ml) was added. After stirring for 1 hour, at 45°C.


The reaction mixture was diluted with ethyl acetate, washed with saturated aqueous sodium chloride, and dried. The solvent was distilled off under reduced pressure. The residue was purified by silica gel chromatography eluting with ethyl acetate-methanol (4:1) to give the titled compound (0.10g).

Recrystallization solvent : ethyl ether

Melting point : 152°C to 154°C

Example 32:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(piperidin-4-yl)urea




To a solution of 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-[1-(trifluoroacetyl)piperidin-4-yl]urea (2.35g) described in Example 27-19 in THF (10.5ml) was added 0.5N-sodium hydroxide solution (10.5ml) and the mixture was stirred for 2 hours at room temperature. The reaction mixture was diluted with ethyl acetate, washed with saturated aqueous sodium chloride, and dried. The solvent was distilled off under reduced pressure. The residue was crystallized from ethyl acetate-ether (4:1) to give the titled compound (1.92 g).

Melting point : 194°C to 196°C

Example 33-1:

Ethyl 4-[4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino]piperidino-4-oxobutyrate



To a mixture of 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(piperidin-4-yl)urea described in Example 32 in THF (10ml) and triethylamine (0.21ml) was added ethyl succinylchloride (0.077ml). After stirring for 2 hours at room temperature, the reaction mixture was diluted with ethyl acetate, washed with saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure. The residue was purified by silica gel chromatography eluting with ethyl acetate-methanol (4:1) to give the titled compound (0.28g).


Recrystallization solvent : ethyl acetate/ethyl ether

Melting point : 173°C to 175°C

The compound 33-2 to 33-5 were synthesized in the same manner as Example 33-1.

Example 33-2:

N-Ethyl-4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino-1-piperidinecarboxamide




Recrystallization solvent : ethyl acetate/ethyl ether

Melting point : 193°C to 195°C

Example 33-3:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(1-acetylpiperidin-4-yl)urea




10 Recrystallization solvent : ethyl ether

Melting point : 146°C to 148°C

Example 33-4:

N-Ethoxycarbonylmethyl-4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonyl-amino-1-piperidinecarboxamide




Recrystallization solvent : ethyl ether

Melting point : 220°C to 221°C

25 Example 33-5:

Ethyl 3-[4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino]piperidino-3-oxopropionate




Recrystallization solvent : ethyl ether

Melting point : 173°C to 175°C

Example 34-1:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(1-ethylpiperidin-4-yl)urea




To a mixture of 1-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(piperidin-4-yl)urea (0.29g) described in Example 32 in DMF (5ml) and potassium carbonate (0.14g) was added ethyl iodide (0.12ml). After stirring for 8 hours at room temperature, the reaction mixture was diluted with ethyl acetate, washed with saturated aqueous sodium chloride solution, and dried. The solvent was distilled off under reduced pressure. The residue was crystallized from ethyl acetate to give the titled compound (0.14g).

Melting point : 154°C to 157°C

The compound 34-2 to 34-4 were synthesized in the same manner as Example 34-1.

Example 34-2:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-[1-(2-hydroxyethyl)piperidin-4-yl]urea




Recrystallization solvent : ethyl ether

Melting point : 177°C to 180°C

Example 34-3:

Ethyl 3-[4-[5-[4-(4-chlorophenyl)-4-hydroxy-  
piperidino]-2,2-diphenylpentyl]aminocarbonylamino  
piperidino]propionate




Recrystallization solvent : ethyl ether/ethyl ether

Melting point : 148°C to 151°C

Example 34-4:

1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-  
2,2-diphenylpentyl]-3-[1-(3-hydroxypropyl)piperidin-4-yl]  
urea




Recrystallization solvent : ethyl ether/ethyl ether

Melting point : 194°C to 197°C

Example 35:

1-[(Piperidin-4-yl)carboxamido]-5-[4-(4-  
chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane  
dihydrochloride




To a mixture of 1-trifluoroacetylpiperidin-4-carboxylic acid (1.2 g) in acetonitrile (30 ml) and triethylamine (0.6 g) was added chloroisopropylcarbonate (0.67 g) slowly and the mixture was stirred for 2 minutes at  $-15^{\circ}\text{C}$ . A solution of 1-amino-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane (2.5 g) and triethylamine (0.5 g) in THF (50 ml) was added to the mixture and the mixture was stirred at room temperature for 24 hours. The reaction mixture was diluted with ethyl acetate, washed with saturated aqueous sodium hydrogen carbonate and water, and dried. The solvent was distilled off under reduced pressure. The residue was dissolved in ethanol-water (2:1), followed by sodium hydroxide (2 g) was added to the mixture. The mixture was stirred at room temperature for 18 hours. The reaction mixture was distilled off under reduced pressure. Water was added to the residue and the mixture was extracted with ethyl acetate, washed, and dried. The solvent was distilled off under reduced pressure and the residue was purified by alumina column eluting with ethyl acetate-ethanol (4:1). The eluted solution was distilled off and the residue was dissolved in ethyl acetate. To the solution was added an excess amount of 4N-hydrochloric acid/ethyl acetate and the solvent was distilled off under reduced pressure to give the titled compound.

Noncrystalline powder

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.20-1.70(9H,m), 1.87-2.18(6H,m), 2.20-2.39(4H,m), 2.42-2.70(4H,m), 3.05(2H,dt), 3.98(2H,d), 5.02(1H,t), 7.14-7.48(14H,m).

## Example 36-1:

1-[(N-Ethylpiperidin-4-yl)carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane dihydrochloride



To a solution of 1-[(Piperidin-4-yl)carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane (0.4g) in acetonitrile (10ml) were added potassium carbonate (0.5g) and ethyl iodide (0.2ml) and the mixture was stirred at room temperature for 18 hours. The reaction mixture was dissolved in ethyl acetate, washed with water, and dried. The solvent was distilled off under reduced pressure. The residue was purified by aluminum oxide column eluting with ethyl acetate. The eluted solution was distilled off and the residue was dissolved in ethyl acetate. An excess amount of 4N-hydrochloric acid/ethyl acetate was added to the solution and the solvent distilled off under reduced pressure to give the titled compound. Noncrystalline powder


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05(3H,t), 1.20-1.85(14H,m), 1.80-2.35(8H,m), 2.65(2H,m), 2.90(2H,m), 3.99(2H,d), 5.04(1H,t), 7.14-7.48(14H,m).

The compound 36-2 to 36-4 were synthesized in the same manner as Example 36-1.

## Example 36-2:

1-[N-(Ethoxycarbonylmethyl)piperidin-4-yl]carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane dihydrochloride






Noncrystalline powder

10  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.26(3H,t), 1.20-1.85(10H,m), 1.80-2.40(8H,m), 2.65(2H,m), 2.90(2H,m), 3.16(2H,s), 3.98(2H,d), 4.16(2H,q), 5.04(1H,t), 7.14-7.48(14H,m).

Example 36-3:

15 1-[[N-(2-Morpholinoethyl)piperidin-4-yl]carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane trihydrochloride




Noncrystalline powder

25  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.20-1.85(12H,m), 1.80-2.50(12H,m), 2.46(4H,s), 2.65(2H,m), 2.90(2H,m), 3.70(4H,m), 3.98(2H,d), 5.01(1H,t), 7.14-7.48(14H,m).

Example 36-4:

30 1-[[N-(2-Dimethylaminoethyl)piperidin-4-yl]carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane trihydrochloride




Noncrystalline powder

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.20-1.85(12H,m), 2.12(6H,s), 2.40(4H,s), 1.80-2.50(12H,m), 2.65(2H,m), 2.89(2H,m), 3.98(2H,d), 5.02(1H,t), 7.14-7.48(14H,m).

The compound 37-1 to 37-8 were synthesized in the same manner as Example 33-1.

Example 37-1:

1-[[[2,2-diphenylpentan-1-yl]carbamoyl]piperidin-4-yl]-4-(4-chlorophenyl)-4-hydroxypiperidine-5-carboxamide hydrochloride




Noncrystalline powder

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.12(3H,t), 1.10-2.10(14H,m), 2.20-2.40(4H,m), 2.45-2.95(4H,m), 3.20(2H,m), 3.85-3.91(2H,m), 4.00(2H,d), 4.38(1H,t), 5.02(1H,t), 7.14-7.48(14H,m).

Example 37-2:

1-[[[2,2-diphenylpentan-1-yl]carbamoyl]piperidin-4-yl]-4-(4-chlorophenyl)-4-hydroxypiperidine-5-carboxamide hydrochloride




5

Noncrystalline powder

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.10-1.80(8H,m), 1.90-2.15(4H,m),  
 10 2.20-2.40(4H,m), 2.60-2.85(4H,m), 3.67(3H,s), 3.95-  
 4.20(2H,d), 4.00(2H,d), 4.38(1H,t), 5.02(1H,t), 7.14-  
 7.48(14H,m).

Example 37-3:

1-[[N-Phenylcarbamoyl]piperidin-4-yl]carbo-  
 15 xamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-  
 diphenylpentane hydrochloride




20

Noncrystalline powder

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.10-2.10(14H,m), 2.20-2.40(4H,m),  
 25 2.50-2.90(4H,m), 4.00(2H,d), 4.00-4.20(2H,m),  
 5.04(1H,t), 6.43(1H,m), 7.14-7.48(19H,m).

Example 37-4:

1-[[N-(4-Chlorobenzoyl)piperidin-4-yl]carbo-  
 30 amido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-  
 diphenylpentane hydrochloride




10 Noncrystalline powder

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.10-2.40(14H,m), 2.50-2.90(5H,m), 3.75(1H,m), 3.98(2H,d), 3.90-4.20(2H,m), 4.50(1H,m), 5.04(1H,t), 7.14-7.48(17H,m), 7.93(2H,d).

Example 37-5:

15 1-[[N-(Ethoxycarbonylacetyl)piperidin-4-yl]carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane hydrochloride




25 Noncrystalline powder

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.25(3H,t), 1.10-2.30(15H,m), 2.42-2.80(4H,m), 3.00(1H,t), 3.43(2H,s), 3.60(1H,m), 3.98(2H,d), 3.80-4.00(2H,m), 4.19(2H,q), 4.43(1H,m), 5.04(1H,t), 7.10-7.58(14H,m).

30 Example 37-6:

1-[[N-(3-Methoxycarbonylpropionyl)piperidin-4-yl]carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane hydrochloride

5




Monocrystalline powder

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.10-1.80(10H,m), 1.90-2.50(8H,m),  
 10 2.59(4H,m), 2.60-2.80(2H,m), 2.96(1H,t), 3.68(3H,s),  
 3.80-4.00(4H,m), 4.40(1H,d), 5.18(1H,m), 7.00-  
 7.50(14H,m).

Example 37-7:

1-[[N-(Nicotinoyl)piperidin-4-yl]carboxamido]-5-  
 15 [4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-  
 diphenylpentane dihydrochloride

20



25


Noncrystalline powder

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.10-1.80(10H,m), 1.90-2.50(8H,m),  
 2.60(2H,m), 2.80-3.10(2H,m), 3.70(1H,m), 4.02(2H,d),  
 4.50(1H,m), 5.04(1H,m), 7.00-7.50(15H,m), 7.73(1H,dt),  
 8.61(1H,d), 8.66(1H,dd).

30

Example 37-8:

1-[[N-(4-Dimethylaminobutyl) piperidin-4-  
 yl]carboxamido]-5-[4-(4-chlorophenyl)-4-  
 hydroxypiperidino]-2,2-diphenylpentane dihydrochloride




Noncrystalline powder

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.10-1.80(10H,m), 2.16(6H,s), 1.90-2.50(15H,m), 2.60(2H,m), 2.80-3.10(1H,m), 3.70-3.90(1H,m), 4.01(2H,m), 4.50(1H,d), 5.04(1H,t), 7.00-7.50(14H,m).

Example 38:

1-[(N-Propylpiperidin-4-yl)carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane dihydrochloride



Noncrystalline powder


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 0.86(3H,t), 1.20-1.85(16H,m), 1.80-2.35(8H,m), 2.64(2H,m), 2.87(2H,m), 3.98(2H,d), 5.05(1H,t), 7.14-7.48(14H,m).

The compound 39 and 40 was synthesized in the same manner as Example 37-1.

Example 39:

1-[[N-(3-Pyridylacetyl)piperidin-4-yl]carboxamido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentane dihydrochloride

5




Noncrystalline powder

10  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.10-1.80(8H,m), 1.90-2.40(8H,m),  
2.60(2H,m), 2.80(3H,m), 3.04(1H,m), 3.68(2H,s), 3.80-  
4.00(2H,m), 4.00(2H,d), 4.48(1H,m), 5.04(1H,m), 7.00-  
7.50(15H,m), 7.65(1H,d), 8.54(2H,d).

Example 40

15 1-[[ (N-Ethylcarbamoyl)piperidin-4-  
yl]carboxamide]5-[4-(4-chlorophenyl)-4-  
hydroxypiperidino]-2,2-diphenylpentane hydrochloride

20



Noncrystalline powder

25  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.27(3H,t), 1.10-1.80(8H,m), 1.90-  
2.15(4H,m), 2.20-2.40(4H,m), 2.60-2.85(4H,m),  
3.05(1H,m), 3.75(1H,m), 3.95-4.20(2H,d), 4.22(2H,q),  
4.47(1H,t), 5.07(1H,t), 7.14-7.48(14H,m).

Formulation Example 1

30	(1) Compound of Example 4-2	10.0 g
	(2) Lactose	60.0 g
	(3) Corn starch	35.0 g
	(4) Gelatin	3.0 g
	(5) Magnesium stearate	2.0 g

35 Using 30 ml of an 10 weight% aqueous solution of  
gelatin (3.0 g as gelatin), a mixture of 10.0 g of the

compound obtained in Example 4-2, 60.0 g of lactose and 35.0 g of corn starch was granulated by means of a 1 mm-mesh sieve, dried at 40°C, and re-sieved. The granules thus prepared were mixed with 2.0 g of magnesium stearate and the mixture was compressed. The core tablets thus obtained were coated using an aqueous suspension containing sucrose, titanium dioxide, talc and gum arabic. The coated tablet were then glazed with beenwax to provide 1000 finished tablets.

10 Formulation Example 2

(1)	Compound of Example 4-2	10.0 g
(2)	Lactose	70.0 g
(3)	Corn starch	50.0 g
(4)	Soluble starch	7.0 g
15 (5)	Magnesium stearate	2.0 g

Using 70 ml of an aqueous solution of soluble starch (7.0 g as soluble starch), a mixture of 10.0 g of Compound obtained in Example 4-2 and 3.0 g of magnesium stearate was granulated, dried, and mixed with 70.0 g of lactose and 50.0 g of corn starch. The whole mixture was then compressed to provide 1000 tablets.

Test Example 1

Determination of inhibitory activity of <sup>125</sup>I-RANTES binding using human MIP-1α/RANTES receptor-expressing CHO cells (CHO (CCR) cells)

CHO (CCR) cells were inoculated on 96 well microplates (CulturPlate, manufactured by Packard Instrument Company, Meriden, CT. U.S.A.) in an amount of  $5 \times 10^4$ /100 μl/well and then cultured for 24 hours. After removing the medium, 35 μl/well of DMEM/0.5% BSA, 5 μl/well of a test compound diluted with DMEM/0.5% BSA and 10 μl/well of <sup>125</sup>I-RANTES (final concentration of 200 pM) were added in order, followed by incubation at room temperature for 40 minutes. Then, the cells were washed twice with 200 μl/well of PBS and 25 μl/well of



ethanol was added and the mixture was stirred. Furthermore, 200  $\mu$ l/well of a scintillator (MicroScint-20, by Packard Instrument Company) was added and stirred, and then the radioactivity of  $^{125}$ I-RANTES bound to the cells was measured using a TopCount (Packard Instrument Company). Assuming that the amount of binding in case that no test compound is added is 100 % and the amount bound to the CHO cells to which vector plasmid pAKKO-111H has been transfected is 0%, the concentration at which 50% inhibition of binding of  $^{125}$ I-RANTES arises ( $IC_{50}$  value) was determined.

15	Compound No.	Binding inhibitory activity toward human RANTES receptor $IC_{50}$ ( $\mu$ M)	Binding inhibitory activity toward human MIP-1 $\alpha$ receptor $IC_{50}$ ( $\mu$ M)
20	Example 1-1	0.04	0.2
	Example 1-9	0.2	
	Example 3-4	0.01	
	Example 3-5	0.04	
	Example 4-2	0.02	0.05
25	Example 4-3	0.01	
	Example 4-5	0.02	
	Example 4-7	0.04	
	Example 4-8	0.05	
	Example 32	0.006	5
30	Example 33-2	0.02	0.6
	Example 33-5	0.01	3
	Example 34-1	<0.01	
	Example 35	0.03	5
	Example 37-1	0.03	0.1
35	Example 37-5	0.03	0.1
	Example 37-6	0.05	0.09
	Ioperamide	3	

## Test Example 2

Determination of inhibition activity of compound by migratrion assay using CHO (CCR) cells

Using a 96 well microchemotaxis chamber  
5 (NeuroProbe, Inc., Cabin John, MD, U.S.A), migration assay was conducted. A pretreatment was conducted by dipping a polycarbonate frame filter (NeuroProbe) having a pore size of 5  $\mu$ m in a bovine fibronectin solution (10  $\mu$ g/ml) diluted with PBS at room  
10 temperature for 10 minutes, followed by air-drying. A solution prepared by dissolving 40 nM RANTES (37  $\mu$ l) in DMEM/0.5% BSA was added to the lower chamber. A solution (100  $\mu$ l) prepared by diluting the test compound with DMEM/0.5% BSA was firstly added to the  
15 upper chamber and then CHO (CCR) cells ( $2 \times 10^6$  cells/ml, 100  $\mu$ l) were added. After incubating at 37°C for 4 hours, the absorbance at 595 nm the CHO cells which migrated to the bottom surface of the filter was fixed and stained with Diff-Quick, and was measured.  
20 Assuming that the absorbance in case that 40 nM RANTES is added to the lower chamber and no test compound is added to the upper chamber is 100 % and the absorbance in case that only DMEM/0.5% BSA is added to the lower chamber and no test compound is added to the upper  
25 chamber is 0%, the concentration at which 50% inhibition of wandering of the CHO (CCR) cells arises ( $IC_{50}$  value) was determined.


The respective test compounds inhibited migration of the CHO (CCR) cells in the  $IC_{50}$  value of less than  
30 10  $\mu$ M.

INDUSTRIAL APPLICABILITY

The present invention provides an excellent MIP-1 $\alpha$ /RANTES-receptor antagonist useful as prophylactic and therapeutic agent for allergic and inflammatory  
35 diseases, etc., which comprises a diphenylmethane derivative or pharmaceutically acceptable salt thereof.


Japanese Patent Application No.343905/1995 filed  
December 28, 1995 and Japanese Patent Application No.  
187375/1996 filed July 17, 1996, which are the priority  
documents of the present application, are hereby  
5 incorporated by reference in their entirety.

Table 1



Reference Example No.	X	Y	n	<sup>1</sup> H-NMR ( $\delta_{ppm}$ , CDCl <sub>3</sub> )
1-1	H	H	2	3.24 (2H, s), 7.19-7.50 (10H, m), 9.79(1H, s)
1-2	H	H	3	2.04-2.13 (2H, m), 2.64-2.74(2H, m), 7.11-7.45 (10H, m), 9.79(1H, s)


Table 2



Reference	X	Y	n	<sup>1</sup> H-NMR (δ <sub>ppm</sub> , CDCl <sub>3</sub> )
2-1	H	H	2	1.29 (3H, t), 3.28 (2H, s), 4.21 (2H, q), 5.71, 7.52 (1H each, d), 7.14-7.41 (10H, m)
2-2	H	H	3	1.29 (3H, t), 2.09-2.18 (2H, m), 2.65-2.77 (2H, m), 4.20 (2H, q), 5.63, 7.47 (1H each, d), 7.09-7.19 (4H, m), 7.22-7.40 (6H, m)


Example No.

Table 3




Reference Example No.	X	Y	<sup>1</sup> H-NMR (δ, ppm, CDCl <sub>3</sub> )
3-1	4-Cl	H	5.11 (1H, s), 7.23-7.42 (9H, m)
3-2	4-MeO	H	3.80 (3H, s), 5.10 (1H, s), 6.85-6.94 (2H, m), 7.20-7.40 (7H, m)
3-3	4-Cl	4-Cl	5.10 (1H, s), 7.20-7.40 (8H, m)

Table 4



Reference Example No.	X	Y	n	<sup>1</sup> H-NMR (δ, ppm, CDCl <sub>3</sub> )	
4-1	H	H	1	3	1.23 (3H, t), 2.40-2.51 (2H, m), 2.71-2.82 (2H, m), 4.11 (2H, q), 7.26-7.43 (10H, m)
4-2	4-Cl	H	1	3	1.23 (3H, t), 2.38-2.48 (2H, m), 2.78-2.88 (2H, m), 4.10 (2H, q), 7.29-7.40 (9H, m)
4-3	4-MeO	H	1	3	1.60-1.85 (2H, m), 2.23-2.50 (2H, m), 3.69 (2H, t), 3.79 (3H, s), 5.88-6.90 (2H, m), 7.10-7.40 (7H, m)
4-4	4-Cl	4-Cl	1	3	1.23 (3H, t), 2.41 (2H, m), 2.70 (2H, m), 4.10 (2H, q), 7.20-7.40 (8H, m)
4-5	H	H	1	4	1.24 (3H, t), 1.69-1.86 (2H, m), 2.38 (2H, t), 2.38-2.48 (2H, m), 4.12 (2H, q), 7.23-7.43 (10H, m)
4-6	H	H	2	3	1.21 (3H, t), 2.09, 2.66 (2H each, t), 3.09 (2H, s), 4.06 (2H, q), 7.16-7.39 (10H, m)
4-7	H	H	3	3	1.22 (3H, t), 1.95-2.06 (4H, m), 2.36-2.48 (4H, m), 4.07 (2H, q), 7.09-7.37 (10H, m)


Table 5



Reference Example No.	X	Y	n	m	m.p.(°C)	<sup>1</sup> H-NMR (δ ppm, CDCl <sub>3</sub> )
5-1	U	H	1	3	Syrup	
5-2	4-Cl	H	1	3	Syrup	1.17-1.33 (2H, m), 1.55 (2H, br s), 2.14-2.44 (2H, m), 3.31 (2H, s), 3.56 (2H, t), 7.07-7.38 (9H, m)
5-3	4-MeO	H	1	3	Syrup	1.20-1.35 (2H, m), 2.15-2.25 (2H, m), 3.31 (2H, s), 3.57 (2H, t), 3.79 (3H, s), 6.78-6.85 (2H, m), 7.05-7.35 (7H, m)
5-4	4-Cl	4-Cl	1	3	Syrup	1.10-1.30 (2H, m), 1.55 (2H, br s), 2.14-2.24 (2H, m), 3.29 (2H, s), 3.55 (2H, t), 7.00-7.30 (8H, m)
5-5	H	H	1	4	Syrup	1.01-1.18 (2H, m), 1.42-1.65 (4H, m), 2.09-2.20 (2H, m), 3.33 (2H, s), 3.56 (2H, t), 7.12-7.35 (10H, m)
5-6	H	H	2	3	Syrup	1.14-1.32 (2H, m), 2.10-2.26 (2H, m), 2.24-2.39 (2H, m), 2.37-2.51 (2H, m), 3.15 (3H, s), 3.51 (2H, t), 7.07-7.30 (10H, m)
5-7	H	H	3	3	Syrup	1.10-1.31 (4H, m), 2.05-2.22 (4H, m), 2.66, 3.53 (2H each, t), 3.01 (3H, br s), 7.06-7.30 (10H, m)




Table 6



Reference Example No.	X	Y	R	n	m	p.	<sup>1</sup> H-NMR (δ, ppm, CDCl <sub>3</sub> )
6-1	H	H	CHO	1	3	151-152	1.32 (2H, m), 2.16 (2H, m), 3.55 (2H, t), 4.05 (2H, d), 5.10-5.30 (1H, m), 7.10-7.40 (10H, m), 8.08 (1H, d)
6-2	4-Cl	H	CHO	1	3	159-161	1.00-1.23 (2H, m), 2.03 (2H, t), 3.30 (2H, q), 3.88 (2H, dd), 4.33 (1H, t), 7.10-7.37 (9H, m), 7.48 (1H, br t), 7.87 (1H, d)
6-3	4-MeO	H	CHO	1	3	Syrup	1.20-1.40 (2H, m), 2.08-2.20 (2H, m), 3.54 (2H, br t), 3.79 (3H, s), 4.01 (2H, dt), 5.20-5.30 (1H, br s), 6.80-6.88 (2H, m), 7.00-7.35 (7H, m), 8.07 (1H, d)
6-4	4-Cl	4-Cl	CHO	1	3	175-178	1.00-1.20 (2H, m), 2.04 (2H, t), 3.30 (2H, q), 3.86 (2H, d), 4.34 (1H, t), 7.10-7.40 (8H, m), 7.55 (1H, br t), 7.88 (1H, d)
6-5	H	H	CHO	1	4	Syrup	1.04-1.22 (2H, m), 1.40-1.56 (2H, m), 1.90-2.18 (2H, m), 3.54 (2H, t), 4.06 (2H, d), 5.20 (1H, br t), 7.10-7.37 (10H, m), 8.08 (1H, d)
6-6	H	H	CHO	2	3	Syrup	1.20-1.38 (2H, m), 2.20-2.40 (4H, m), 3.06 (2H, q), 3.57 (2H, t), 5.49 (1H, br), 7.10-7.34 (10H, m), 7.99 (1H, d)
6-7	H	H	Ac	3	3	Syrup	1.12-1.30 (4H, m), 1.90 (3H, s), 2.02-2.21 (5H, m), 3.15 (2H, q), 3.55 (2H, t), 5.49 (1H, br t), 7.11-7.30 (10H, m)

Table 7




Reference Example No.	X	Y	R	L	n	m	p.	<sup>1</sup> H-NMR (δ, ... CDCl <sub>3</sub> )
							(°C)	
7-1	H	H	CHO	I	1	3	Syrup	1.49-1.65 (2H, m), 2.12-2.25 (2H, m), 3.10 (2H, t), 4.04 (2H, d), 5.07 (1H, br t), 7.11-7.40 (10H, m), 8.11 (1H, d)
7-2	H	H	Ts	I	1	3	Syrup	1.31-1.49 (2H, m), 2.15-2.26 (2H, m), 2.44 (3H, s), 2.99 (2H, t), 3.54 (2H, d), 3.87 (1H, t), 7.05, 7.06, 7.64 (2H each, d), 7.20-7.35 (8H, m)
7-3	4-Cl	H	CHO	I	1	3	Syrup	1.46-1.63 (2H, m), 2.10-2.22 (2H, m), 3.10 (2H, t), 4.01 (2H, d), 5.08 (1H, br t), 7.06-7.39 (9H, m), 8.10 (1H, d)
7-4	4-veO	H	CHO	I	1	3	Syrup	1.45-1.65 (2H, m), 2.09-2.22 (2H, m), 3.09 (2H, t), 3.80 (3H, s), 3.99 (2H, d), 5.00-5.15 (1H, br s), 6.80-6.90 (2H, m), 7.00-7.38 (7H, m), 8.10 (1H, d)
7-5	4-Cl	4-Cl	CHO	OTs	1	3	Syrup	1.30-1.60 (2H, m), 2.00-2.20 (2H, m), 2.45 (3H, s), 3.90-4.00 (4H, m), 5.00-5.20 (1H, br s), 7.00-7.40 (10H, m), 7.72 (2H, m), 8.08 (1H, d)
7-6	H	H	CHO	I	1	4	Syrup	1.07-1.28 (2H, m), 1.65-1.85 (2H, m), 2.00-2.13 (2H, m), 3.07 (2H, t), 4.04 (2H, d), 5.02 (1H, br s), 7.10-7.39 (10H, m), 8.10 (1H, d)
7-7	H	H	CHO	I	2	3	Syrup	1.43-1.61 (2H, m), 2.19-2.40 (4H, m), 3.08 (2H, q), 3.11 (2H, t), 5.19 (1H, br s), 7.12-7.36 (10H, m), 7.99 (1H, d)
7-8	H	H	Ts	I	2	3	Syrup	1.32-1.49 (2H, m), 2.02-2.17 (2H, m), 2.21-2.34 (2H, m), 2.42 (3H, s), 2.62-2.78 (2H, m), 3.05 (2H, t), 4.26 (1H, t), 7.06, 7.07, 7.61 (2H each, d), 7.12-7.32 (8H, m)
7-9	H	H	Ac	I	3	3	Syrup	1.12-1.28 (2H, m), 1.40-1.56 (2H, m), 1.91 (3H, s), 2.04-2.23 (4H, m), 3.10 (2H, t), 3.17 (2H, q), 5.25 (1H, br s), 7.11-7.32 (10H, m)

## CLAIMS

What is claimed is

1. A composition for antagonizing MIP-1 $\alpha$ /RANTES receptor comprising a compound of the formula:




wherein Ar<sup>1</sup> and Ar<sup>2</sup> independently represent an optionally substituted aromatic group;

Q<sup>1</sup> and Q<sup>2</sup> independently represent an optionally substituted divalent C<sub>1-6</sub> aliphatic hydrocarbon group which may have oxygen or sulfur within the carbon chain;

R<sup>1</sup> is a hydrogen atom, an optionally substituted lower alkyl group or an optionally substituted lower alkyl-carbonyl group;

R<sup>2</sup> is an optionally substituted hydrocarbon group or an acyl group, or R<sup>1</sup> and R<sup>2</sup>, taken together with the adjacent nitrogen atom, may form an optionally substituted nitrogen-containing heterocyclic ring; and a group of the formula:



is an optionally substituted monocyclic or fused nitrogen-containing heterocyclic group, or a salt thereof.

2. A composition as claimed in claim 1, wherein

Ar<sup>1</sup> and Ar<sup>2</sup> independently represent (A) a monocyclic or fused polycyclic aromatic hydrocarbon group having 6 to 14 carbon atoms, or (B) a 5- to 11-membered monocyclic or fused heteroaromatic group having at least one of 1 or 2 kinds of hetero atoms

selected from nitrogen, sulfur and oxygen in addition to carbon atoms, said heterocyclic group being optionally fused with the monocyclic or fused polycyclic aromatic hydrocarbon group having 6 to 14 carbon atoms, each of which may have a substituent selected from the group consisting of

- (1) a halogen atom,
- (2) a C<sub>1-3</sub> alkylenedioxy group,
- (3) a nitro group,
- (4) a cyano group,
- (5) a C<sub>1-6</sub> alkyl group optionally having 1 to 3 halogen atoms,
- (6) a C<sub>2-6</sub> alkenyl group optionally having 1 to 3 halogen atoms,
- (7) a C<sub>2-6</sub> alkynyl group optionally having 1 to 3 halogen atoms,
- (8) a C<sub>3-6</sub> cycloalkyl group,
- (9) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms,
- (10) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms,
- (11) a hydroxyl group,
- (12) an amino group,
- (13) a mono-C<sub>1-6</sub> alkylamino group,
- (14) a di-C<sub>1-6</sub> alkylamino group,
- (15) a 5- to 7-membered cyclic amino group,
- (16) an acylamino group which is shown by  
(i) -NHCOOR<sup>3</sup>, (ii) -NHCONHR<sup>3</sup>, (iii) -NHCOR<sup>3</sup> or (iv) -NHSO<sub>2</sub>R<sup>3</sup> wherein R<sup>3</sup> is (1) a C<sub>1-6</sub> alkyl group, (2) a C<sub>2-6</sub> alkenyl group, (3) a C<sub>2-6</sub> alkynyl group, (4) a C<sub>3-6</sub> cycloalkyl group which may be fused with a benzene ring optionally having 1 to 3 C<sub>1-6</sub> alkoxy groups, (5) a C<sub>6-10</sub> aryl group or (6) a C<sub>7-16</sub> aralkyl group, each of a group shown by the above items (1) to (6) optionally having 1 to 5 substituents selected from the group consisting of

(a) a halogen atom, (b) a C<sub>1-3</sub> alkylenedioxy group, (c) a nitro group, (d) a cyano group, (e) a C<sub>1-6</sub> alkyl group optionally having 1 to 3 halogen atoms, (f) a C<sub>3-6</sub> cycloalkyl group, (g) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (h) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (i) a hydroxyl group, (j) an amino group, (k) a mono-C<sub>1-6</sub> alkylamino group, (l) a di-C<sub>1-6</sub> alkylamino group, (m) a C<sub>1-6</sub> alkyl-carbonyl group, (n) a carboxyl group, (o) a C<sub>1-6</sub> alkoxy-carbonyl group, (p) a carbamoyl group, (q) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (r) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (s) a C<sub>6-10</sub> aryl-carbamoyl group, (t) a sulfo group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, (w) a C<sub>6-10</sub> aryloxy group and (x) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,

- (17) a C<sub>1-6</sub> alkyl-carbonyl group,
- (18) a carboxyl group,
- (19) a C<sub>1-6</sub> alkoxy-carbonyl group,
- (20) a carbamoyl group,
- (21) a mono-C<sub>1-6</sub> alkyl-carbamoyl group,
- (22) a di-C<sub>1-6</sub> alkyl-carbamoyl group,
- (23) a C<sub>6-10</sub> aryl-carbamoyl group,
- (24) a sulfo group,
- (25) a C<sub>1-6</sub> alkylsulfonyl group,
- (26) a C<sub>6-10</sub> aryl group, and
- (27) a C<sub>6-10</sub> aryloxy group;

Q<sup>1</sup> and Q<sup>2</sup> independently represent

- (1) a C<sub>1-6</sub> alkylene group,
- (2) a C<sub>2-6</sub> alkenylene group, or
- (3) a C<sub>2-6</sub> alkynylene group, each of a group shown by the above items (1) to (3) may have oxygen or

optionally oxydized sulfur within the carbon chain;

$R^1$  is

- (1) a hydrogen atom,
- (2) a  $C_{1-6}$  alkyl group which may have 1 to 5 substituents selected from the group consisting of (a) a halogen atom, (b) a  $C_{1-3}$  alkylenedioxy group, (c) a nitro group, (d) a cyano group, (e) a  $C_{1-6}$  alkyl group optionally having 1 to 3 halogen atoms, (f) a  $C_{3-6}$  cycloalkyl group, (g) a  $C_{1-6}$  alkoxy group optionally having 1 to 3 halogen atoms, (h) a  $C_{1-6}$  alkylthio group optionally having 1 to 3 halogen atoms, (i) a hydroxyl group, (j) an amino group, (k) a mono- $C_{1-6}$  alkylamino group, (l) a di- $C_{1-6}$  alkylamino group, (m) a  $C_{1-6}$  alkyl-carbonyl group, (n) a carboxyl group, (o) a  $C_{1-6}$  alkoxy-carbonyl group, (p) a carbamoyl group, (q) a mono- $C_{1-6}$  alkyl-carbamoyl group, (r) a di- $C_{1-6}$  alkyl-carbamoyl group, (s) a  $C_{6-10}$  aryl-carbamoyl group, (t) a sulfo group, (u) a  $C_{1-6}$  alkylsulfonyl group, (v) a  $C_{6-10}$  aryl group, (w) a  $C_{6-10}$  aryloxy group and (x) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, or
- (3) a  $C_{1-6}$  alkyl-carbonyl group which may have 1 to 5 substituents selected from (a) a halogen atom, (b) a  $C_{1-3}$  alkylenedioxy group, (c) a nitro group, (d) a cyano group, (e) a  $C_{1-6}$  alkyl group optionally having 1 to 3 halogen atoms, (f) a  $C_{3-6}$  cycloalkyl group, (g) a  $C_{1-6}$  alkoxy group optionally having 1 to 3 halogen atoms, (h) a  $C_{1-6}$  alkylthio group optionally having 1 to 3 halogen atoms, (i) a hydroxyl group, (j) an amino group, (k) a mono- $C_{1-6}$  alkylamino group, (l) a di- $C_{1-6}$  alkylamino group, (m) a  $C_{1-6}$  alkyl-carbonyl group, (n) a carboxyl group, (o) a  $C_{1-6}$  alkoxy-carbonyl group, (p) a

carbamoyl group, (q) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (r) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (s) a C<sub>6-10</sub> aryl-carbamoyl group, (t) a sulfo group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, (w) a C<sub>6-10</sub> aryloxy group and (x) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring;

R<sup>2</sup> is

- (1) a C<sub>1-6</sub> alkyl group,
- (2) a C<sub>2-6</sub> alkenyl group,
- (3) a C<sub>2-6</sub> alkynyl group,
- (4) a C<sub>3-6</sub> cycloalkyl group which may be fused with a benzene ring optionally having 1 to 3 C<sub>1-6</sub> alkoxy groups,

- (5) a C<sub>6-10</sub> aryl group,
- (6) a C<sub>7-16</sub> aralkyl group,

each of a group shown by the above items (1) to (6) optionally having 1 to 5 substituents selected from the group consisting of (a) a halogen atom, (b) a C<sub>1-3</sub> alkylenedioxy group, (c) a nitro group, (d) a cyano group, (e) a C<sub>1-6</sub> alkyl group optionally having 1 to 3 halogen atoms, (f) a C<sub>3-6</sub> cycloalkyl group, (g) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (h) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (i) a hydroxyl group, (j) an amino group, (k) a mono-C<sub>1-6</sub> alkylamino group, (l) a di-C<sub>1-6</sub> alkylamino group, (m) a C<sub>1-6</sub> alkyl-carbonyl group, (n) a carboxyl group, (o) a C<sub>1-6</sub> alkoxy-carbonyl group, (p) a carbamoyl group, (q) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (r) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (s) a C<sub>6-10</sub> aryl-carbamoyl group, (t) a sulfo group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, (w) a C<sub>6-10</sub>

aryloxy group and (x) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, or

(7) an acyl group which is shown by the formula:

$-(C=O)-R^4$ ,  $-SO_2-R^4$ ,  $-(C=O)NR^5R^4$ ,  $-(C=O)O-R^4$ ,  $-(C=S)O-R^4$ ,  
or  $-(C=S)NR^5R^4$ , wherein  $R^4$  is

(i) a hydrogen atom,

(ii) a  $C_{1-6}$  alkyl group,

(iii) a  $C_{2-6}$  alkenyl group,

(iv) a  $C_{2-6}$  alkynyl group,

(v) a  $C_{3-6}$  cycloalkyl group which may be fused with a benzene ring optionally having 1 to 3  $C_{1-6}$  alkoxy groups,

(vi) a  $C_{6-10}$  aryl group,

(vii) a  $C_{7-16}$  aralkyl group,

(viii) a 5- to 11-membered heterocyclic group having at least one hetero atom selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,

(ix) a  $C_{1-6}$  alkyl-carbonyl group,

(x) a carboxyl group,

(xi) a  $C_{1-6}$  alkoxy-carbonyl group,

(xii) a mono- $C_{1-6}$  alkyl-carbamoyl group,

(xiii) a di- $C_{1-6}$  alkyl-carbamoyl group,

(xiv) a 5- to 7-membered cyclic amino group, or

(xv) a  $C_{6-10}$  aryloxy group,

each of a group shown by the above items (ii) to (xv) optionally having 1 to 5 substituents selected from the group consisting of (a) a halogen atom, (b) a  $C_{1-3}$  alkylenedioxy group, (c) a nitro group, (d) a cyano group, (e) a  $C_{1-6}$  alkyl group optionally substituted with (e-1) a halogen atom, (e-2) a  $C_{1-3}$  alkylenedioxy



group, (e-3) a nitro group, (e-4) a cyano group, (e-5) a C<sub>3-6</sub> cycloalkyl group, (e-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (e-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (e-8) a hydroxyl group, (e-9) an amino group, (e-10) a mono-C<sub>1-6</sub> alkylamino group, (e-11) a di-C<sub>1-6</sub> alkylamino group, (e-12) a C<sub>1-6</sub> alkyl-carbonyl group, (e-13) a carboxyl group, (e-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (e-15) a carbamoyl group, (e-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (e-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (e-18) a C<sub>6-10</sub> aryl-carbamoyl group, (e-19) a sulfo group, (e-20) a C<sub>1-6</sub> alkylsulfonyl group, (e-21) a C<sub>6-10</sub> aryl group, (e-22) a C<sub>6-10</sub> aryloxy group or (e-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (f) a C<sub>3-6</sub> cycloalkyl group, (g) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (h) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (i) a C<sub>7-16</sub> aralkyl group, (j) a hydroxyl group, (k) an amino group which may be substituted with a C<sub>1-6</sub> alkyl carbonyl group, (l) a mono-C<sub>1-6</sub> alkylamino group, (m) a di-C<sub>1-6</sub> alkylamino group, (n) a C<sub>1-6</sub> alkyl-carbonyl group whose alkyl portion may be substituted with (n-1) a halogen atom, (n-2) a C<sub>1-3</sub> alkylenedioxy group, (n-3) a nitro group, (n-4) a cyano group, (n-5) a C<sub>3-6</sub> cycloalkyl group, (n-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (n-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (n-8) a hydroxyl group, (n-9) an amino group, (n-10) a mono-C<sub>1-6</sub> alkylamino group, (n-11) a di-C<sub>1-6</sub> alkylamino group, (n-12) a C<sub>1-6</sub> alkyl-carbonyl group, (n-13) a carboxyl group, (n-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (n-15) a carbamoyl group,


(n-16) a mono- $C_{1-6}$  alkyl-carbamoyl group, (n-17) a di- $C_{1-6}$  alkyl-carbamoyl group, (n-18) a  $C_{6-10}$  aryl-carbamoyl group, (n-19) a sulfo group, (n-20) a  $C_{1-6}$  alkylsulfonyl group, (n-21) a  $C_{6-10}$  aryl group, (n-22) a  $C_{6-10}$  aryloxy group or (n-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (o) a carboxyl group, (p) a  $C_{1-6}$  alkoxy-carbonyl group, (q) a formyl group which may be substituted with 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (r) a carbamoyl group, (s) a mono- $C_{1-6}$  alkyl-carbamoyl group whose alkyl portion may be substituted with (s-1) a halogen atom, (s-2) a  $C_{1-3}$  alkylenedioxy group, (s-3) a nitro group, (s-4) a cyano group, (s-5) a  $C_{3-6}$  cycloalkyl group, (s-6) a  $C_{1-6}$  alkoxy group optionally having 1 to 3 halogen atoms, (s-7) a  $C_{1-6}$  alkylthio group optionally having 1 to 3 halogen atoms, (s-8) a hydroxyl group, (s-9) an amino group, (s-10) a mono- $C_{1-6}$  alkylamino group, (s-11) a di- $C_{1-6}$  alkylamino group, (s-12) a  $C_{1-6}$  alkyl-carbonyl group, (s-13) a carboxyl group, (s-14) a  $C_{1-6}$  alkoxy-carbonyl group, (s-15) a carbamoyl group, (s-16) a mono- $C_{1-6}$  alkyl-carbamoyl group, (s-17) a di- $C_{1-6}$  alkyl-carbamoyl group, (s-18) a  $C_{6-10}$  aryl-carbamoyl group, (s-19) a sulfo group, (s-20) a  $C_{1-6}$  alkylsulfonyl group, (s-21) a  $C_{6-10}$  aryl group, (s-22) a  $C_{6-10}$  aryloxy group or (s-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (t) a di- $C_{1-6}$  alkyl-carbamoyl group whose alkyl portion may be

substituted with (t-1) a halogen atom, (t-2) a C<sub>1-3</sub> alkylenedioxy group, (t-3) a nitro group, (t-4) a cyano group, (t-5) a C<sub>3-6</sub> cycloalkyl group, (t-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (t-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (t-8) a hydroxyl group, (t-9) an amino group, (t-10) a mono-C<sub>1-6</sub> alkylamino group, (t-11) a di-C<sub>1-6</sub> alkylamino group, (t-12) a C<sub>1-6</sub> alkyl-carbonyl group, (t-13) a carboxyl group, (t-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (t-15) a carbamoyl group, (t-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (t-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (t-18) a C<sub>6-10</sub> aryl-carbamoyl group, (t-19) a sulfo group, (t-20) a C<sub>1-6</sub> alkylsulfonyl group, (t-21) a C<sub>6-10</sub> aryl group, (t-22) a C<sub>6-10</sub> aryloxy group or (t-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (u) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (v) an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group, (w) a sulfo group which may be substituted with an amino group, (x) a C<sub>1-6</sub> alkylsulfonyl group, (y) a C<sub>6-10</sub> aryl group, (z) a C<sub>6-10</sub> aryloxy group, (aa) a C<sub>2-6</sub> alkenylamino group, (bb) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (cc) a 5- to 7-membered cyclic amino group which may have an oxo group or which may be substituted with a hydroxyl group, (dd) a C<sub>1-6</sub> alkoxy-carbamoyl group, (ee) a carbamoyloxy group, (ff) a sulfamoyl group, (gg) a mono-C<sub>1-6</sub> alkyl-sulfamoyl group, and (hh) a di-C<sub>1-6</sub> alkyl-sulfamoyl group;

R<sup>5</sup> is

1) a hydrogen atom or

2) a C<sub>1-6</sub> alkyl group;  
 or R<sup>1</sup> and R<sup>2</sup>, taken together with the adjacent nitrogen atom, form a 4- to 8-membered heterocyclic group optionally having at least one nitrogen and 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, which may have 1 to 5 substituents selected from the group consisting of (a) a halogen atom, (b) a C<sub>1-3</sub> alkylenedioxy group, (c) a nitro group, (d) a cyano group, (e) a C<sub>1-6</sub> alkyl group optionally having 1 to 3 halogen atoms, (f) a C<sub>3-6</sub> cycloalkyl group, (g) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (h) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (i) a hydroxyl group, (j) an amino group, (k) a mono-C<sub>1-6</sub> alkylamino group, (l) a di-C<sub>1-6</sub> alkylamino group, (m) a C<sub>1-6</sub> alkyl-carbonyl group, (n) a carboxyl group, (o) a C<sub>1-6</sub> alkoxy-carbonyl group, (p) a carbamoyl group, (q) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (r) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (s) a C<sub>6-10</sub> aryl-carbamoyl group, (t) a sulfo group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, and (w) a C<sub>6-10</sub> aryloxy group;  
 a group of the formula:



is (1) a 4- to 9-membered monocyclic ring or (2) 6- to 14-membered bicyclic ring, each of which may have 1 or 2 unsaturated bonds and optionally having 1 or 2 substituents selected from the group consisting of  
 (i) a C<sub>1-6</sub> alkyl group,  
 (ii) a C<sub>1-6</sub> alkoxy group,  
 (iii) a C<sub>1-6</sub> alkylthio group, each of a group shown by the above items (i) to (iii) may have 1 to 5


substituents selected from (a) a halogen atom, (b) a C<sub>1-3</sub> alkylenedioxy group, (c) a nitro group, (d) a cyano group, (e) a C<sub>1-6</sub> alkyl group optionally having 1 to 3 halogen atoms, (f) a C<sub>3-6</sub> cycloalkyl group, (g) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (h) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (i) a hydroxyl group, (j) an amino group, (k) a mono-C<sub>1-6</sub> alkylamino group, (l) a di-C<sub>1-6</sub> alkylamino group, (m) a C<sub>1-6</sub> alkyl-carbonyl group, (n) a carboxyl group, (o) a C<sub>1-6</sub> alkyl-carbamoyl group, (p) a carbamoyl group, (q) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (r) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (s) a C<sub>6-10</sub> aryl-carbamoyl group, (t) a sulfo group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, (w) a C<sub>6-10</sub> aryloxy group and (x) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,

(iv) a hydroxyl group,  
(v) an amino group,  
(vi) a mono-C<sub>1-6</sub> alkylamino group,  
(vii) a di-C<sub>1-6</sub> alkylamino group,  
(viii) a C<sub>1-6</sub> alkyl-carbonyl group,  
(ix) a carboxyl group,  
(x) a C<sub>1-6</sub> alkoxy-carbonyl group,  
(xi) a carbamoyl group,  
(xii) a mono-C<sub>1-6</sub> alkyl-carbamoyl group,  
(xiii) a di-C<sub>1-6</sub> alkyl-carbamoyl group,  
(xiv) a C<sub>6-10</sub> aryl-carbamoyl group,  
(xv) a sulfo group,  
(xvi) a C<sub>1-6</sub> alkylsulfonyl group,  
(xv) a C<sub>6-10</sub> aryl group, and  
(xvi) a C<sub>6-10</sub> aryloxy group.

3. A composition as claimed in Claim 1 wherein  $R^1$  is a hydrogen atom or a  $C_{1-6}$  alkyl group.
4. A composition as claimed in Claim 1 wherein  $R^1$  is a hydrogen atom or methyl.
5. A composition as claimed in Claim 1 wherein  $R^1$  is a hydrogen atom.
6. A composition as claimed in Claim 1 wherein  $R^2$  is an acyl group.
7. A composition as claimed in Claim 6 wherein the acyl group is of the formula  $-(C=O)-R^4$ ,  $-SO_2-R^4$ ,  $-SO-R^4$ ,  $-(C=O)NR^5R^4$ ,  $-(C=O)O-R^4$ ,  $-(C=S)O-R^4$ , or  $-(C=S)NR^5R^4$ , wherein  $R^4$  is a hydrogen atom, an optionally substituted hydrocarbon group, an optionally substituted heterocyclic group, an optionally substituted lower alkyl-carbonyl group, a carboxyl group, an optionally substituted lower alkoxy-carbonyl group, an optionally substituted mono-lower alkylaminocarbonyl group, an optionally substituted di-lower alkylaminocarbonyl group, an optionally substituted 5- or 7-membered cyclic amino group or an optionally substituted aryloxy group; and  $R^5$  is a hydrogen atom or a lower alkyl group.
8. A composition as claimed in Claim 6, wherein the acyl group is of the formula  $-(C=O)-R^4$  or  $-(C=O)NHR^4$ , wherein  $R^4$  is a hydrogen atom, an optionally substituted hydrocarbon group, an optionally substituted heterocyclic group, an optionally substituted lower alkyl-carbonyl group, a carboxyl group, an optionally substituted lower alkoxy-carbonyl group, an optionally substituted mono-lower alkylaminocarbonyl group, an optionally substituted di-lower alkylaminocarbonyl group, an optionally substituted 5- or 7-membered cyclic amino group or an optionally substituted aryloxy group; and  $R^5$  is a hydrogen atom or a lower alkyl group.


9. A composition as claimed in Claim 8, wherein  $R^4$  is a group of the formula:

(1)



or

(2)



wherein  $R^6$  and  $R^7$  independently represent (a) a hydrogen atom, (b) a  $C_{1-6}$  alkyl group optionally substituted with

(b-1) a halogen atom, (b-2) a  $C_{1-3}$  alkylenedioxy group, (b-3) a nitro group, (b-4) a cyano group, (b-5) a  $C_{3-6}$  cycloalkyl group, (b-6) a  $C_{1-6}$  alkoxy group optionally having 1 to 3 halogen atoms, (b-7) a  $C_{1-6}$  alkylthio group optionally having 1 to 3 halogen atoms, (b-8) a hydroxyl group, (b-9) an amino group, (b-10) a mono- $C_{1-6}$  alkylamino group, (b-11) a di- $C_{1-6}$  alkylamino group, (b-12) a  $C_{1-6}$  alkyl-carbonyl group, (b-13) a carboxyl group, (b-14) a  $C_{1-6}$  alkoxy-carbonyl group, (b-15) a carbamoyl group, (b-16) a mono- $C_{1-6}$  alkyl-carbamoyl group, (b-17) a di- $C_{1-6}$  alkyl-carbamoyl group, (b-18) a  $C_{6-10}$  aryl-carbamoyl group, (b-19) a sulfo group, (b-20) a  $C_{1-6}$  alkylsulfonyl group, (b-21) a  $C_{6-10}$  aryl group, (b-22) a  $C_{6-10}$  aryloxy group or (b-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (c) a  $C_{3-6}$  cycloalkyl group, (d) a  $C_{1-6}$  alkoxy group optionally having 1 to 3 halogen atoms, (e) a  $C_{1-6}$  alkylthio group optionally having 1 to 3 halogen atoms, (f) a  $C_{7-16}$  aralkyl group, (g) a hydroxyl group, (h) an amino

group, (i) a mono- $C_{1-6}$  alkylamino group, (j) a di- $C_{1-6}$  alkylamino group, (k) a  $C_{1-6}$  alkyl-carbonyl group whose alkyl portion may be substituted with (k-1) a halogen atom, (k-2) a  $C_{1-3}$  alkylenedioxy group, (k-3) a nitro group, (k-4) a cyano group, (k-5) a  $C_{3-6}$  cycloalkyl group, (k-6) a  $C_{1-6}$  alkoxy group optionally having 1 to 3 halogen atoms, (k-7) a  $C_{1-6}$  alkylthio group optionally having 1 to 3 halogen atoms, (k-8) a hydroxyl group, (k-9) an amino group, (k-10) a mono- $C_{1-6}$  alkylamino group, (k-11) a di- $C_{1-6}$  alkylamino group, (k-12) a  $C_{1-6}$  alkyl-carbonyl group, (k-13) a carboxyl group, (k-14) a  $C_{1-6}$  alkoxy-carbonyl group, (k-15) a carbamoyl group, (k-16) a mono- $C_{1-6}$  alkyl-carbamoyl group, (k-17) a di- $C_{1-6}$  alkyl-carbamoyl group, (k-18) a  $C_{6-10}$  aryl-carbamoyl group, (k-19) a sulfo group, (k-20) a  $C_{1-6}$  alkylsulfonyl group, or (k-21) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (l) a carboxyl group, (m) a  $C_{1-6}$  alkoxy-carbonyl group, (n) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (o) a carbamoyl group, (p) a mono- $C_{1-6}$  alkyl-carbamoyl group whose alkyl portion may be substituted with (p-1) a halogen atom, (p-2) a  $C_{1-3}$  alkylenedioxy group, (p-3) a nitro group, (p-4) a cyano group, (p-5) a  $C_{3-6}$  cycloalkyl group, (p-6) a  $C_{1-6}$  alkoxy group optionally having 1 to 3 halogen atoms, (p-7) a  $C_{1-6}$  alkylthio group optionally having 1 to 3 halogen atoms, (p-8) a hydroxyl group, (p-9) an amino group, (p-10) a mono- $C_{1-6}$  alkylamino group, (p-11) a di- $C_{1-6}$  alkylamino group, (p-12) a  $C_{1-6}$  alkyl-carbonyl group,




(p-13) a carboxyl group, (p-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (p-15) a carbamoyl group, (p-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (p-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (p-18) a C<sub>6-10</sub> aryl-carbamoyl group, (p-19) a sulfo group, (p-20) a C<sub>1-6</sub> alkylsulfonyl group, (p-21) a C<sub>6-10</sub> aryl group, (p-22) a C<sub>6-10</sub> aryloxy group or (p-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (q) a di-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (q-1) a halogen atom, (q-2) a C<sub>1-3</sub> alkylenedioxy group, (q-3) a nitro group, (q-4) a cyano group, (q-5) a C<sub>3-6</sub> cycloalkyl group, (q-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (q-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (q-8) a hydroxyl group, (q-9) an amino group, (q-10) a mono-C<sub>1-6</sub> alkylamino group, (q-11) a di-C<sub>1-6</sub> alkylamino group, (q-12) a C<sub>1-6</sub> alkyl-carbonyl group, (q-13) a carboxyl group, (q-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (q-15) a carbamoyl group, (q-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (q-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (q-18) a C<sub>6-10</sub> aryl-carbamoyl group, (q-19) a sulfo group, (q-20) a C<sub>1-6</sub> alkylsulfonyl group, (q-21) a C<sub>6-10</sub> aryl group, (q-22) a C<sub>6-10</sub> aryloxy group or (q-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (r) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (s) an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group, (t) a sulfo group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, (w) a C<sub>6-10</sub> aryloxy group, (x) a C<sub>2-6</sub> alkenylamino group or (y) a 5- to 7-membered

heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring.


10. A composition as claimed in Claim 8, wherein  $R^4$  is a group of the formula:

(1)



or

(2)



wherein  $R^6$  and  $R^7$  independently represent (a) a hydrogen atom, (b) a  $C_{1-6}$  alkyl group optionally substituted with (b-1) a hydroxyl group, (b-2) a di- $C_{1-6}$  alkylamino group, (b-3) a  $C_{1-6}$  alkoxy-carbonyl group, or (b-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (c) a  $C_{7-16}$  aralkyl group, (d) a  $C_{1-6}$  alkyl-carbonyl group whose alkyl portion may be substituted with (d-1) a halogen atom, (d-2) a mono- $C_{1-6}$  alkylamino group, (d-3) a  $C_{1-6}$  alkoxy-carbonyl group, or (d-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (e) a  $C_{1-6}$  alkoxy-carbonyl group, (f) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being

optionally fused with benzene ring, (g) a mono- $C_{1-6}$  alkyl-carbamoyl group whose alkyl portion may be substituted with (g-1) a halogen atom, or (g-2) a  $C_{1-6}$  alkyl-carbonyl group, (h) an optionally halogenated  $C_{6-10}$  aryl-carbamoyl group, (i) an optionally halogenated  $C_{6-10}$  aryl-carbonyl group, or (j) a  $C_{6-10}$  aryloxy group.

11. A composition as claimed in Claim 1 wherein  $Q^1$  and  $Q^2$  are independently a  $C_{1-6}$  alkylene group which may have an oxo group.

12. A composition as claimed in Claim 1 wherein  $Q^1$  is a  $C_{1-4}$  alkylene group and  $Q^2$  is a methylene group.

13. A composition as claimed in Claim 1 wherein the ring of the formula:




is a 4- to 9-membered monocyclic ring or 6- to 14-membered bicyclic ring, which may have 1 or 2 unsaturated bonds and may have 1 or 2 substituents in any position other than N and Z.

14. A composition as claimed in Claim 1 wherein the ring of the formula:



is




15. A composition as claimed in Claim 1 wherein the

ring of the formula:



is




16. A composition as claimed in Claim 1 wherein the ring of the formula:



is




17. A composition as claimed in Claim 13 wherein Z is  
 (1) an optionally substituted 1, 2-phenylene,  
 (2) a group of the formula:



wherein  $Ar^3$  is an optionally substituted aromatic group, and n is an integer of 0 to 3,


(3) a group of the formula:



wherein  $Ar^3$  and n have the same meanings as defined above; and Y is (i) a hydrogen atom, (ii) an optionally halogenated lower alkyl group, (iii) an


optionally halogenated lower alkoxy group, (iv) an optionally halogenated lower alkylthio group, (v) a hydroxyl group, (vi) a cyano group, (vii) an alkyl-carbonyl group, (viii) a lower alkyl-carbonyloxy group, (ix) a formylamino group, (x) an amino group, (xi) a mono-lower alkylamino group, (xii) a di-lower alkylamino group, (xiii) a carboxyl group, (xiv) a lower alkoxy-carbonyl group or (xv) a lower alkyl-carbonylamino group, or

(4) a group of the formula:



wherein  $\text{Ar}^3$  and  $n$  have the same meanings as defined above, or

(5) a group of the formula:




wherein  $\text{Ar}^3$  and  $n$  have the same meanings as defined above.

18. A composition as claimed in Claim 1 wherein the ring of the formula:



is pyrrolidine, piperidine, piperazine, azepine or azocine, each of which may be fused with a benzene ring and may have a substituent.

19. A composition as claimed in Claim 13 wherein  $Z$  is a group of the formula:



wherein  $\text{Ar}^3$  is an optionally substituted aromatic group,  $n$  is an integer of 0 to 3, and  $Y$  is a hydrogen atom or a hydroxyl group.

20. A composition as claimed in Claim 19 wherein  $\text{Ar}^3$  is a  $\text{C}_{6-14}$  aryl group or a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms of 1 or 2 kinds selected from nitrogen, oxygen and sulfur in addition to a carbon atom, each of which may have 1 to 3 substituents selected from a halogen atom, an optionally halogenated  $\text{C}_{1-6}$  alkyl group, and an optionally halogenated  $\text{C}_{1-6}$  alkoxy group.

21. A composition as claimed in Claim 19 wherein  $\text{Ar}^3$  is a phenyl group optionally substituted with a halogen atom.

22. A composition as claimed in Claim 19 wherein  $n$  is 0.


23. A composition as claimed in Claim 19 wherein  $Y$  is a hydroxyl group.

24. A composition as claimed in claim 1 wherein  $\text{Ar}^1$  and  $\text{Ar}^2$  independently represent a  $\text{C}_{6-14}$  aryl group or a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms of 1 or 2 kinds selected from nitrogen, oxygen and sulfur in addition to a carbon atom, each of which may have 1 to 3 substituents selected from a halogen atom, an optionally halogenated  $\text{C}_{1-6}$  alkyl group, and an optionally halogenated  $\text{C}_{1-6}$  alkoxy group.


25. A composition as claimed in Claim 1 wherein  $\text{Ar}^1$  and  $\text{Ar}^2$  independently represent phenyl, 4-chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl.

26. A composition as claimed in claim 1, wherein  $\text{Ar}^1$  and  $\text{Ar}^2$  independently represent phenyl, 4-chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl;  $Q^1$  is a  $\text{C}_{1-4}$  alkylene group;  $Q^2$  is a methylene group;


the group of the formula:



is




wherein Z is a group of the formula:



wherein  $Ar^3$  is a phenyl group optionally substituted with a halogen atom,  $n$  is an integer of 0 to 3, and  $Y$  is a hydrogen atom or a hydroxyl group;

$R^1$  is a hydrogen atom or methyl;

$R^2$  is (1) an  $C_{1-6}$  alkyl group which may be substituted with a  $C_{1-6}$  alkoxy-carbonyl group, a carboxyl group, a  $C_{1-6}$  alkyl-carbonyl group or a formyl group or (2) an acyl group represented by the formula:



wherein  $R^4$  is

- (i) a hydrogen atom,
- (ii) a  $C_{1-6}$  alkyl group which may have 1 to 5 substituents selected from (a) a hydroxyl group, (b) an amino group which may be substituted with a  $C_{1-6}$  alkyl-carbonyl group, (c) a mono- $C_{1-6}$  alkylamino group, (d) a di- $C_{1-6}$  alkylamino group, (e) a carboxyl group, (f) a  $C_{1-6}$  alkoxy-carbonyl group, (g) a mono- $C_{1-6}$  alkyl-carbamoyl group, (h) a sulfo group which may be substituted with amino group, (i) a 5- to 7-membered cyclic amino group which may have an oxo group or which may be substituted with a hydroxyl group, (j) a  $C_{1-6}$  alkoxy-carbamoyl group, and (k) a carbamoyloxy group,

- (iii) a C<sub>2-6</sub> alkenyl group,
- (iv) a C<sub>6-10</sub> aryl group,
- (v) a 5- to 11-membered heterocyclic group having at least one hetero atom selected from nitrogen, oxygen and sulfur in addition to a carbon atom, said heterocyclic group being optionally fused with a benzene ring,
- (vi) a C<sub>1-6</sub> alkyl group which may be substituted with a C<sub>1-6</sub> alkyl-carbonyl group,
- (vii) a carboxyl group which may be substituted with a C<sub>1-6</sub> alkyl group,
- (viii) a 5- to 7-membered cyclic amino group which may be substituted with
  - (a) a C<sub>1-6</sub> alkyl group optionally substituted with (a-1) a hydroxyl group, (a-2) a di-C<sub>1-6</sub> alkylamino group, (a-3) a C<sub>1-6</sub> alkoxy-carbonyl group or (a-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to a carbon atom, said heterocyclic group being optionally fused with a benzene ring,
  - (b) a C<sub>7-16</sub> aralkyl group, (c) a C<sub>1-6</sub> alkyl-carbonyl group whose alkyl portion may be substituted with (c-1) a halogen atom, (c-2) a mono-C<sub>1-6</sub> alkylamino group, (c-3) a C<sub>1-6</sub> alkoxy-carbonyl group or (c-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to a carbon atom, said heterocyclic group being optionally fused with a benzene ring,
  - (d) a C<sub>1-6</sub> alkoxy-carbonyl group, (e) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to a carbon atom, said heterocyclic group being optionally fused with a benzene ring,
  - (f) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl




portion may be substituted with a halogen atom or a C<sub>1-6</sub> alkyl-carbonyl group, (g) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (h) an optionally halogenated C<sub>6-10</sub> aryl carbonyl group or (i) a C<sub>1-6</sub> alkoxy-carbamoyl group, or

(ix) a C<sub>6-10</sub> aryloxy group; and

R<sup>5</sup> is a hydrogen atom or a C<sub>1-6</sub> alkyl group.

27. A compound of the formula:



wherein Ar<sup>1</sup>, Ar<sup>2</sup> and Ar<sup>3</sup> independently represent an optionally substituted aromatic group;

Q<sup>1</sup> and Q<sup>2</sup> independently represent a divalent C<sub>1-6</sub> aliphatic hydrocarbon group, which may have oxygen or sulfur within the carbon chain; and

R<sup>2</sup> is an optionally substituted hydrocarbon group or an acyl group or a salt thereof (except N-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino-2,2-diphenylpentyl]-1-methanesulfonamide hydrochloride, N-[5-[4-chlorophenyl)-4-hydroxypiperidino-2,2-diphenylpentyl]-1-(p-toluene)sulfonamide hydrochloride and N-[5-(4-(4-chlorophenyl)-4-hydroxypiperidino-2,2-diphenylpentyl]-1-(2-thiophene)sulfonamide hydrochloride).


28. The compound of Claim 27 wherein R<sup>2</sup> is a group of the formula -(C=O)-R<sup>4</sup>, -(C=O)NR<sup>5</sup>R<sup>4</sup>, -(C=O)O-R<sup>4</sup>, -(C=S)O-R<sup>4</sup> or -(C=S)NR<sup>5</sup>R<sup>4</sup> wherein R<sup>4</sup> is a hydrogen atom, an optionally substituted hydrocarbon group, an optionally substituted heterocyclic group, an optionally substituted lower alkyl-carbonyl group, a carboxyl group, an optionally substituted lower alkoxy carbonyl group, an optionally substituted

mono-lower alkylaminocarbonyl group, an optionally substituted di-lower alkylaminocarbonyl group or an optionally substituted 5- or 7-membered cyclic amino group; and  $R^5$  is a hydrogen atom or a lower alkyl group.

29. A compound as claimed in Claim 27, wherein  $R^2$  is the formula  $-(C=O)-R^4$  or  $-(C=O)NH-R^4$ , wherein  $R^4$  is a hydrogen atom, an optionally substituted hydrocarbon group, an optionally substituted heterocyclic group, an optionally substituted lower alkyl-carbonyl group, a carboxyl group, an optionally substituted lower alkoxycarbonyl group, an optionally substituted mono-lower alkylaminocarbonyl group, an optionally substituted di-lower alkylaminocarbonyl group or an optionally substituted 5- or 7-membered cyclic amino group


30. A compound as claimed in Claim 28, wherein  $R^4$  is of the formula:

(1)



or

(2)



wherein  $R^6$  and  $R^7$  independently represent (a) a hydrogen atom, (b) a  $C_{1-6}$  alkyl group optionally substituted with

(b-1) a halogen atom, (b-2) a  $C_{1-3}$  alkylenedioxy group, (b-3) a nitro group, (b-4) a cyano group, (b-5) a  $C_{3-6}$  cycloalkyl group, (b-6) a  $C_{1-6}$  alkoxy group optionally having 1 to 3 halogen atoms, (b-7) a  $C_{1-6}$  alkylthio group optionally having 1 to 3 halogen atoms, (b-8) a hydroxyl group, (b-9) an amino group, (b-10) a mono- $C_{1-6}$

alkylamino group, (b-11) a di- $C_{1-6}$  alkylamino group, (b-12) a  $C_{1-6}$  alkyl-carbonyl group, (b-13) a carboxyl group, (b-14) a  $C_{1-6}$  alkoxy-carbonyl group, (b-15) a carbamoyl group, (b-16) a mono- $C_{1-6}$  alkyl-carbamoyl group, (b-17) a di- $C_{1-6}$  alkyl-carbamoyl group, (b-18) a  $C_{6-10}$  aryl-carbamoyl group, (b-19) a sulfo group, (b-20) a  $C_{1-6}$  alkylsulfonyl group, (b-21) a  $C_{6-10}$  aryl group, (b-22) a  $C_{6-10}$  aryloxy group or (b-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (c) a  $C_{3-6}$  cycloalkyl group, (d) a  $C_{1-6}$  alkoxy group optionally having 1 to 3 halogen atoms, (e) a  $C_{1-6}$  alkylthio group optionally having 1 to 3 halogen atoms, (f) a  $C_{7-16}$  aralkyl group, (g) a hydroxyl group, (h) an amino group, (i) a mono- $C_{1-6}$  alkylamino group, (j) a di- $C_{1-6}$  alkylamino group, (k) a  $C_{1-6}$  alkyl-carbonyl group whose alkyl portion may be substituted with (k-1) a halogen atom, (k-2) a  $C_{1-3}$  alkylenedioxy group, (k-3) a nitro group, (k-4) a cyano group, (k-5) a  $C_{3-6}$  cycloalkyl group, (k-6) a  $C_{1-6}$  alkoxy group optionally having 1 to 3 halogen atoms, (k-7) a  $C_{1-6}$  alkylthio group optionally having 1 to 3 halogen atoms, (k-8) a hydroxyl group, (k-9) an amino group, (k-10) a mono- $C_{1-6}$  alkylamino group, (k-11) a di- $C_{1-6}$  alkylamino group, (k-12) a  $C_{1-6}$  alkyl-carbonyl group, (k-13) a carboxyl group, (k-14) a  $C_{1-6}$  alkoxy-carbonyl group, (k-15) a carbamoyl group, (k-16) a mono- $C_{1-6}$  alkyl-carbamoyl group, (k-17) a di- $C_{1-6}$  alkyl-carbamoyl group, (k-18) a  $C_{6-10}$  aryl-carbamoyl group, (k-19) a sulfo group, (k-20) a  $C_{1-6}$  alkylsulfonyl group, or (k-21) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said

heterocyclic group being optionally fused with a benzene ring, (l) a carboxyl group, (m) a C<sub>1-6</sub> alkoxy-carbonyl group, (n) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (o) a carbamoyl group, (p) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (p-1) a halogen atom, (p-2) a C<sub>1-3</sub> alkylenedioxy group, (p-3) a nitro group, (p-4) a cyano group, (p-5) a C<sub>3-6</sub> cycloalkyl group, (p-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (p-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (p-8) a hydroxyl group, (p-9) an amino group, (p-10) a mono-C<sub>1-6</sub> alkylamino group, (p-11) a di-C<sub>1-6</sub> alkylamino group, (p-12) a C<sub>1-6</sub> alkyl-carbonyl group, (p-13) a carboxyl group, (p-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (p-15) a carbamoyl group, (p-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (p-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (p-18) a C<sub>6-10</sub> aryl-carbamoyl group, (p-19) a sulfo group, (p-20) a C<sub>1-6</sub> alkylsulfonyl group, (p-21) a C<sub>6-10</sub> aryl group, (p-22) a C<sub>6-10</sub> aryloxy group or (p-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (q) a di-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with (q-1) a halogen atom, (q-2) a C<sub>1-3</sub> alkylenedioxy group, (q-3) a nitro group, (q-4) a cyano group, (q-5) a C<sub>3-6</sub> cycloalkyl group, (q-6) a C<sub>1-6</sub> alkoxy group optionally having 1 to 3 halogen atoms, (q-7) a C<sub>1-6</sub> alkylthio group optionally having 1 to 3 halogen atoms, (q-8) a hydroxyl group, (q-9) an amino group, (q-10) a mono-C<sub>1-6</sub> alkylamino group, (q-11) a di-C<sub>1-6</sub>

alkylamino group, (q-12) a C<sub>1-6</sub> alkyl-carbonyl group, (q-13) a carboxyl group, (q-14) a C<sub>1-6</sub> alkoxy-carbonyl group, (q-15) a carbamoyl group, (q-16) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (q-17) a di-C<sub>1-6</sub> alkyl-carbamoyl group, (q-18) a C<sub>6-10</sub> aryl-carbamoyl group, (q-19) a sulfo group, (q-20) a C<sub>1-6</sub> alkylsulfonyl group, (q-21) a C<sub>6-10</sub> aryl group, (q-22) a C<sub>6-10</sub> aryloxy group or (q-23) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring, (r) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (s) an optionally halogenated C<sub>6-10</sub> aryl-carbonyl group, (t) a sulfo group, (u) a C<sub>1-6</sub> alkylsulfonyl group, (v) a C<sub>6-10</sub> aryl group, (w) a C<sub>6-10</sub> aryloxy group, (x) a C<sub>2-6</sub> alkenylamino group or (y) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring.

31. A compound as claimed in Claim 27 wherein Q<sup>1</sup> and Q<sup>2</sup> are independently a C<sub>1-6</sub> alkylene group which may have an oxo group.

32. A compound as claimed in Claim 27 wherein Q<sup>1</sup> is a C<sub>1-4</sub> alkylene group and Q<sup>2</sup> is a methylene group.

33. A compound as claimed in Claim 27 wherein Ar<sup>3</sup> is a phenyl group optionally substituted with a halogen atom.

34. A compound as claimed in claim 27 wherein Ar<sup>1</sup> and Ar<sup>2</sup> independently represent a C<sub>6-14</sub> aryl group or a 5- to 7-membered heterocyclic groups having 1 to 3 hetero atoms of 1 or 2 kinds selected from nitrogen, oxygen and sulfur in addition to a carbon atom, each of which may have 1 to 3 substituents selected from a halogen atom, an optionally halogenated C<sub>1-6</sub> alkyl group, and an


optionally halogenated C<sub>1-6</sub> alkoxy group.

35. A compound as claimed in Claim 27 wherein Ar<sup>1</sup> and Ar<sup>2</sup> independently represent phenyl, 4-chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl.

36. A compound as claimed in claim 27, wherein Ar<sup>1</sup> and Ar<sup>2</sup> independently represent phenyl, 4-chlorophenyl, 4-fluorophenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl;

Q<sup>1</sup> is a C<sub>1-4</sub> alkylene group; Q<sup>2</sup> is a methylene group;

R<sup>2</sup> is (1) a C<sub>1-6</sub> alkyl group which may be substituted with a C<sub>1-6</sub> alkoxy-carbonyl group, a carboxyl group, a C<sub>1-6</sub> alkyl-carbonyl group or a formyl group or (2) an acyl group represented by the formula:



wherein R<sup>4</sup> is

(i) a hydrogen atom,

(ii) a C<sub>1-6</sub> alkyl group which may have 1 to 5

substituents selected from (a) a hydroxyl group, (b) an amino group which may be substituted with a C<sub>1-6</sub> alkyl-carbonyl group, (c) a mono-C<sub>1-6</sub> alkylamino group, (d) a di-C<sub>1-6</sub> alkylamino group, (e) a carboxyl group, (f) a C<sub>1-6</sub> alkoxy-carbonyl group, (g) a mono-C<sub>1-6</sub> alkyl-carbamoyl group, (h) a sulfo group which may be substituted with amino group (i) a 5- to 7-membered cyclic amino group which may have an oxo group or which may be substituted with a hydroxyl group, (j) a C<sub>1-6</sub> alkoxy-carbamoyl group, and (k) a carbamoyloxy group.

(iii) a C<sub>2-6</sub> alkenyl group,

(iv) a C<sub>6-10</sub> aryl group,

(v) a 5- to 11-membered heterocyclic group having at least one hetero atom selected from nitrogen, oxygen and sulfur in addition to a carbon atom, said heterocyclic group being optionally fused with a benzene ring,

(vi) a C<sub>1-6</sub> alkyl group which may be substituted with a C<sub>1-6</sub> alkyl-carbonyl group,

(vii) a carboxyl group which may be substituted with a C<sub>1-6</sub> alkyl group,

(viii) a 5- to 7-membered cyclic amino group which may be substituted with

(a) a C<sub>1-6</sub> alkyl group optionally substituted with (a-1) a hydroxyl group, (a-2) a di-C<sub>1-6</sub> alkylamino group, (a-3) a C<sub>1-6</sub> alkoxy-carbonyl group or (a-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,

(b) a C<sub>7-16</sub> aralkyl group, (c) a C<sub>1-6</sub> alkyl-carbonyl group whose alkyl portion may be substituted with (c-1) a halogen atom, (c-2) a mono-C<sub>1-6</sub> alkylamino group, (c-3) a C<sub>1-6</sub> alkoxy-carbonyl group or (c-4) a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,

(d) a C<sub>1-6</sub> alkoxy-carbonyl group, (e) a formyl group which may be substituted with a 5- to 7-membered heterocyclic group having 1 to 3 hetero atoms selected from nitrogen, oxygen and sulfur in addition to carbon atoms, said heterocyclic group being optionally fused with a benzene ring,


(f) a mono-C<sub>1-6</sub> alkyl-carbamoyl group whose alkyl portion may be substituted with a halogen atom or a C<sub>1-6</sub> alkyl-carbonyl group, (g) an optionally halogenated C<sub>6-10</sub> aryl-carbamoyl group, (h) an optionally halogenated C<sub>6-10</sub> aryl carbonyl group or (i) a C<sub>1-6</sub> alkoxy-carbamoyl group, or

(ix) a C<sub>6-10</sub> aryloxy group;


R<sup>5</sup> is a hydrogen atom or a C<sub>1-6</sub> alkyl group; and

Ar<sup>3</sup> is a phenyl group optionally substituted with a halogen atom.

37. A process for producing a compound of the formula:




wherein  $R^2$  is an acyl group, and the other symbols have the same meanings as defined in Claim 27 or a salt thereof, which comprises subjecting a compound of the formula:



wherein the all symbols have the same meanings as defined in Claim 27 or a salt thereof to the acylation reaction.


38. A process for producing a compound of the formula:




wherein  $R^4$  represents a hydrogen atom, an optionally substituted hydrocarbon group, an optionally substituted heterocyclic group, an optionally substituted lower alkyl-carbonyl group, a carboxyl group, an optionally substituted lower alkoxy-carbonyl group, an optionally substituted mono-lower alkylaminocarbonyl group, an optionally substituted di-lower alkylaminocarbonyl group or an optionally



substituted 5- or 6-membered cyclic amino group; and R<sup>5</sup> is a hydrogen atom or a lower alkyl group, and the other symbols have the same meanings as defined in Claim 27 or a salt thereof, which comprises reacting a compound of the formula:



wherein Ph is a phenyl group, and the other symbols have the same meanings as defined above or a salt thereof with a compound of the formula:



wherein R<sup>4</sup> and R<sup>5</sup> have the same meanings as defined above or a salt thereof.

39. A composition as claimed in Claim 1 which is a prophylactic or therapeutic agent for inflammatory diseases.

40. A composition as claimed in Claim 1 which is a prophylactic or therapeutic agent for allergic diseases.


41. A composition as claimed in Claim 1 which is a prophylactic or therapeutic agent for arteriosclerosis, bronchial asthma, atopy, multiple sclerosis or rheumatoid arthritis.

42. A pharmaceutical composition comprising the compound of Claim 27.

43. A MIP-1 $\alpha$ /RANTES receptor antagonist comprising the compound of claim 27.

44. A method of treating or preventing inflammatory diseases or allergic diseases which comprises

administering to a mammal in need an effective amount of a compound of the formula:




wherein  $\text{Ar}^1$  and  $\text{Ar}^2$  independently represent an optionally substituted aromatic group;

$\text{Q}^1$  and  $\text{Q}^2$  independently represent an optionally substituted divalent  $\text{C}_{1-6}$  aliphatic hydrocarbon group which may have oxygen or sulfur within the carbon chain;


$\text{R}^1$  is a hydrogen atom, an optionally substituted lower alkyl group or an optionally substituted lower alkyl-carbonyl group;

$\text{R}^2$  is an optionally substituted hydrocarbon group or an acyl group, or  $\text{R}^1$  and  $\text{R}^2$ , taken together with the adjacent nitrogen atom, form an optionally substituted nitrogen-containing heterocyclic ring; and a group of the formula:



is an optionally substituted monocyclic or fused nitrogen-containing heterocyclic ring, or a salt thereof.

45. Use of a compound of the formula:




wherein  $\text{Ar}^1$  and  $\text{Ar}^2$  independently represent an optionally substituted aromatic group;

$Q^1$  and  $Q^2$  independently represent an optionally substituted divalent  $C_{1-6}$  aliphatic hydrocarbon group which may have oxygen or sulfur within the carbon chain;

$R^1$  is a hydrogen atom, an optionally substituted lower alkyl group or an optionally substituted lower alkyl-carbonyl group;

$R^2$  is an optionally substituted hydrocarbon group or an acyl group, or  $R^1$  and  $R^2$ , taken together with the adjacent nitrogen atom, form an optionally substituted nitrogen-containing heterocyclic ring; and a group of the formula:



is an optionally substituted monocyclic or fused nitrogen-containing heterocyclic ring or a salt thereof, for the manufacture of a medicament for treating or preventing inflammatory diseases or allergic diseases.

46. A compound as claimed in claim 27, which is  
 1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]-3-(piperidin-4-yl)urea,  
 Ethyl 4-[4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentylaminocarbonylamino]piperidino-4-oxobutyrate,  
 N-Ethyl-4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino-1-piperidinecarboxamide,  
 N-Ethoxycarbonylmethyl-4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino-1-piperidinecarboxamide,  
 Ethyl 3-[4-[5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpentyl]aminocarbonylamino]piperidino-3-oxopropionate,  
 1-[5-[4-(4-Chlorophenyl)-4-hydroxypiperidino]-

2,2-diphenylpentyl]-3-(1-ethylpiperidin-4-yl)urea,  
1-[(Piperidin-4-yl)carboxamido]-5-[4-(4-chlorophenyl)-4-  
-hydroxypiperidino]-2,2-diphenylpentane,  
1-[(N-Ethylcarbamoylethyl)piperidin-4-yl]carboxamido]-5-[4-  
(4-chlorophenyl)-4-hydroxypiperidino]-2,2-diphenylpenta  
ne,  
1-[[N-(Ethoxycarbonylacetyl)piperidin-4-yl]carboxamido]  
-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-dipheny  
lpentane,  
1-[[N-(3-Methoxycarbonylpropionyl)piperidin-4-yl]carbox  
amido]-5-[4-(4-chlorophenyl)-4-hydroxypiperidino]-2,2-  
diphenylpentane, or a salt thereof.

# INTERNATIONAL SEARCH REPORT

Int'l Application No  
PCT/JP 96/03820

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D211/52 C07D295/12 C07D211/70 C07D401/04 C07D401/06  
C07D401/12 C07D223/08 C07D409/06 C07D211/58 A61K31/445

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	EP 0 712 845 A (TAKEDA) 22 May 1996 see the whole document ---	1-46
X	EP 0 219 898 A (JANSSEN) 29 April 1987 see the whole document ---	1-46
X	FR 2 100 711 A (JANSSEN) 24 March 1972 see the whole document ---	1-46
X	FR 2 408 599 A (DEVINTER SA) 8 June 1979 see the whole document ---	1-46
A	WO 94 11504 A (GENENTECH INC) 26 May 1994 see the whole document ---	1-46
A	WO 94 07521 A (UNIVERSITY OF TEXAS) 14 April 1994 see the whole document -----	1-46

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

4 April 1997

Date of mailing of the international search report

14. 04. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Kissler, B

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 96/03820

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 712845 A	22-05-96	JP 8253447 A	01-10-96
EP 219898 A	29-04-87	AU 583429 B	27-04-89
		AU 6381586 A	16-04-87
		CA 1332236 A	04-10-94
		CY 1603 A	03-04-92
		EG 18126 A	30-08-92
		HK 52491 A	19-07-91
		IE 59519 B	09-03-94
		JP 1810892 C	27-12-93
		JP 5017909 B	10-03-93
		JP 62087569 A	22-04-87
		KR 9401772 B	05-03-94
		SU 1443798 A	07-12-88
		US 4898873 A	06-02-90
		US 4824853 A	25-04-89
FR 2100711 A	24-03-72	AT 312606 A,B	15-12-73
		BE 767798 A	29-11-71
		CH 549015 A	15-05-74
		CH 553182 A	30-08-74
		DE 2126559 A	05-10-72
		DE 2167193 C	19-05-83
		GB 1319040 A	31-05-73
		HK 78076 A	17-12-76
		JP 1242513 C	26-11-84
		JP 55120576 A	17-09-80
		JP 59014467 B	04-04-84
		JP 1258821 C	12-04-85
		JP 55120577 A	17-09-80
		JP 59036991 B	06-09-84
		JP 1182918 C	27-12-83
		JP 56127355 A	06-10-81
		JP 58017749 B	09-04-83
		NL 7106829 A,C	03-12-71
		SE 369418 B	26-08-74
		ZA 7103471 A	31-01-73
		US 3714159 A	30-01-73
		US 3884916 A	20-05-75

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/JP 96/03820

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
FR 2408599 A	08-06-79	NONE	
-----	-----	-----	-----
WO 9411504 A	26-05-94	EP 0669979 A	06-09-95
		JP 8503463 T	16-04-96
-----	-----	-----	-----
WO 9407521 A	14-04-94	AU 5293293 A	26-04-94
-----	-----	-----	-----